

**ANNEX II**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Carton 50 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Revozyn RTU 400 mg/ml suspension for injection

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each ml contains:  
308.8 mg penethamate equivalent to 400 mg penethamate hydriodide

**3. PACKAGE SIZE**

50 ml

**4. TARGET SPECIES**

Cattle (lactating cows)

**5. INDICATION(S)**

**6. ROUTES OF ADMINISTRATION**

Intramuscular administration.

**7. WITHDRAWAL PERIODS**

Withdrawal period(s):  
Milk: 4 days.  
Meat and offal: 10 days.

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached, use within 28 days. Use by: \_\_\_\_/\_\_\_\_/\_\_\_\_

**9. SPECIAL STORAGE PRECAUTIONS**

Keep the vial in the outer carton.  
Store below 30 °C.  
Keep upright.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Eurovet Animal Health B.V.  
Handelsweg 25  
5531 AE Bladel  
The Netherlands

**14. MARKETING AUTHORISATION NUMBERS**

Vm 16849/5000

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

User warnings  
Penicillins and cephalosporins may occasionally cause severe allergic reactions.  
See package leaflet for user warnings.

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-V (To be supplied only on veterinary prescription')

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**Glass vial 50 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Revozyn RTU

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCE(S)**

Each ml contains:  
308.8 mg penethamate equivalent to 400 mg penethamate hydriodide

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/ yyyy}  
Once broached, use by: \_\_\_\_/\_\_\_\_/\_\_\_\_

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

50 ml

**6. ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**7. WITHDRAWAL PERIOD**

Withdrawal period(s):  
Milk: 4 days.  
Meat and offal: 10 days.

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET:

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Revozyn RTU 400 mg/ml suspension for injection for cattle

### 2. COMPOSITION

Each ml contains:

**Active substance:**

308.8 mg penethamate equivalent to 400 mg penethamate hydriodide

A white to yellowish white oily suspension.

### 3. TARGET SPECIES

Cattle (lactating cows)

### 4. INDICATIONS FOR USE

For the treatment of clinical and subclinical mastitis in lactating cows caused by staphylococci and streptococci, susceptible to penicillin.

### 5. CONTRAINDICATIONS

Do not use in cases of known hypersensitivity to the active substance, or to any of the excipients.

Do not administer by intravenous injection.

### 6. SPECIAL WARNING(S)

Cross-resistance has been shown between benzylpenicillin and penicillins and beta-lactam antimicrobials in staphylococci and streptococci. Use of benzylpenicillin should be carefully considered when susceptibility testing has shown resistance to penicillins or beta-lactam antimicrobials because its effectiveness may be reduced.

Special precautions for safe use in the target species:

Use of the product should be based on identification and susceptibility testing of the target pathogens. If this is not possible, therapy should be based on epidemiological information and knowledge of susceptibility of the target pathogens at farm level, or at local/regional level.

Use of the product should be in accordance with official, national and regional antimicrobial policies.

The feeding of waste milk containing residues of penicillin to calves should be avoided up to the end of the milk withdrawal period (except during the colostral phase), because it could select antimicrobial-resistant bacteria (e.g. ESBL) within the intestinal microbiota of the calf and increase the faecal shedding of these bacteria.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can cause sensitisation and contact dermatitis. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins, and *vice versa*.

Allergic reactions to these substances may occasionally be serious.

Handle this product with great care to avoid direct skin contact or self-injection.

People with known hypersensitivity to penicillin should avoid contact with the veterinary medicinal product.

Personal protective equipment consisting of gloves should be worn when handling the veterinary medicinal product.

Wash hands after use.

In case of accidental contact with the skin, wash immediately with plenty of water. If symptoms following exposure such as skin rash develop or in case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician. Swelling of the face, lips or eyes, or difficulty with breathing are more serious symptoms and require urgent medical attention.

Pregnancy and lactation:

Can be used during pregnancy or lactation.

Interactions with other medicinal products and other forms of interaction:

The veterinary medicinal product should not be administered concurrently with bacteriostatic antibiotics.

Overdose:

In case of overdose, no adverse effects other than those mentioned in section "Adverse events" are to be expected.

Special restrictions for use and special conditions for use:

For administration only by a veterinarian.

Major incompatibilities:

Do not mix with any other veterinary medicinal product.

## **7. ADVERSE EVENTS**

**Cattle (lactating cows):**

Very rare (<1 animal / 10,000 animals treated, including isolated reports): Urticaria, Anaphylactic shock<sup>a</sup>, death<sup>a</sup>. Sensitisation against penicillins.

Undetermined frequency (cannot be estimated from the available data): Skin reactions (mild), such as dermatitis.

<sup>a</sup> Anaphylactic shock can be fatal, very rarely

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to (the local representative of) the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: <https://www.gov.uk/report-veterinary-medicine-problem>.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For intramuscular administration only, preferably in the neck.  
Administer alternately on the left and the right side.

Administer 10-15 mg penethamate hydriodide per kg body weight per day, once daily for 3 consecutive days, corresponding to 2.5-3.75 ml of the veterinary medicinal product per 100 kg body weight per day, once daily for 3 consecutive days.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Shake well before use.

Avoid underdosing. To ensure a correct dosage, body weight should be determined as accurately as possible.

## **10. WITHDRAWAL PERIOD(S)**

Milk: 4 days.  
Meat and offal: 10 days.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store below 30 °C.  
Keep upright.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: 28 days.

When the container is breached/opened for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the container should be discarded should be determined. This discard date should be written in the space provided.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater.  
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

## **14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

Vm 16849/5000

Cardboard box with 1 x 50 ml vial.

## **15 DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED**

February 2023

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk).

## **16 CONTACT DETAILS**

Marketing authorisation holder and contact details to report suspected adverse reactions:

Eurovet Animal Health B.V.  
Handelsweg 25  
5531 AE Bladel  
The Netherlands  
Tel: +31 (0)348-563434

Manufacturer responsible for batch release:

Eurovet Animal Health B.V.  
Handelsweg 25  
5531 AE Bladel  
The Netherlands

Produlab Pharma B.V.  
Forellenweg 16  
4941 SJ Raamsdonksveer  
The Netherlands

**17. OTHER INFORMATION**

Approved 17 February 2023

A handwritten signature in black ink, consisting of a stylized initial 'A' followed by the name 'Hunter.' with a period.