PARTICULARS TO APPEAR ON <THE OUTER PACKAGE> <AND> <THE IMMEDIATE PACKAGE> {Carton}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

JOHNSON'S Veterinary PUPPY EASY-WORM SYRUP 4% w/v

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Contains Piperazine Hexahydrate 4% w/v (syrup)

3. PHARMACEUTICAL FORM

ROUNDWORM SYRUP FOR DOGS & PUPPIES OVER 2 WEEKS OF AGE & A MINIMUM BODYWEIGHT OF 0.25kg

4. PACKAGE SIZE

50ml

5. TARGET SPECIES

FOR DOGS & PUPPIES OVER 2 WEEKS OF AGE & A MINIMUM BODYWEIGHT OF 0.25kg (½ lb)

6. INDICATION(S)

Roundworm treatment for the control of ascarids (Toxascaris & Toxocara)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

ROUNDWORM SYRUP FOR DOGS & PUPPIES OVER 2 WEEKS OF AGE & A MINIMUM BODYWEIGHT OF 0.25kg (½ lb). Give before or during a main meal or add to a small portion of tasty food. Give at rate of 1ml for every ½kg (1.1 lb) bodyweight as per table, using plastic measuring cup provided, which must be washed immediately after use.

Less than 0.25kg (½ lb) - Consult a Veterinary Surgeon 0.25 to 0.5kg (½- 1lb) - 1 ml 0.6 to 1.0kg (1½- 2lb) - 2 ml 1.1 to 1.5kg (2 - 3lb) - 3 ml 1.6 to 2.0kg (3 - 4lb) - 4 ml 2.1 to 2.5kg (4 - 5lb) - 5 ml 2.6 to 5.0kg (5 -11lb) - 10 ml 5.1 to 7.5kg (11 -16lb) - 15 ml REPEAT ABOVE DOSE IN 14 DAYS TIME.

Do not repeat treatment if vomiting occurs shortly after dosing.

PUPPIES should be treated routinely for roundworms at 2 weeks of age and over 0.25kg bodyweight and every 2 weeks until 12 weeks of age; thereafter treat at 3 month intervals.

NURSING BITCHES should be dosed 2 weeks after giving birth & every 2 weeks until weaning. It is advisable to treat the bitch at same time as puppies. For medium & larger Bitches, Johnson's One Dose Easy Wormers Tablets are more suitable. It is advisable to consult a Veterinary Surgeon before treating pregnant animals & those with a history of epilepsy or disease of the liver or kidneys.

8. WITHDRAWAL PERIOD

Not applicable.

9. SPECIAL WARNING(S), IF NECESSARY

OPERATOR WARNINGS: Avoid contact with skin and eyes. In case of accidental skin or eye contact, wash the affected area with plenty of water. If irritation persists, seek medical advice. In case of accidental ingestion by humans, drink plenty of water and seek medical attention. Wash hands after handling product. Keep out of the reach of children.

USE WITHIN 3 MONTHS OF OPENING.	Date 1st opened:

10. EXPIRY DATE

Exp DD/MM/YY

11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C. Protect from light. Dispose of used packaging and remaining liquid in the household refuse. For animal treatment only. Keep container in outer carton.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of used packaging and remaining liquid in the household refuse.

13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]

For animal treatment only

A \	
AVM-GSL	

14. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"Keep out of the reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Johnson's Veterinary Products Limited

5 Reddicap Trading Estate

Sutton Coldfield, West Midlands

B75 7DF. UK

16. MARKETING AUTHORISATION NUMBER(S)

Vm 01759/4061

17. MANUFACTURER'S BATCH NUMBER

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS {Label}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

JOHNSON'S Veterinary PUPPY EASY-WORM SYRUP 4% w/v

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Contains Piperazine Hexahydrate 4% w/v (syrup)

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

50ml

4. ROUTE(S) OF ADMINISTRATION

ROUNDWORM SYRUP FOR DOGS & PUPPIES OVER 2 WEEKS OF AGE & A MINIMUM BODYWEIGHT OF 0.25kg (½ lb). Give before or during a main meal or add to a small portion of tasty food. Give at rate of 1ml for every ½kg (1.1 lb) bodyweight as per table, using plastic measuring cup provided, which must be washed immediately after use.

Less than 0.25kg (½ lb) - Consult a Veterinary Surgeon

$$0.25 \text{ to } 0.5\text{kg} (\frac{1}{2} - 1\text{lb}) - 1 \text{ ml}$$

0.6 to 1.0kg
$$(1\frac{1}{2}$$
- 2lb) - 2 ml

REPEAT ABOVE DOSE IN 14 DAYS TIME.

Do not repeat treatment if vomiting occurs shortly after dosing.

PUPPIES should be treated routinely for roundworms at 2 weeks of age and over 0.25kg bodyweight and every 2 weeks until 12 weeks of age; thereafter treat at 3 month intervals.

NURSING BITCHES should be dosed 2 weeks after giving birth & every 2 weeks until weaning. It is advisable to treat the bitch at same time as puppies. For medium & larger Bitches, Johnson's One Dose Easy Wormers Tablets are more suitable. It is advisable to consult a Veterinary Surgeon before treating pregnant animals & those with a history of epilepsy or disease of the liver or kidneys.

5. WITHDRAWAL PERIOD

Not Applicable.

- **6. BATCH NUMBER**
- 7. EXPIRY DATE

Exp DD/MM/YY

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only

Approved: 03/08/2017