

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE {50 ML, 100 ML, 250 ML  
/ CARDBOARD BOX}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Diatrim 200 mg/ml + 40 mg/ml solution for injection.

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Active substances:

Sulfadiazine	200 mg
Trimethoprim	40 mg

Excipient:

N-Methylpyrrolidone	510 mg
---------------------	--------

**3. PACKAGE SIZE**

50 ml, 100 ml, 250 ml

**4. TARGET SPECIES**

Cattle, pigs, dogs and cats.

**5. INDICATION(S)**

**6. ROUTES OF ADMINISTRATION**

For subcutaneous (dogs, cats), intramuscular or slow intravenous injection (cattle, pigs).

**7. WITHDRAWAL PERIODS**

Withdrawal periods:

*Cattle:*

Meat and offal: 12 days

Milk: 48 hours

*Pig:*

Meat and offal: 20 days

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached, use within 28 days.

Once broached use by: \_\_\_\_/\_\_\_\_/\_\_\_\_

#### **9. SPECIAL STORAGE PRECAUTIONS**

Do not store in a refrigerator after broaching.

#### **10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

#### **11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

#### **12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

#### **13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Eurovet Animal Health B.V.

#### **14. MARKETING AUTHORISATION NUMBERS**

Vm 16849/5011

#### **15. BATCH NUMBER**

Lot: {number}

#### **16. SPECIAL WARNING(S), IF NECESSARY**

##### Special warnings:

Pregnant women should take extra care when handling this product. See full user warnings in the package leaflet for details.

#### **17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

#### **18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

POM-V (Veterinary medicinal product subject to prescription)

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {50 ML/100 ML,  
250 ML / VIAL LABEL}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Diatrim 200 mg/ml + 40 mg/ml, solution for injection.

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:

Active substances:

Sulfadiazine	200 mg
Trimethoprim	40 mg

Excipient:

N-Methylpyrrolidone	510 mg
---------------------	--------

**3. TARGET SPECIES**

Cattle, pigs, dogs and cats.

**4. ROUTES OF ADMINISTRATION**

For subcutaneous (dogs, cats), intramuscular or slow intravenous injection (cattle, pigs).

**5. WITHDRAWAL PERIODS**

Withdrawal periods:

*Cattle:*

Meat and offal: 12 days

Milk: 48 hours

*Pig:*

Meat and offal: 20 days

**6. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached, use within 28 days.

Once broached use by \_\_\_\_ / \_\_\_\_ / \_\_\_\_

**7. SPECIAL STORAGE PRECAUTIONS**

Do not store in a refrigerator after broaching.  
Keep the vial in the outer carton.

## **8. NAME OF THE MARKETING AUTHORISATION HOLDER**

Eurovet Animal Health B.V.

## **9. BATCH NUMBER**

Lot: {number}

## **10. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

## **11. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

## **12. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

For animal treatment only.

POM-V (Veterinary medicinal product subject to prescription)

## **PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Diatrim 200 mg/ml + 40 mg/ml, solution for injection for cattle, pigs, dogs and cats.

### **2. COMPOSITION**

Each ml contains:

Active substances:

Sulfadiazine	200 mg
Trimethoprim	40 mg

Excipient:

N-Methylpyrrolidone	510 mg
---------------------	--------

Clear, greenish yellow to brownish yellow solution, practically free from particles.

### **3. TARGET SPECIES**

Cattle, pigs, dogs and cats.

### **4. INDICATIONS FOR USE**

Treatment of infections caused by, or associated with, organisms sensitive to the trimethoprim-sulfadiazine combination.

### **5. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

Do not use in animals with severe liver or renal damage or blood dyscrasias.

Do not use in case of reduced water intake or losses of body fluid.

### **6. SPECIAL WARNINGS**

Special precautions for safe use in the target species:

Use of the veterinary medicinal product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Use of the veterinary medicinal product deviating from the instructions given in the product leaflet may increase the prevalence of bacteria resistant to the veterinary medicinal product and may decrease the effectiveness of treatment with other antimicrobials or classes of antimicrobials due to the potential for cross-resistance. Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

The veterinary medicinal product may cause an allergic reaction in people sensitised to sulfonamides.

People with known hypersensitivity to sulfonamides should avoid contact with the veterinary medicinal product.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

This veterinary medicinal product may cause skin and eye irritation.

Avoid contact with skin or eyes.

In case of skin or eye contact, wash exposed area with plenty of clean water. If symptoms persist, seek medical advice.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

Laboratory studies in rabbits and rats with the excipient N-Methylpyrrolidone have shown evidence of foetotoxic effects. Women of childbearing age, pregnant women or women suspected of being pregnant should use the veterinary medicinal product with serious caution to avoid accidental self-injection.

Use during pregnancy or lactation:

Can be used during lactation.

The safety of the veterinary medicinal product has not been established in cattle, pigs, dogs and cats during pregnancy. Laboratory studies in rabbits and rats with the excipient N-Methylpyrrolidone have shown evidence of foetotoxic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

Local anaesthetics from the group of para-aminobenzoic acid esters (procaine, tetracaine) can locally inhibit the effect of sulfonamides.

Do not combine with other veterinary medicinal products.

Overdose (symptoms, emergency procedures, antidotes):

None known.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## 7. ADVERSE EVENTS

Cattle, pigs, dogs and cats:

Rare (1 to 10 animals / 10,000 animals treated):	Anaphylactic shock <sup>ab</sup>
---	----------------------------------

<sup>a</sup> Potentially fatal

<sup>b</sup> Mostly after intravenous injection. For intravenous administration the product should be warmed to body temperature and injected slowly over as long a period as is reasonably practical. At the first sign of intolerance the injection should be interrupted and shock treatment initiated.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in the package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or its local representative using the contact details at the end of this leaflet, or via your national reporting system {national system details}.

## **8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

For intramuscular, intravenous or subcutaneous use.

To ensure a correct dosage, the body weight of animals to be treated should be determined as accurately as possible.

### Cattle and pigs:

The recommended dose rate is 2.5 mg trimethoprim / 12.5 mg sulfadiazine per kilogram body weight (1 ml veterinary medicinal product per 16 kg body weight) by intramuscular or slow intravenous injection, once daily until 2 days after symptoms resolve up to a maximum of 5 days. The maximum intramuscular volume of injection per injection site is 5 ml for pigs and 15 ml in cattle. The veterinary medicinal product may be administered by intravenous injection when blood levels of trimethoprim and sulfadiazine are required more rapidly.

### Dogs and cats:

The recommended dose rate is 5 mg trimethoprim / 25 mg sulfadiazine per kilogram body weight (1 ml veterinary medicinal product per 8 kg body weight), by subcutaneous injection only, once daily until 2 days after symptoms resolve up to a maximum of 5 days. The recommended injection site in dogs is the loose skin at the top of the neck.

The closures must not be punctured more than 40 times.

## **9. ADVICE ON CORRECT ADMINISTRATION**

For intravenous administration the veterinary medicinal product should be warmed to body temperature and injected slowly over as long a period as is reasonably practical. At the first sign of intolerance the injection should be interrupted and shock treatment initiated. Intravenous administration should be used with extreme caution and only if therapeutically justified.

## 10. WITHDRAWAL PERIODS

### *Cattle:*

Meat and offal: 12 days  
Milk: 48 hours

### *Pig:*

Meat and offal: 20 days

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.  
Do not store in a refrigerator after broaching.  
Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton after Exp. The expiry date refers to the last day of that month.  
Shelf life after first opening the container: 28 days.

## 12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## 13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

## 14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 16849/5011

Pack sizes: 50 ml, 100 ml or 250 ml vials of colourless glass type II with a fluoropolymer coated chlorobutyl stopper type I secured with an aluminium cap.

1 vial in a cardboard box.  
Not all pack sizes may be marketed.

## 15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED

To be added at print

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk).



## 16. CONTACT DETAILS

Marketing authorisation holder and manufacturer responsible for batch release:

Eurovet Animal Health B.V.  
Handelsweg 25  
5531 AE Bladel  
The Netherlands

Local representative and contact details to report suspected adverse events:

Dechra Veterinary Products Limited  
Sansaw Business Park  
Hadnall  
Shrewsbury  
Shropshire  
SY4 4AS  
United Kingdom  
Tel: +44 (0)1939 211200

## 17. OTHER INFORMATION

Environmental properties:

Trimethoprim is persistent in soils.

POM-V (Veterinary medicinal product subject to prescription).

For animal treatment only.

Approved: 18 May 2024

*Gavin Hall*