

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Firodyl 62.5 mg chewable tablets for dogs
Firocoxib

2. STATEMENT OF ACTIVE SUBSTANCES

Firocoxib 62.5 mg

3. PHARMACEUTICAL FORM

Chewable tablet

4. PACKAGE SIZE

12 tablets
36 tablets
96 tablets

5. TARGET SPECIES

Dogs

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use
Read the package leaflet before use.

8. WITHDRAWAL PERIOD(S)

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}

11. SPECIAL STORAGE CONDITIONS

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Ceva Animal Health Ltd
Explorer House
Mercury Park
Wycombe Lane
Wooburn Green
High Wycombe
Buckinghamshire
HP10 0HH
United Kingdom

16. MARKETING AUTHORISATION NUMBER(S)

Vm 15052/4155

17. MANUFACTURER’S BATCH NUMBER

<Batch><Lot> {number}

MINIMUM PARTICULARS TO APPEAR ON BLISTERS

Blister

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Firodyl 62.5 mg chewable tablets for dogs
Firocoxib

2. NAME OF THE MARKETING AUTHORISATION HOLDER



3. EXPIRY DATE

EXP {month/year}

4. BATCH NUMBER

<Batch><Lot> {number}

5. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

B. PACKAGE LEAFLET

PACKAGE LEAFLET:

Firodyl 62.5 mg chewable tablets for dogs

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Ceva Animal Health Ltd
Explorer House
Mercury Park
Wycombe Lane
Wooburn Green
High Wycombe
Buckinghamshire
HP10 0HH
United Kingdom

Manufacturer responsible for batch release:

Ceva Santé Animale
Boulevard de la communication
Zone Autoroutière
53950 LOUVERNE
FRANCE

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Firodyl 62.5 mg chewable tablets for dogs
Firocoxib

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each tablet contains:

Active substance:

Firocoxib 62.5 mg

Round clover shaped tablet, Beige to light brown. Double scored on one side.

The tablets can be divided into equal quarters.

4. INDICATION(S)

For the relief of pain and inflammation associated with osteoarthritis in dogs.

For the relief of post-operative pain and inflammation associated with soft-tissue, orthopaedic and dental surgery in dogs.

5. CONTRAINDICATIONS

Do not use in pregnant or lactating bitches.

Do not use in animals less than 10 weeks of age or less than 3 kg body weight.

Do not use in animals suffering from gastrointestinal bleeding, blood dyscrasia or haemorrhagic disorders.

Do not use concomitantly with corticosteroids or other non-steroidal anti-inflammatory drugs (NSAIDs).

Do not use in known cases of hypersensitivity to the active substance or to any of the excipients.

6. ADVERSE REACTIONS

Emesis and diarrhoea have occasionally been reported. These reactions are generally of a transitory nature and are reversible when the treatment is stopped. Renal and/or hepatic disorders have been reported in very rare cases in dogs administered the recommended treatment dose. Rarely, nervous system disorders have been reported in treated dogs.

If adverse reactions like vomiting, repeated diarrhoea, faecal occult blood, sudden weight loss, anorexia, lethargy, degradation of renal or hepatic biochemistry parameters occur, use of the product should be stopped and the advice of a veterinarian should be sought. As with other NSAIDs, serious adverse effects can occur and, in very rare cases, may be fatal.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

Alternatively you can report via your national reporting system {national system details}.

7. TARGET SPECIES

Dogs

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Oral use.

5 mg/kg once daily, as presented in the table below.

For the reduction of post-operative pain and inflammation, the animals can be dosed starting approximately 2 hours before surgery for up to 3 consecutive days as needed. Following orthopaedic surgery and depending on the response observed, treatment using the same daily dosing schedule may be continued after the first 3 days, upon judgement of the attending veterinarian.

| Bodyweight (kg) | Number of tablets | | Dose range (mg/kg BW) |
|-----------------|-------------------|--------|-----------------------|
| | 62.5 mg | 250 mg | |
| 3.1 | 0.25 | | 5.0 |
| 3.2-6.2 | 0.5 | | 5.0-9.8 |
| 6.3-9.3 | 0.75 | | 5.0-7.4 |
| 9.4-12.5 | 1 | 0.25 | 5.0-6.6 |
| 12.6-15.5 | 1.25 | | 5.0-6.2 |
| 15.6-18.5 | 1.5 | | 5.1-6.0 |
| 18.6-21.5 | 1.75 | | 5.1-5.9 |
| 21.6-25 | | 0.5 | 5.0-5.8 |
| 25.1-37.5 | | 0.75 | 5.0-7.5 |
| 37.6-50 | | 1 | 5.0-6.6 |
| 50.1-62.5 | | 1.25 | 5.0-6.2 |
| 62.6-75 | | 1.5 | 5.0-6.0 |
| 75.1-87.5 | | 1.75 | 5.0-5.8 |
| 87.6-100 | | 2 | 5.0-5.7 |

9. ADVICE ON CORRECT ADMINISTRATION

The tablets are palatable, i.e. they are usually taken voluntarily by dogs (voluntary consumption of 76% of occasions in animals studied). If not, they can be given directly in the dog's mouth.

Tablets can be administered with or without food.

Instruction on how to divide the tablet: Put the tablet on an even surface, with its scored side facing down (convex face up). With the tip of the forefinger, exert slight vertical pressure on the middle of the tablet to break it along its width into halves. Then, in order to obtain quarters, exert slight pressure on the middle of one half with the forefinger to break it into two parts.

Do not exceed the recommended dose.

Duration of treatment will be dependent on the response observed. As field studies were limited to 90 days, longer-term treatment should be considered carefully and regular monitoring undertaken by the veterinarian.

10. WITHDRAWAL PERIOD(S)

Not applicable

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Any part-used tablet should be returned to the opened blister and used within 4 days.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

12. SPECIAL WARNING(S)

Special precautions for use in animals

As the tablets are flavoured, they should be stored in a safe place out of the reach of animals.

The recommended dose, as indicated in the dosing table, should not be exceeded.

Use in very young animals, or animals with suspected or confirmed impairment of renal, cardiac or hepatic function may involve additional risk. If such use cannot be avoided, those dogs require careful veterinary monitoring.

Avoid use in any dehydrated, hypovolaemic or hypotensive animals, as there is a potential risk of increased renal toxicity. Concurrent administration of potentially nephrotoxic drugs should be avoided.

Use this product under strict veterinary monitoring where there is a risk of gastrointestinal bleeding, or if the animal previously displayed intolerance to NSAIDs. Renal and/or hepatic disorders have been reported in very rare cases in dogs administered the recommended treatment dose. It is possible that a proportion of such cases had sub-clinical renal or hepatic disease prior to the commencement of therapy. Therefore, appropriate laboratory testing to establish baseline renal or hepatic biochemistry parameters is recommended prior to and periodically during administration.

The treatment should be discontinued if any of these signs are observed: repeated diarrhoea, vomiting, faecal occult blood, sudden weight loss, anorexia, lethargy, degradation of renal or hepatic biochemistry parameters.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product may be harmful following accidental ingestion.

In order to prevent children from accessing the product, tablets should be administered and stored out of sight and reach of children. Halved or quartered tablets should be returned to the open blister pocket and inserted into the outer carton.

Laboratory studies in rats and rabbits have shown evidence that firocoxib has the potential to effect reproduction and to induce malformations in fetuses. Pregnant women or women who are intending to become pregnant should administer the product with caution.

Wash hands after use of the product.

In the event of accidental ingestion of one or more tablets, seek medical advice immediately and show the package leaflet or the label to the doctor.

Pregnancy and lactation

Do not use in pregnant or lactating bitches.

Laboratory studies in rabbits have shown evidence of maternotoxic and foetotoxic effects at dose rates approximating the recommended treatment dose for the dog.

Interaction with other medicinal products and other forms of interaction

Pre-treatment with other anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before the commencement of treatment with the product. The treatment-free period, however, should take into account the pharmacokinetic properties of the products used previously.

The product must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Gastrointestinal tract ulceration may be exacerbated by corticosteroids in animals given non-steroidal anti-inflammatory drugs.

Concomitant treatment with molecules displaying action on renal flow, e.g. diuretics or Angiotensin Converting Enzyme (ACE) inhibitors, should be subject to clinical monitoring. Concurrent administration of potentially nephrotoxic drugs should be avoided as there might be an increased risk of renal toxicity. As anaesthetic drugs may affect renal perfusion, the use of parenteral fluid therapy during surgery should be considered to decrease potential renal complications when using NSAIDs peri-operatively.

Concurrent use of other active substances that have a high degree of protein binding may compete with firocoxib for binding and thus lead to toxic effects.

Overdose (symptoms, emergency procedures, antidotes):

In dogs ten weeks of age at the start of treatment at dose rates equal or greater to 25 mg/kg/day (5 times the recommended dose) for three months, the following signs of toxicity were observed: bodyweight loss, poor appetite, changes in the liver (accumulation of lipid), brain (vacuolisation), duodenum (ulcers) and death. At dose rates equal or greater to 15 mg/kg/day (3 times the recommended dose) for six months, similar clinical signs were observed, albeit that the severity and frequency were less and duodenal ulcers were absent.

In those target animal safety studies, clinical signs of toxicity were reversible in some dogs following cessation of therapy.

In dogs seven months of age at the start of treatment at dose rates greater than or equal to 25 mg/kg/day (5 times the recommended dose) for six months, gastrointestinal adverse effects, i.e. vomiting were observed.

Overdose studies were not conducted in animals over 14 months of age.
If clinical signs of overdosing are observed, discontinue treatment.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

September 2022

15. OTHER INFORMATION

Cardboard box with 12, 36, 96 tablets.

Not all pack sizes may be marketed.

Revised: September 2022
AN: 01507/2022

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Approved 27 September 2022

A handwritten signature in black ink, consisting of several loops and a long horizontal stroke at the end.