

ANNEX III
LABELLING AND PACKAGE LEAFLET

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

{BOX}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Katkin DE:WORM 230 mg/20 mg Film-coated Tablets for Cats
Pyrantel embonate/praziquantel

2. STATEMENT OF ACTIVE SUBSTANCES

Each film-coated tablet contains:

Pyrantel embonate	230 mg (equivalent to 80 mg pyrantel)
Praziquantel	20 mg

3. PHARMACEUTICAL FORM

Film-coated tablet

4. PACKAGE SIZE

2 x TABLETS

5. TARGET SPECIES

Cats.

6. INDICATION(S)

For the treatment of mixed infestations with roundworms, hookworms and tapeworms in cats.



ROUNDWORMS



TAPEWORMS



HOOKWORMS

MULTIWORMER FOR CATS

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Check your cat's weight in the table to see which dose they need.
Katkin DE:WORM is made for oral use only.

Dose:

Use 1 tablet per 4 kg body weight.

(5 mg praziquantel and 20 mg pyrantel base (57.5 mg pyrantel embonate) per kg of body weight.)

Cat's weight	Tablets
1.0 - 2.0 kg	1/2
2.1 - 4.0 kg	1
4.1 - 6.0 kg	1 1/2
6.1 - 8.0 kg	2

8. WITHDRAWAL PERIOD

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

Do not use in cats less than 6 weeks old or weighing less than 1 kg. Do not use during pregnancy. Do not use at the same time as other wormers, or if your cat has an allergy to any of the ingredients. If your cat is receiving other medication, check with your vet before using this product.

User Warnings

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician. Wash hands after use.

Keep the product in the outer package until ready to use.

10. EXPIRY DATE

EXP {month/year}

Shelf life of halved tablets after first opening the blister: 1 month.

11. SPECIAL STORAGE CONDITIONS

Storage:

Keep the blister in the outer carton.

Store unused parts of the halved tablets below 25°C. Each time an unused part-tablet is stored until next use, it should be returned to the open blister pocket and kept in a safe place out of the sight and reach of children.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal – see the package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.

AVM – GSL

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto
Šmarkeška cesta 6
8501 Novo mesto
Slovenia

16. MARKETING AUTHORISATION NUMBER

Vm 01656/3005

17. MANUFACTURER’S BATCH NUMBER

Lot

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

{BLISTER}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Katkin DE:WORM 230 mg/20 mg Film-coated Tablets for Cats
Pyrantel embonate/praziquantel

2. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

PACKAGE LEAFLET FOR:

Katkin DE:WORM 230 mg/20 mg Film-coated Tablets for Cats

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

KRKA, d.d., Novo mesto
Šmarkeška cesta 6
8501 Novo mesto
Slovenia

Manufacturer responsible for batch release:

KRKA - FARMA d.o.o., V. Holjevca 20/E, 10450 Jastrebarsko, Croatia
KRKA, d.d., Novo mesto, Šmarkeška cesta 6, 8501 Novo mesto, Slovenia

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Katkin DE:WORM 230 mg/20 mg Film-coated Tablets for Cats
Pyrantel embonate/praziquantel

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each film-coated tablet contains:

Active substances:

Pyrantel embonate	230 mg (equivalent to 80 mg pyrantel)
Praziquantel	20 mg

White to almost white, biconvex, oval film-coated tablet, scored on one side. The tablet can be divided into halves.

4. INDICATION(S)

For the treatment of mixed infestations with roundworms, hookworms and tapeworms in cats, caused by:

- adult stages of roundworms: *Toxocara cati* (*syn. mystax*)
- adult stages of hookworms: *Ancylostoma tubaeforme*, *Ancylostoma braziliense*
- tapeworms: *Echinococcus multilocularis*, *Dipylidium caninum*, *Hydatigera* (*Taenia*) *taeniaeformis*, *Mesocestoides spp.*, *Joyeuxiella pasqualei*.

Roundworms and tapeworms are parasites that live in the gut. If left untreated they can cause your cat to become unwell. Treating these infestations will reduce the risk of other animals and humans being exposed to worm infections.

5. CONTRAINDICATIONS

Do not use simultaneously with products containing piperazine as piperazine may block the action of pyrantel embonate contained in the product. Other worming products may contain piperazine.

Do not use simultaneously with other deworming products without veterinary advice.

Do not use in kittens less than 6 weeks of age,

Do not use in animals known to be allergic to the active substances (pyrantel embonate and praziquantel) or to any of the other ingredients (excipients).

Do not use during pregnancy.

6. ADVERSE REACTIONS

Mild and short-lived digestive tract disorders such as excessive salivation and/or vomiting and mild and short-lived nervous system disorders such as loss of balance may occur in very rare cases.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Cats.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Dosage:

5 mg praziquantel and 20 mg pyrantel base (57.5 mg pyrantel embonate) per kg of body weight. This corresponds to 1 tablet per 4 kg of body weight.

Body weight	Tablets
1.0 - 2.0 kg	1/2
2.1 - 4.0 kg	1
4.1 - 6.0 kg	1 1/2
6.1 - 8.0 kg	2

Kittens weighing less than 1 kg should not be treated with the product, because correct dosing of such cats may not be feasible.

Route of administration:

Oral use.

The tablets are to be given directly into the mouth but can be administered in a small amount of food, if necessary.

Duration of use:

Single treatment

In ascarid infestation, especially in kittens, complete elimination cannot be expected, so a risk of infection for humans can persist. Repeat treatments should therefore be carried out with a suitable roundworm product at 14 day intervals until 2-3 weeks after weaning. If signs of disease persist or appear, consult a veterinary surgeon.

9. ADVICE ON CORRECT ADMINISTRATION

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

10. WITHDRAWAL PERIOD

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store unused parts of the halved tablets below 25°C.

Each time an unused part-tablet is stored until next use, it should be returned to the open blister pocket and kept in a safe place out of the sight and reach of children.

Shelf life of halved tablets after first opening the immediate packaging: 1 month.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

12. SPECIAL WARNING(S)

Special warnings for each target species:

Tapeworm infestation occurs in cats at the earliest in the third week of life.

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*.

Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.

- Underdosing, this may be due to underestimation of body weight or misadministration of the product.

Do not use in cats less than 6 weeks of age or weighing less than 1 kg bodyweight.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

In the interest of good hygiene, persons administering the tablets directly to the cat or by adding them to the cat's food, should wash their hands afterwards.

Other precautions:

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), it must be reported to your veterinary surgeon, as specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

Pregnancy:

The safety of the veterinary medicinal product has not been established during pregnancy. The product should not be used during pregnancy but may be used during lactation.

Interaction with other medicinal products and other forms of interaction:

Do not use simultaneously with piperazine compounds, because the specific activities of piperazine (neuromuscular paralysis of the parasites) can inhibit the efficacy of pyrantel (spastic paralysis of the parasites). Do not use simultaneously with any other deworming products. If your cat receives other veterinary medicinal products, check with a veterinary surgeon or pharmacist before using this product.

Overdose (symptoms, emergency procedures, antidotes):

Symptoms of overdoses do not occur less than 5 times the recommended dose. The first expected sign of intoxication is vomiting.

Incompatibilities:

Not applicable.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials should be disposed of in accordance with local requirements. Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

14. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

15. OTHER INFORMATION

Box with 1 blister of 2 tablets.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

For Animal Treatment Only

AVM – GSL

Local representative:
KRKA UK Ltd, United Kingdom, Tel: 02071 646 156, Email: info.uk@krka.biz

Approved: 17 July 2024

Gavin Hall