

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**  
**cardboard box with 1 or 10 vials**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

AviPro ND C131  
Lyophilisate for suspension.

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each dose contains Newcastle Disease Virus, live attenuated, strain clone 13-1  
 $10^{6.0} - 10^{7.2}$  EID<sub>50</sub>

**3. PACKAGE SIZE**

10x 2,000, 10x 5,000 doses  
1x 2,000, 1x 5,000 doses

**4. TARGET SPECIES**

Chickens and turkeys.

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

For drinking water, ocular and nebulisation use.

**7. WITHDRAWAL PERIODS**

Withdrawal period: Zero days

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once reconstituted use within 2 hours.

**9. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.  
Do not freeze.  
Protect from light.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Elanco Europe Ltd

**14. MARKETING AUTHORISATION NUMBER**

Vm 00879/3031

**15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

glass vial

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

AviPro ND C131 Lyophilisate for suspension.

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

2000, 5000 doses live NDV, strain clone 13-1

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

## **B. PACKAGE LEAFLET**

## **PACKAGE LEAFLET**

### **1. Name of the veterinary medicinal product**

AviPro ND C131  
Lyophilisate for suspension for chickens and turkeys.

### **2. Composition**

#### **Active substance:**

Each dose contains  
Newcastle Disease Virus, live attenuated, strain clone 13-1  $10^{6.0}$  -  
 $10^{7.2}$  EID<sub>50</sub>

\*EID<sub>50</sub> = 50%-embryo infectious dose: the virus titre causing infection in 50% of the embryos inoculated with the virus.

Appearance: white-beige pellet

### **3. Target species**

Chickens and turkeys.

### **4. Indications for use**

Active immunisation of chickens and turkeys against Newcastle disease to reduce clinical signs and mortality.

#### **Chickens:**

Onset of immunity: 3 weeks after vaccination (7 days in seronegative chickens when vaccinated at 14 days of age).

Duration of immunity: 8 weeks after vaccination

#### **Turkeys:**

Onset of immunity: 2 weeks after vaccination

Duration of immunity: 8 weeks after vaccination

### **5. Contraindications**

None.

### **6. Special warnings**

#### Special warnings:

Vaccinate healthy animals only.

Maternally derived antibodies (MDA) may interfere with the development of a protective immune response following vaccination.

Special precautions for safe use in the target species:

**Chickens:**

The vaccine virus is excreted with faeces up to 12 days and may spread to susceptible animals by contact infection. However, ND negative contact animals do not show sero-conversion until 15 days after contact.

**Turkeys:**

The vaccine virus is excreted for less than 14 days after vaccination.

The vaccine virus may spread to susceptible non-vaccinated turkeys without inducing any clinical symptoms.

Transmission of the vaccine strain to ducks and geese is harmless. In pigeons slight pathological findings were observed in the respiratory tract, but no clinical symptoms occurred.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

ND-virus can induce conjunctivitis upon contact to eyes. Therefore during spraying eye- and inhalation protection (**face mask/visors**) must be worn.

Upon contact of the veterinary medicinal product to eyes seek medical advice.

Wash and disinfect hands and equipment after application.

Laying birds:

**Chickens:**

Safety data demonstrate that layers can be vaccinated during the laying period according to the recommended vaccine schedule (see "Dosage for each species, routes and method of administration").

In non-primed birds ND-vaccine virus was found in the oviduct after 10-fold over dosage. No egg transmission is observed in laying birds after basic immunisation.

The safety of the veterinary medicinal product has not been established in breeders during lay.

**Turkeys:**

The safety of the veterinary medicinal product has not been established during lay.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

**Chickens:**

Severity and duration of adverse reactions after the administration of a 10-fold dose are dependent on the (maternal) immune status as well as the general health condition of the chickens at the time point of vaccination.

**Turkeys:**

None.

**Special restrictions for use and special conditions for use:**

**Major incompatibilities:**

Do not mix with any substance other than water and skimmed milk or skimmed milk powder.

Ensure that the drinking water is cold, clean, non-chlorinated and free from detergents, disinfectants and metal ions.

**7. Adverse events**

**Chickens:**

Common (1 to 10 animals / 100 animals treated):	Respiratory tract disorders* like cough and sneezing
--	--

\*3 – 15 days after vaccination. This does not influence the performance of the birds. Severity and duration of adverse reactions are dependent on the (maternal) immune status as well as the general health condition of the chickens at the time point of vaccination.

**Turkeys:**

None.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

**8. Dosage for each species, routes and method of administration**

Species	Vaccination age	Administration route
Chickens	from 1 day onwards	nebulisation
	from 14 days onwards	nebulisation, ocular use, in drinking water use
Turkeys	from 21 days onwards	in drinking water use

## **9. Advice on correct administration**

Ensure that the drinking water is cold, clean, non-chlorinated and free from detergents, disinfectants and metal ions.

- Remove sealing cap and stopper from vaccine container.
- Suspend the vaccine in the corresponding amount of water and mix carefully.
- Prepare only the amount of vaccine that can be administered within 2 hours.
- The vaccine is ready for use.

### Ocular use (chickens)

The equipment used for eye drop application should be clean, free of detergents and disinfectants and should be used for vaccination purposes only.

For preparation of the vaccine use 34 ml of boiled and cooled drinking water per 1000 doses of vaccine.

Administer 1 droplet (corresponding to approximately 34 µl) into one eye of each chicken to be vaccinated by use of a pipette or dropper.

#### a) Nebulisation (chickens)

The amount of drinking water to be used for nebulisation depends on local and husbandry conditions.

After removing the stopper under water 1000 doses of vaccine are diluted as follows:

- 500 ml for 1000 chickens up to the 4<sup>th</sup> week of life
- 750 – 1000 ml for 1000 chickens after the 4<sup>th</sup> week of life.

The chickens are sprayed uniformly with a distance of 30 – 40 cm.

During and after vaccination ventilation should be switched off in order to avoid turbulences.

For primary vaccination during the 1<sup>st</sup> weeks of life a coarse spray having a droplet size of 100 µm and more should be used to avoid penetration into the lower parts of the respiratory tract and increased vaccination reactions.

#### b) In drinking water use (chickens and turkeys)

1. all equipment used for vaccination (tubes, drinkers etc.) are carefully cleaned and are free of detergents and disinfectants.
2. Estimate the amount of water according to the number of birds to be vaccinated (see 5.) Only cold clean water of drinking water quality should be used.

The addition of skimmed milk powder (2 – 4 g/l water) or skimmed milk (20 – 40 ml/l water) may positively influence the stability of the vaccine. Skimmed milk powder or skimmed milk must be carefully mixed with the water before dilution of the vaccine.

3. Remove aluminium-cap. Open the stopper of vaccine bottle under water and dilute the contents completely.

4. For easy handling the vaccine should be prepared in a small container (about 1 litre). Rinse the vial carefully and empty it completely. The vaccine suspension is then diluted in a larger vessel (5 – 10 l) and mixed well again.  
  
The complete content of the vaccine vials should be used for one flock or drinking water system only. Splitting of the diluted vaccine may lead to dosage errors.
5. To the vaccine suspension fresh cold water is added to a final volume that will be consumed by the birds within 1 – 2 hours. In case of doubt the uptake of water should be established the day before vaccination.
6. Drinker lines still filled with water must be drained before application of the vaccine suspension. The vaccine should be consumed within 2 hours. Since drinking behaviour of birds is varying it may be necessary to withdraw the drinking water for 2-3 hours before vaccination to ensure that all birds will drink during the vaccination phase. Every bird should receive an adequate dose of the vaccine.

#### **10. Withdrawal periods**

Zero days.

#### **11. Special storage precautions**

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

Shelf life after reconstitution according to directions: 2 hours

The complete content of opened containers should be used at once.

#### **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon how to dispose of medicines no longer required.

#### **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.

#### 14. Marketing authorisation numbers and pack sizes

Vm 00879/3031

The vaccine is available in the following packaging sizes:

- Box with 1 vial with 2000 doses
- Box with 10 vials with 2000 doses
- Box with 1 vial with 5000 doses
- Box with 10 vials with 5000 doses

Not all pack sizes may be marketed.

#### 15. Date on which the package leaflet was last revised

September 2023

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

#### 16. Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Elanco Europe Ltd,  
Form 2, Bartley Way,  
Bartley Wood Business Park,  
Hook,  
RG27 9XA,  
United Kingdom

##### **België/Belgique/Belgien**

{Nom/Naam/Name}  
<{Adresse/Adres/Anschrift }  
BE-0000 {Localité/Stad/Stadt}>  
Tél/Tel: + {N° de  
téléphone/Telefoonnummer/  
Telefonnummer}  
<{E-mail}>

##### **Lietuva**

{pavadinimas}  
<{adresas}  
LT {pašto indeksas} {miestas}>  
Tel: +370{telefono numeris}  
<{E-mail}>

##### **Република България**

{Наименование}  
<{Адрес}  
BG {Град} {Пощенски код}>  
Тел: + 359 {Телефонен номер}  
<{E-mail}>

##### **Luxembourg/Luxemburg**

{Nom}  
<{Adresse}  
L-0000 {Localité/Stadt}>  
Tél/Tel: + {N° de  
téléphone/Telefonnummer}  
<{E-mail}>

### **Česká republika**

{Název}  
<{Adresa}>  
CZ {město}>  
Tel: +{telefonní číslo}  
<{E-mail}>

### **Danmark**

{Navn}  
<{Adresse}>  
DK-0000 {by}>  
Tlf: + {Telefonnummer}  
<{E-mail}>

### **Deutschland**

{Name}  
<{Anschrift}>  
DE-00000 {Stadt}>  
Tel: + {Telefonnummer}  
<{E-mail}>

### **Eesti**

{Nimi}  
<{Address}>  
EE - (Postiindeks) (Linn)>  
Tel: +(Telefoninumber)  
<{E-mail}>

### **Ελλάδα**

{Όνομα}  
<{Διεύθυνση}>  
EL-000 00 {πόλη}>  
Τηλ: + {Αριθμός τηλεφώνου}  
<{E-mail}>

### **España**

{Nombre}  
<{Dirección}>  
ES-00000 {Ciudad}>  
Tel: + {Teléfono}  
<{E-mail}>

### **France**

{Nom}  
<{Adresse}>  
FR-00000 {Localité}>  
Tél: + {Numéro de téléphone}  
<{E-mail}>

### **Magyarország**

{Név}  
<{Cím}>  
HU-0000 {Város}>  
Tel.: + {Telefonszám}  
<{E-mail}>

### **Malta**

{Isem}  
<{Indirizz}>  
MT-0000 {Belt/Raħal}>  
Tel: + {Numru tat-telefon}  
<{E-mail}>

### **Nederland**

{Naam}  
<{Adres}>  
NL-0000 XX {stad}>  
Tel: + {Telefoonnummer}  
<{E-mail}>

### **Norge**

{Navn}  
<{Adresse}>  
N-0000 {poststed}>  
Tlf: + {Telefonnummer}  
<{E-mail}>

### **Österreich**

{Name}  
<{Anschrift}>  
A-00000 {Stadt}>  
Tel: + {Telefonnummer}  
<{E-mail}>

### **Polska**

{Nazwa/ Nazwisko:}  
<{Adres:}>  
PL – 00 000{Miasto:}>  
Tel.: + {Numer telefonu:}  
<{E-mail}>

### **Portugal**

{Nome}  
<{Morada}>  
PT-0000-000 {Cidade}>  
Tel: + {Número de telefone}  
<{E-mail}>

**Hrvatska**

{Ime}  
<{Adresa}>  
{Poštanski broj} {grad}>  
Tel: + {Telefonski broj}  
<{e-mail}>

**Ireland**

{Name}  
<{Address}>  
IE - {Town} {Code for Dublin}>  
Tel: + {Telephone number}  
<{E-mail}>

**Ísland**

{Nafn}  
<{Heimilisfang}>  
IS-000 {Borg/Bær}>  
Sími: + {Símanúmer}  
<{Netfang}>

**Italia**

{Nome}  
<{Indirizzo}>  
IT-00000 {Località}>  
Tel: + {Numero di telefono}>  
<{E-mail}>

**Κύπρος**

{Όνομα}  
<{Διεύθυνση}>  
CY-000 00 {πόλη}>  
Τηλ: + {Αριθμός τηλεφώνου}  
<{E-mail}>

**Latvija**

{Nosaukums}  
<{Adrese}>  
{Pilsēta}, LV{Pasta indekss }>  
Tel: + {Telefona numurs}  
<{E-mail}>

**România**

{Nume}  
<{Adresă}>  
{Oraș} {Cod poștal} – RO>  
Tel: + {Număr de telefon}  
<{E-mail}>

**Slovenija**

{Ime}  
<{Naslov}>  
SI-0000 {Mesto}>  
Tel: + {telefonska številka}  
<{E-mail}>

**Slovenská republika**

{Meno}  
<{Adresa}>  
SK-000 00 {Mesto}>  
Tel: + {Telefónne číslo}  
<{E-mail}>

**Suomi/Finland**

{Nimi/Namn}  
<{Osoite/Adress}>  
FI-00000 {Postitoimipaikka/Stad}>  
Puh/Tel: +  
{Puhelinnumero/Telefonnummer}  
<{E-mail}>

**Sverige**

{Namn}  
<{Adress}>  
SE-000 00 {Stad}>  
Tel: + {Telefonnummer}  
<{E-mail}>

**United Kingdom (Northern Ireland)**

{Name}  
<{Address}>  
{Town} {Postal code} – UK>  
Tel: + {Telephone number}  
<{E-mail}>>

Manufacturer responsible for batch release:

Lohmann Animal Health GmbH, Heinz-Lohmann-Str. 4, 27472 Cuxhaven, Germany

Approved 10 November 2023

