

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

Cardboard box 50 ml bottle  
Cardboard box 100 ml bottle

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Continance 40 mg/ml syrup

**2. STATEMENT OF ACTIVE SUBSTANCES**

phenylpropanolamine 40.28 mg/ml equivalent to phenylpropanolamine hydrochloride 50 mg/ml.

**3. PACKAGE SIZE**

50 ml + dosing syringe  
100 ml + dosing syringe

**4. TARGET SPECIES**

Dogs.

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Oral use.

**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

Exp. {mm/yyyy}  
Once broached use within 3 months.  
Use by...

**9. SPECIAL STORAGE PRECAUTIONS**

Keep the bottle and the syringe in the outer carton.

**10. THE WORDS "READ THE PACKAGE LEAFLET BEFORE USE"**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

FATRO S.p.A.

**14. MARKETING AUTHORISATION NUMBERS**

Vm 11557/3006

**15. BATCH NUMBER**

Lot {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**100 ml label**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Continence 40 mg/ml syrup

**2. STATEMENT OF ACTIVE SUBSTANCES**

phenylpropanolamine 40.28 mg/ml equivalent to phenylpropanolamine hydrochloride 50 mg/ml.

**3. TARGET SPECIES**

Dogs.

**4. ROUTES OF ADMINISTRATION**

Oral use.

**5. WITHDRAWAL PERIODS**

**6. EXPIRY DATE**

Exp. {mm/yyyy}  
Once broached use within 3 months.  
Use by...

**7. SPECIAL STORAGE PRECAUTIONS**

Keep the bottle and the syringe in the outer carton.

**8. NAME OF THE MARKETING AUTHORISATION HOLDER**

FATRO S.p.A.

**9. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**50 ml label**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Continence

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

phenylpropanolamine 40.28 mg/ml equivalent to phenylpropanolamine hydrochloride  
50 mg/ml.

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached use within 3 months.

Use by...

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET

### 1. Name of the veterinary medicinal product

Continence 40 mg/ml syrup for dogs

### 2. Composition

Each ml contains:

#### **Active substance:**

phenylpropanolamine 40.28 mg equivalent to phenylpropanolamine hydrochloride 50 mg

Clear colourless to pale yellow solution.

### 3. Target species

Dogs.

### 4. Indications for use

For the management of urinary incontinence associated with urethral sphincter incompetence in the bitch, particularly that associated with ovariohysterectomy.

### 5. Contraindications

Do not administer to patients treated with non-selective monoamine oxidase inhibitors.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not administer to pregnant or lactating bitches.

### 6. Special warnings

#### Special warnings:

The veterinary medicinal product should be avoided in hypertensive individuals.

In bitches less than 1 year old the possibility of anatomical disorders contributing to incontinence should be considered prior to treatment.

The use of the veterinary medicinal product is not appropriate for the treatment of behavioural causes of inappropriate urination.

#### Special precautions for safe use in the target species:

Phenylpropanolamine, a sympathomimetic drug, may affect the cardiovascular system, especially blood pressure and heart rate, and should be used with caution in animals with cardiovascular diseases.

Care should be exercised in treating animals with severe renal or hepatic insufficiency, diabetes mellitus, hyperadrenocorticism, glaucoma, hyperthyroidism or other metabolic disorders.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Phenylpropanolamine hydrochloride is toxic when ingested in higher doses. Adverse effects may include dizziness, headache, nausea, insomnia or restlessness, and increased blood pressure.

Accidental ingestion by a child may be fatal.

To avoid accidental ingestion, the veterinary medicinal product must be used and kept out of sight and reach of children. Always replace the cap securely after use and store the syringe and bottle inside the cardboard box at all times.

In case of accidental ingestion, seek medical attention immediately and show the package leaflet or the label to the physician.

In the event of accidental skin contact, wash the contaminated area with soap and water. Wash hands after handling the veterinary medicinal product.

This veterinary medicinal product may cause eye irritation. Avoid eye contact.

In case of accidental eye contact, rinse the eye thoroughly with clean water for about 15 minutes and consult a physician.

Pregnancy and lactation:

Do not use (during the whole or part of the pregnancy and lactation).

Interaction with other medicinal products and other forms of interaction:

Care should be exercised in administering the veterinary medicinal product with other sympathomimetic drugs, anticholinergic drugs, tricyclic antidepressants or specific type B monoamine oxidase inhibitors.

Overdose:

Lethargy and inappetence have been reported following an overdose of 2.5 mg/kg 3 times daily.

An overdose of phenylpropanolamine could produce symptoms of excessive stimulation of the sympathetic nervous system. Treatment should be symptomatic. Alpha-adrenergic blockers may be appropriate in the case of severe overdose.

Major incompatibilities:

None known.

**7. Adverse events**

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Hypersensitivity reaction.
Undetermined frequency (cannot be estimated from the available data):	Loose stool; Diarrhoea; Decreased appetite; Arrhythmia; Collapse; Dizziness; Aggression; Restlessness.

\*Sympathomimetics may produce a wide range of effects, most of which mimic the results of excessive stimulation of the sympathetic nervous system (e.g. effects on heart rate and blood pressure).

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: [adverse.events@vmd.gov](mailto:adverse.events@vmd.gov)

## 8. Dosage for each species, routes and method of administration

For oral use.

The recommended dose of phenylpropanolamine is 1.5 mg/kg bodyweight (equivalent to 0.15 ml per 5 kg bodyweight) twice daily in the feed. Alternatively, 1 mg/kg bodyweight (equivalent to 0.1 ml per 5 kg bodyweight) may be administered three times daily in the feed. The absorption rate is increased if the veterinary medicinal product is administered to fasted dogs.

## 9. Advice on correct administration


To ensure the correct dosage is administered the body weight of the animals should be determined as accurately as possible to avoid underdosing.



**1.** Remove the child-proof security cap by pressing firmly down and rotating anticlockwise.

**2.** Take the dosing syringe with the piston all the way down and insert the tip into the dosing syringe adaptor. Firmly push down.

**3.** Invert the bottle and, holding the syringe, pull the piston downwards aspirating the product slowly into the dosing syringe, to avoid the formation of air bubbles. Stop at the mark shown on the plunger corresponding to the required volume of the product.



**4.** Straighten the bottle and grasp the lower part of the syringe, close to the neck of the bottle. Remove the dosing syringe from the bottle by turning carefully.

**5.** Hold the dosing syringe on top of the dog's food and push the piston to the bottom to ensure delivery of the full dose of the product.

**6.** Replace the cap on the bottle and screw clockwise to close. Keep the bottle out of sight and reach of children.

**7.** Dry the tip with a clean cloth or paper. Wash the dosing syringe by removing the piston and rinse the items with hot water.

**8.** Dry carefully, making sure that the inside of the syringe is dry before reinserting the piston. Store the syringe inside the cardboard box to avoid access by children.

## **10. Withdrawal periods**

Not applicable.

## **11. Special storage precautions**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Keep the bottle and the syringe in the outer carton.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after Exp. The expiry date refers to the last day of that month.

Shelf-life after first opening the immediate packaging: 3 months.

## **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.

## **14. Marketing authorisation numbers and pack sizes**

Vm 11557/3006

### Pack sizes:

Cardboard box with 1 bottle of 50 ml + 1.5 ml dosing syringe

Cardboard box with 1 bottle of 100 ml + 1.5 ml dosing syringe

Not all pack sizes may be marketed.

## **15. PID LINK (Do not print heading)**

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

*Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).*

## 16. Contact details

### Marketing authorisation holder and manufacturer responsible for batch release:

FATRO S.p.A.  
Via Emilia, 285  
40064 Ozzano dell'Emilia BO  
Italy

### Local representatives and contact details to report suspected adverse reactions:

DUGV (UK) Ltd.  
Union House  
111 New Union Street  
Coventry, CV1 2NT, UK  
Phone: +353 (0) 504 43169  
uksales@dugganvet.com

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

UK(NI): POM-V

IE: POM (Prescription Only)

*Gavin Hall*

Approved: 05 June 2025