**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING WHERE THERE IS NO PACKAGE LEAFLET, i.e. Combined label and package leaflet {Bottle Labels}**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Dysect Sheep Pour-On solution 12.5 g/l

## 2. COMPOSITION

Each litre contains:

Active substance:
Alphacypermethrin 12.5 g

Excipient:
Copper Chlorophyll Complex (E141) 0.6 g

Clear dark green liquid free from solids.

## 3. PACKAGE SIZE

2.5 litres

5 litres

## 4. TARGET SPECIES

Sheep.

## 5. INDICATIONS FOR USE

### Indications for use

For the treatment and control of blowfly strike in sheep and lambs, and for the treatment of lice and ticks on sheep.

## 6. CONTRAINDICATIONS

### Contraindications

Do not treat lambs under one week old.
Do not treat animals in very hot weather.

## 7. SPECIAL WARNINGS

### Special warnings

Special precautions for safe use in the target species:
For external use only.

Because alphacypermethrin is retained in the wool grease of the fleece, it is recommended that the product be applied to sheep with a minimum wool length of 1.0cm.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:
Personal protective equipment consisting of waterproof apron, boots, and impervious gloves should be worn when handling the veterinary medicinal product and recently treated animals.
Wash hands and exposed skin before eating, drinking or smoking, and after work. Wash splashes from skin and eyes immediately. Use in a well-ventilated area. Remove heavily contaminated clothing immediately and wash before re-use.

Pregnancy and lactation:
The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Overdose:
No adverse reactions were recorded in double dose exposure trials.

## 8. ADVERSE EVENTS

### Adverse events

Sheep:

| | |
|---|---|
| Very rare (<1 animal / 10,000 animals treated, including isolated reports): | Fleece discolouration[1] |

[1] Observed from approximately 6 to 8 weeks after treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system at:

Website:https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine
e-mail: adverse.events@vmd.gov.uk

## 9. DOSAGE FOR EACH TARGET SPECIES, ROUTES, AND METHOD OF ADMINISTRATION

### Dosage for each species, routes, and method of administration

READ THE INSTRUCTIONS CAREFULLY BEFORE USE

| Pest | Dosage, ml<br>Sheep (> 25kg) | Lambs (< 25kg) |
|---|---|---|
| Blowfly | 40 | 25 |
| Lice | 40 | |
| Ticks | 40 | |
| Headfly | 5 | 5 |

For dosages of 40 ml and 25 ml, apply as two half-doses. Apply the first half-dose evenly along the back line from the neck to the saddle, and the second half-dose evenly from the saddle to the tail head and around the rump.

For the control of blowfly it is important that there is good coverage around the tail area. The T-bar must be held against the fleece so that the dose is applied directly into it. For the treatment of existing strike, apply the dose directly to the struck area. Most strikes will require the application of 5ml (saucer size) or 10ml (dinner plate size).

For the reduction of headfly strike, 5ml per animal is applied between the horns. Because alphacypermethrin is retained in the wool grease of the fleece, it is recommended that the product be applied to sheep with a minimum wool length of 1.0cm.

## 10. ADVICE ON CORRECT ADMINISTRATION

### Advice on correct administration

Remove the transit cap from the bottle and pierce the seal. Fit the spouted cap to the bottle. Fit the tube from the applicator gun to the spouted cap and fit the metal spring over the tube on the cap spout to ensure it is fixed securely. Check the gun is fitted with the T-bar and that it is working correctly. Before use the gun must be primed and set to the required volume. Clean the gun after use and maintain the gun to ensure accurate dosing.

The SimcroTech 25ml Pour-On Applicator gun is recommended as the product may have a detrimental effect on certain parts of conventional dosing guns.

Ready to use topical insecticidal solution. Treats a maximum of 124 ewes or a maximum of 200 lambs.

Contains the pyrethroid alphacypermethrin, which has both a repellent and insecticidal action to protect ewes and lambs from blowfly strike. The product has been formulated as a ready to use pour-on. Treated animals are easily identified by the natural green colour of the product which will fade after a few days without staining the fleece.

For the best results, apply before the spring or autumn tick rise and before the first appearance of headfly or blowfly; treat for lice when they are present. To ensure the best results the whole flock should be treated.

**Use:**

Blowfly – For the control of blowfly a single application will normally provide 8 to 10 weeks protection. Re-treatment may be necessary after this period.

For the treatment of blowfly strike, a single treatment applied to the infected area will ensure blowfly larvae are killed.

Ticks – A single treatment will normally provide control of ticks for 8 to 12 weeks.

Headfly – A single treatment will reduce the incidence for up to 6 weeks.

Lice – A single treatment will normally kill all lice.

## 11. WITHDRAWAL PERIODS

**Withdrawal periods**
Meat and offal: 49 days.
Milk: not permitted for use in animals producing milk for human consumption.

## 12. SPECIAL STORAGE PRECAUTIONS

**Special storage precautions**

Keep out of the sight and reach of children.

Do not store above 25 °C.
Store in a dry place.
Store in the original container.
Keep the container tightly closed.
Protect from direct sunlight.
Do not freeze.
If contents freeze, they must be thawed and thoroughly mixed before use.
Store away from food, drink, and animal feeding stuffs.

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

## 13. SPECIAL PRECAUTIONS FOR DISPOSAL

### Special precautions for disposal

Medicines should not be disposed of via wastewater.
DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the product or empty container.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## 14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

## 15. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 60021/3007

**Pack sizes: 2.5 litre and 5 litres.**

Not all pack sizes may be marketed.

## 16. PID LINK (Do not print heading)

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

## 17. CONTACT DETAILS

### Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:
Zoetis Belgium S.A.
2nd Floor, Building 10
Cherrywood Business Park
Loughlinstown
Dublin 18
D18 T3Y1
Ireland

Manufacturer responsible for batch release:
Zoetis Manufacturing Italia S.R.L.
Catania Branch
Via Franco Gorgone – Zona Industriale
1-95100 Catania CT
Italy

## 18. OTHER INFORMATION

POM-VPS

## 19. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

## 20. EXPIRY DATE

Exp {mm/yyyy}

## 21. BATCH NUMBER

Lot {number}

*Gavin Hall*

Approved: 18 March 2026