

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

White high density polyethylene bottles containing 250 ml of suspension with a white high density polyethylene screw cap.

White high density polyethylene bottles containing 1000 ml of suspension with a white high density polyethylene screw cap.

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Toltramax 50 mg/ml oral suspension for pigs  
Toltrazuril

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

1 ml contains:

**Active substance:**

Toltrazuril 50 mg

**Excipients:**

Sodium benzoate (E211) 2 mg

Sodium propionate (E281) 2 mg

**3. PHARMACEUTICAL FORM**

Oral suspension.

**4. PACKAGE SIZE**

250 ml

1000 ml

**5. TARGET SPECIES**

Pig (Piglets 3 to 5 days old).

**6. INDICATION(S)**

For the prevention of clinical signs of coccidiosis in neonatal piglets (3 to 5 days old) on farms with a confirmed history of coccidiosis caused by *Isospora suis*.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Shake well before use.

Oral use.

**8. WITHDRAWAL PERIOD**

Withdrawal period: Meat and offal: 77 days

**9. SPECIAL WARNING(S), IF NECESSARY**

Wash any splashes from skin or eyes immediately with water.  
Wash hands after use. Do not smoke, eat or drink whilst handling the product.  
People with known hypersensitivity to toltrazuril should avoid contact with the product.

**10. EXPIRY DATE**

EXP {month/year}  
Shelf-life after first opening the container: 3 months  
Once opened, use by...

**11. SPECIAL STORAGE CONDITIONS**

This veterinary medicinal product does not require any special storage conditions.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of waste material in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only - to be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder:  
Lavet Pharmaceuticals Ltd.  
Batthyány u. 6.  
Kistarcsa  
H-2143

Manufacturer for batch release:  
Lavet Pharmaceuticals Ltd.  
Batthyány u. 6.  
Kistarcsa  
H-2143

**16. MARKETING AUTHORISATION NUMBER**

Vm 32823/4009

**17. MANUFACTURER'S BATCH NUMBER**

Batch: {number}

## PACKAGE LEAFLET FOR:

Toltramax 50 mg/ml oral suspension for pigs

### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Lavet Pharmaceuticals Ltd.  
Batthyány u. 6.  
Kistarcsa  
H-2143

Manufacturer responsible for batch release:

Lavet Pharmaceuticals Ltd.  
Batthyány u. 6.  
Kistarcsa  
H-2143

### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Toltramax 50 mg/ml oral suspension for pigs  
Toltrazuril

### 3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

1 ml contains:

**Active substance:**

Toltrazuril 50 mg

**Excipients:**

Sodium benzoate (E211) 2 mg  
Sodium propionate (E281) 2 mg  
White or almost white suspension.

### 4. INDICATION(S)

For the prevention of clinical signs of coccidiosis in neonatal piglets (3 to 5 days old) on farms with a confirmed history of coccidiosis caused by *Isospora suis*.

### 5. CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

### 6. ADVERSE REACTIONS

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Pig (Piglets 3 to 5 days old).

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For oral use.

The oral suspension must be shaken before use.

Individual animal treatment.

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight. Treatment during an outbreak will be of limited value for the individual piglet because of damage to the small intestine having already occurred.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 ml is recommended.

The weight of animal should be accurately determined before treatment.

## **10. WITHDRAWAL PERIOD**

Meat and offal: 77 days

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of sight and reach of children.

Do not use after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Shelf-life after first opening the container: 3 months

## **12. SPECIAL WARNING(S)**

### **Special warnings for each target species**

As with any antiparasiticide, frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

### **Special precautions for use in animals**

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

In case of accidental ingestion seek medical advice immediately and show the package leaflet or the label to the physician.

People with known hypersensitivity to toltrazuril, or any of the excipients, should avoid contact with the veterinary medicinal product.

Avoid skin and eye contact with the product.

Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke whilst using the product.

**Interaction with other medicinal products and other forms of interaction**

None known, e.g. there is no interaction in combination with iron supplementation.

**Overdose (symptoms, emergency procedures, antidotes)**

No adverse effect has been observed in piglets after administration of a threefold overdose.

**Incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

March 2021

**15. OTHER INFORMATION**

White high density polyethylene bottles containing 250 or 1000 ml of suspension with a white high density polyethylene screw cap.

Not all pack sizes may be marketed.

For animal treatment only.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Approved 24 March 2021

A handwritten signature in black ink, appearing to read "A. Hunter.", is positioned below the approval date.