

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE Label of Cardboard**  
**Carton (Bottle): 1 x 10, 50, 125 Doses Label of 10-Pack Cardboard Carton**  
**(Bottles): 10 x 10, 50, 125 Doses**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Suvaxyn MH-One emulsion for injection.

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each 2 ml dose contains:

Inactivated *Mycoplasma hyopneumoniae*, strain P-5722-3 RP\* (undiluted)  $\geq 1.00$

\*Relative Potency unit determined by ELISA antigen quantification (*in vitro* potency test) compared to a reference vaccine.

**3. PACKAGE SIZE**

1x 10 doses  
1x 50 doses  
1x 125 doses  
10 x 10 doses  
10 x 50 doses  
10 x 125 doses

**4. TARGET SPECIES**

Pigs.

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Intramuscular use.

**7. WITHDRAWAL PERIODS**

Withdrawal period: Zero days.

**8. EXPIRY DATE**

Exp. (mm/yyyy)  
Once broached use immediately.

**9. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.  
Do not freeze.

Store in the original container.  
Protect from light.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited

**14. MARKETING AUTHORISATION NUMBERS**

Vm 42058/5123

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

POM-V Veterinary medicinal product subject to prescription
---

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING UNITS Bottle**  
**label: 10, 50 and 125 doses**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Suvaxyn MH-One emulsion for injection.

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each 2 ml dose contains:

Inactivated *Mycoplasma* RP\* (undiluted)  $\geq 1.00$   
*hyopneumoniae*, strain P-5722-  
3

\*Relative Potency unit determined by ELISA antigen quantification (*in vitro* potency test) compared to a reference vaccine.

10 doses  
50 doses  
125 doses

**3. TARGET SPECIES**

Pigs.

**4. ROUTES OF ADMINISTRATION**

Intramuscular use.  
Read the package leaflet before use.

**5. WITHDRAWAL PERIODS**

Withdrawal period: Zero days.

**6. EXPIRY DATE**

Exp. (mm/yyyy)  
Once broached use immediately.

**7. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.  
Do not freeze.  
Store in the original container.  
Protect from light.

**8. NAME OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited

**9. BATCH NUMBER**

Lot {Number}

**10. SPECIAL WARNING(S), IF NECESSARY**

**11. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**12. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

For animal treatment only.

POM-V Veterinary medicinal product subject to prescription
---

## **PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Suvaxyn MH-One emulsion for injection for pigs

### **2. COMPOSITION**

Each 2 ml dose contains:

#### **Active substance:**

Inactivated *Mycoplasma hyopneumoniae*,  
strain P-5722-3 RP\* (undiluted)  $\geq 1.00$

#### **Adjuvants:**

Carbopol # 941	4.00 mg
Squalane**	3.24 mg

\*Relative Potency unit determined by ELISA antigen quantification (*in vitro* potency test) compared to a reference vaccine.

\*\*As component of MetaStim (that also contains Pluronic L-121 and Polysorbate 80).

#### **Excipients:**

Thiomersal	0.20 mg
------------	---------

Brownish-grey emulsion.

### **3. TARGET SPECIES**

Pigs.

### **4. INDICATIONS FOR USE**

For the active immunisation of pigs of a minimum age of 7 days to reduce lung lesions that are caused by *Mycoplasma hyopneumoniae*.

Onset of immunity: 2 weeks.

Duration of immunity: 6 months after vaccination.

### **5. CONTRAINDICATIONS**

None.

### **6. SPECIAL WARNINGS**

Special warnings:

Vaccinate healthy animals only.

Special precautions for safe use in the target species:  
Avoid stress in the animals around the time of vaccination.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product contains animal oil. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy and lactation:  
Do not use in pregnant or lactating animals.

Interaction with other medicinal products and other forms of interaction:  
No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:  
After administration of a two-fold overdose by the recommended route to 3 weeks-old pigs, no other symptoms than those described under section "Adverse events" can be observed. However, the duration may be prolonged (body temperature increases up to 2 days and local tissue reactions up to 3 days) and the area of local tissue reactions may reach 1.0 cm in diameter. Administration of an overdose of the vaccine has not been investigated in 1 week-old piglets.

Special restrictions for use and special conditions for use:  
Not applicable.

Major incompatibilities:  
Do not mix with any other veterinary medicinal product.

## 7. ADVERSE EVENTS

Pigs:

Very common (>1 animal / 10 animals treated):
Injection site swelling <sup>1</sup>
Shivering <sup>2</sup>
Bristling of hairs <sup>2</sup>
Depression <sup>2</sup> , Elevated temperature <sup>2/3</sup>
Uncommon (1 to 10 animals / 1,000 animals treated):
Anaphylactic-type (severe allergic) reaction
Neurological signs

<sup>1</sup>May reach 0.3 cm in diameter (palpable, but not visible) and last for up to 2 days.

<sup>2</sup>Within 4 hours after vaccination and spontaneously resolving within 24 hours without treatment.

<sup>3</sup>Body temperature increase up to 1.9°C.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

## **8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

One dose (2 ml) per animal should be administered intramuscularly in the neck to pigs from the age of 7 days onwards.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Shake vaccine well before administration and intermittently during the process of vaccination.

It is good practice to allow the vaccine to warm to body temperature in the hand or pocket before administration, to avoid the discomfort of injection of a cold liquid.

## **10. WITHDRAWAL PERIODS**

Zero days.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Store in the original container.

Protect from light.

Do not use this veterinary medicinal product after the expiry date stated on the label and outer package after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: use immediately.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

## 14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 42058/5123

Carton box containing 1 or 10 bottles of 10, 50 or 125 doses.

Not all pack sizes may be marketed.

## 15. PID LINK (Do not print heading)

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

## 16. CONTACT DETAILS

Marketing authorisation holder and contact details to report suspected adverse reactions:

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP  
Tel: +44 (0) 345 300 8034

Manufacturer responsible for batch release:  
Zoetis Manufacturing & Research Spain S.L.  
Carretera De Camprodon S/n  
La Vall De Bianya  
17813 Girona  
Spain

## 17. OTHER INFORMATION

To stimulate active immunity against *Mycoplasma hyopneumoniae*.  
Post-vaccination serum antibody levels are not related to the degree of protection afforded by vaccination.

For animal treatment only.

POM-V Veterinary medicinal product subject to prescription
---

Approved 18 May 2024  
*Gavin Hall*