

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Carprodyl Quadri 120 mg Chewable Tablets for Dogs

Carprofen

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

One tablet contains:

120mg of Carprofen

3. PACKAGE SIZE

12 tablets

120 tablets

240 tablets

480 tablets

4. TARGET SPECIES

Dogs

5. INDICATION(S)

6. ROUTES OF ADMINISTRATION

For oral administration.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

EXP: {month/year}

9. SPECIAL STORAGE PRECAUTIONS

Do not store above 30°C.

Protect from light.

For shelf life of divided tablets: see package leaflet.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER



14. MARKETING AUTHORISATION NUMBERS

Vm 14966/5024

15. BATCH NUMBER

Lot:

16. SPECIAL WARNING(S), IF NECESSARY

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet


18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

POM-V – Veterinary medicinal product subject to prescription

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Carprodyl Quadri



2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

3. BATCH NUMBER

Lot:

4. EXPIRY DATE

EXP: {month/year}

5. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Carprodyl Quadri 120 mg Chewable Tablets for Dogs
Carprofen

2. COMPOSITION

One tablet contains:
Carprofen.....120 mg

3. TARGET SPECIES

Dogs

4. INDICATIONS FOR USE

In the dog:
Reduction of inflammation and pain caused by musculo-skeletal disorders and degenerative joint disease.
As a follow-up to parenteral analgesia in the management of post-operative pain.

5. CONTRAINDICATIONS

Do not use in pregnant and lactating bitches.
Do not use in dogs aged less than 4 months in the absence of specific data.
Do not use in cats.
Do not use in dogs, suffering from cardiac, hepatic or renal disease, when there is a possibility of gastrointestinal ulceration or bleeding or where there is evidence of blood dyscrasia.
Do not use in case of hypersensitivity to the active substance, to other NSAIDs and to any of the excipients.

6. SPECIAL WARNINGS

Use during pregnancy and lactation

Studies in laboratory species (rat and rabbit) have shown evidence of foetotoxic effects of carprofen at doses close to the therapeutic dose. The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Do not use in pregnant or lactating bitches.
For breeding animals, do not use during reproduction period.

Special precautions for use in animals

Use in aged dogs, may involve additional risk. If such a use cannot be avoided, dogs may require a reduced dosage and careful clinical management. Ask your veterinary surgeon for advice.

Avoid use in any dehydrated, hypovolaemic or hypotensive dog, as there is a potential risk of increased renal toxicity.

Concurrent administration of potential nephrotoxic drugs should be avoided.

NSAIDs can cause inhibition of phagocytosis and hence in the treatment of

inflammatory conditions associated with bacterial infection, appropriate concurrent antimicrobial therapy should be instigated.

As with other NSAIDs, photodermatitis during treatment with carprofen has been observed in laboratory animals and in humans. These skin reactions have never been observed in dogs.

Do not administer other NSAIDs concurrently or within 24 hours of each other.

Due to the good palatability of the tablet, they should be stored in a safe place out of the reach of animals. Intake of dose exceeding the recommended number of tablets may lead to severe adverse effects. If this is the case, seek veterinary assistance immediately.

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet to the physician.

Wash hands after handling the product.

Interaction with other medicinal products and other forms of interaction

Carprofen is highly bound to plasma proteins and compete with other highly bound drugs, which can increase their respective toxic effects.

Do not use this veterinary medicinal drug concurrently with other NSAIDs or with glucocorticoids.

Concurrent administration of potentially nephrotoxic drugs (e.g. aminoglycoside antibiotics) should be avoided.

Overdose

Bibliographic data report that carprofen is well tolerated in dogs at twice the recommended dosage for 42 days.

Doses up to 3 times the recommended dose are reported to be without adverse effects.

There is no specific antidote to carprofen but general supportive therapy as applied to clinical overdose with NSAIDs should be applied.

7. ADVERSE EVENTS

Dog:

Rare (1 to 10 animals / 10,000 animals treated):
Renal disorder ¹ Hepatic disorder ^{1,3}
Very rare (<1 animal / 10,000 animals treated, including isolated reports):
Vomiting ² , Soft stool/Diarrhoea ² , Blood in faeces ² Appetite loss ² , Lethargy ²

¹As with other NSAIDs

²Typical undesirable effects associated with NSAIDs, these adverse reactions

generally occur within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

³Idiosyncratic effects

If adverse reactions occur, use of the product should be stopped and the advice of a veterinarian should be sought.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or the national competent authority via the national reporting system. National contact details: <https://www.gov.uk/report-veterinary-medicine-problem>

8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

For oral administration.

4 mg carprofen per kg bodyweight per day.

An initial dose of 4 mg carprofen per kg bodyweight per day given as a single daily dose. The analgesic effect from each dose persists for at least 12 hours.

The daily dose may be reduced, subject to clinical response.

Duration of treatment will be dependant upon the response seen. Long term treatment should be under regular veterinary supervision. To extend analgesic and anti-inflammatory cover post-operatively parenteral pre-operative treatment with an injectable carprofen may be followed with carprofen tablets at 4 mg/kg/day for 5 days.

Do not exceed the stated dose.

Tablets can be broken as follows: Put the tablet on a flat surface, with its scored side facing the surface (convex face up).

With the tip of forefinger, exert a slight vertical pressure on the middle of the tablet to break it into halves. In order to obtain quarters, then exert a slight pressure on the middle of one half with forefinger.

Each chewable tablet can be quartered for accurate dosing according to the individual body weight of the animal.

Number of tablets per day	Dog weight (kg)	
$\frac{1}{4}$	<input type="checkbox"/> 7.5	< 14.4
$\frac{1}{2}$	<input type="checkbox"/> 14.5	< 20.9
$\frac{3}{4}$	<input type="checkbox"/> 21	< 29.9
1	<input type="checkbox"/> 30	< 37.4
$1 \frac{1}{4}$	<input type="checkbox"/> 37.5	< 44.9
$1 \frac{1}{2}$	<input type="checkbox"/> 45	< 52.4
$1 \frac{3}{4}$	<input type="checkbox"/> 52.5	< 59.9
2	<input type="checkbox"/> 60	< 70

9. ADVICE ON CORRECT ADMINISTRATION

The tablets are flavoured, and are accepted by dogs, but they may be administered directly in the mouth of the dog or added to food if necessary.

10. WITHDRAWAL PERIODS

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not store above 30 °C.

Protect from light.

Divided tablets should be stored in the blister pack. Any divided tablet portions remaining after 72 hours should be discarded.

Do not use the veterinary medicinal product after the expiry date which is stated on the blister and outer carton. The expiry date refers to the last day of that month.

12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

POM-V – Veterinary medicinal product subject to prescription

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 14966/5024

Cardboard box with 2 blisters of 6 tablets.

Cardboard box with 20 blisters of 6 tablets.

Cardboard box with 40 blisters of 6 tablets.

Cardboard box with 80 blisters of 6 tablets.

Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder:

Ceva Sante Animale
8 rue de Logrono
33500 Libourne
France

Contact details to report suspected adverse reactions:

Ceva Animal Health Ltd
Explorer House
Mercury Park
Wycombe Lane
Wooburn Green
High Wycombe
Buckinghamshire
HP10 0HH
United Kingdom

Manufacturer responsible for batch release:

Ceva Santé Animale
Boulevard de la Communication
Zone Autoroutière
53950 Louverné
France.

17. OTHER INFORMATION

Gavin Hall
Approved: 26 September 2025