

ANNEX

Registration number of additive	Name and registration number of person responsible for putting additive into circulation	Additive (Trade name)	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions		End of period of authorisation	Provisional Maximum Residue Limit (MRLs) in the relevant food-stuffs of animal origin
								ng of active substance/kg of complete feedingstuff with a moisture content of 12 %			
Coccidiostats and histomonostats											
E 1701	Huvepharma NV Belgium	Monensin sodium Coxidin	Active substance: $C_{36}H_{61}O_{11}Na$ Sodium salt of polyether monocarboxylic acid produced by Streptomyces cinnamoneensis, 28682, LMG S-19095 in powder form. Factor composition: Monensin A: not less than 90 % Monensin: A + B: not less than 95 % Monensin C: 0,2-0,3 %	Chickens for fattening Turkeys	— 16 weeks	100 90	125 100	1. Use prohibited at least three days before slaughter. 2. The additive shall be incorporated in compound feeding-stuffs in form of a premixture. 3. Maximum permitted dose of monensin sodium in complementary feedingstuffs: — 625 mg/kg for chickens for fattening; — 500 mg/kg for turkeys.	6.2.2017	25 µg monensin sodium/kg of wet skin + fat. 8 µg monensin sodium/kg of wet liver, kidney and muscle.	
Additive composition:											
Monensin sodium technical substance equivalent to monensin activity: 25 % Perlite: 15-20 % Wheat bran 55-60 %											
Analytical method⁽¹⁾											
HPLC method											

⁽¹⁾ Details of the analytical methods are available at the following address of the Community Reference Laboratory: [www.irmm.jrc.be/\[html\]/crfaa/](http://www.irmm.jrc.be/[html]/crfaa/)