

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One tablet contains:

**Active substances:**

Milbemycin oxime	4 mg
Praziquantel	10 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Film-coated tablet.

Oblong shaped, beige to brown, artificial beef flavoured tablet with a score on both sides. One side bears the imprint "BC", the other side "NA".

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cats

#### **4.2 Indications for use, specifying the target species**

In cats: treatment of mixed infections by immature and adult cestodes **and** nematodes of the following species:

- Cestodes:
  - Dipylidium caninum*
  - Taenia spp.*
  - Echinococcus multilocularis*
  
- Nematodes:
  - Ancylostoma tubaeforme*
  - Toxocara cati*

Prevention of heartworm disease (*Dirofilaria immitis*) if concomitant treatment against cestodes is indicated.

#### **4.3 Contraindications**

Do not use in cats of less than 6 weeks of age and/or weighing less than 0.5 kg.

#### 4.4 Special warnings for each target species

In order to develop an effective worm control programme local epidemiological information and the risk of exposure of the cat should be taken into account. It is recommended to treat all the animals living in the same household concomitantly. When infection with the cestode *D. caninum* has been confirmed, concomitant treatment against intermediate hosts, such as fleas and lice, should be discussed with a veterinarian to prevent re-infection.

#### 4.5 Special precautions for use

##### Special precautions for use in animals

As per good veterinary practice, animals should be weighed to ensure accurate dosing. Ensure cats and kittens weighing between 0.5 kg and ≤2 kg receive the appropriate tablet strength (4 mg MBO/10 mg praziquantel) and the appropriate dose (1/2 or 1 tablet) for the corresponding weight band (1/2 tablet for cats weighing 0.5 to 1 kg ; 1 tablet for cats weighing >1 to 2 kg – 1 tablet).

Echinococcosis represents a hazard for humans. In case of Echinococcosis, specific guidelines on the treatment and follow up and on the safeguard of persons have to be followed. Experts or institutes of parasitology should be consulted.

No studies have been performed with severely debilitated cats or individuals with seriously compromised kidney or liver function. The product is not recommended for such animals or only according to a benefit/risk assessment by the responsible veterinarian.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

In the event of accidental ingestion of the tablets, particularly by a child, seek medical advice immediately and show the package leaflet or the label to the physician.

#### 4.6 Adverse reactions (frequency and seriousness)

In very rare occasions, especially in young cats, hypersensitivity reactions, systemic signs (such as lethargy), neurological signs (such as ataxia and muscle tremors) and/or gastrointestinal signs (such as emesis and diarrhoea) have been observed after administration of the veterinary medicinal product.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens can be used in breeding cats including pregnant and lactating queens.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

The concurrent use of Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens with selamectin is well tolerated. No interactions were observed when the recommended dose of the macrocyclic lactone selamectin was administered during treatment with Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens at the recommended dose.

Although not recommended, the concomitant use of Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens with a spot on containing moxidectin and imidacloprid at recommended dose rates following a single application was well tolerated in one laboratory study in 10 kittens.

The safety and efficacy of the concurrent use have not been investigated in field studies. In the absence of further studies, caution should be taken in the case of concurrent use of the product with any other macrocyclic lactone. Also, no such studies have been performed with reproducing animals.

#### **4.9 Amounts to be administered and administration route**

Minimum recommended dose rate: 2 mg of milbemycin oxime and 5 mg of praziquantel per kg are given orally as a single dose. The product should be administered with or after some food. Doing so ensures optimum protection against heartworm disease.

Depending on the bodyweight of the cat, the practical dosing is as follows:

Weight	Tablets
0.5 - 1 kg	½ tablet
> 1 - 2 kg	1 tablet

Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens can be inserted into a programme for prevention of heartworm disease if at the same time treatment against tapeworms is indicated. Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens has a duration of heartworm prevention of one month. For prevention of heartworm disease the use of a monosubstance is preferred.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In case of overdose, in addition to signs observed at the recommended dose (see section 4.6), drooling was observed. This sign will usually disappear spontaneously within a day.

#### **4.11 Withdrawal period(s)**

Not applicable.

## 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antiparasitic products, insecticides and repellants - endectocides

ATC vet Code: QP54AB51 (milbemycin oxime, combinations)

### 5.1 Pharmacodynamic properties

Milbemycin oxime belongs to the group of macrocyclic lactones, isolated from the fermentation of *Streptomyces hygroscopicus* var. *aureolacrimosus*. It is active against mites, against larval and adult stages of nematodes as well as against larvae of *Dirofilaria immitis*.

The activity of milbemycin is related to its action on invertebrate neurotransmission: Milbemycin oxime, like avermectins and other milbemycins, increases nematode and insect membrane permeability to chloride ions via glutamate-gated chloride ion channels (related to vertebrate GABA<sub>A</sub> and glycine receptors). This leads to hyperpolarisation of the neuromuscular membrane and flaccid paralysis and death of the parasite.

Praziquantel is an acylated pyrazino-isoquinoline derivative. Praziquantel is active against cestodes and trematodes. It modifies the permeability for calcium (influx of Ca<sup>2+</sup>) in the membranes of the parasite inducing an imbalance in the membrane structures, leading to membrane depolarisation and almost instantaneous contraction of the musculature (tetany), rapid vacuolization of the syncytial tegument and subsequent tegumental disintegration (blebbing), resulting in easier expulsion from the gastrointestinal tract or death of the parasite.

### 5.2 Pharmacokinetic particulars

In the cat, praziquantel reaches peak plasma concentrations within an hour after oral administration.

The half life of elimination is around 3 hours.

In the dog, there is rapid hepatic biotransformation, principally to monohydroxylated derivatives.

The principal route of elimination in the dog is renal.

After oral administration in the cat, milbemycin oxime reaches peak plasma concentrations within 2 hours. The half life of elimination is around 13 hours ( $\pm$  9 hours). In the rat, metabolism appears to be complete although slow, since unchanged milbemycin oxime has not been found in urine or feces. Main metabolites in the rat are monohydroxylated derivatives, attributable to hepatic biotransformation. In addition to relatively high liver concentrations, there is some concentration in fat, reflecting its lipophilicity.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Core:

Cellulose, microcrystalline  
Croscarmellose sodium  
Povidone  
Lactose monohydrate  
Silica, colloidal anhydrous  
Magnesium stearate

Coat:

Hypromellose  
Macrogol  
Talc  
Artificial beef flavour

### **6.2 Major incompatibilities**

Not applicable.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years  
Shelf-life after first opening of the immediate packaging: 6 months

### **6.4 Special precautions for storage**

Do not store above 25 °C  
Keep the blister in the outer carton in order to protect from light

### **6.5 Nature and composition of immediate packaging**

PVC/PE/PVdC/aluminium blister

Available pack sizes:

Box with 2 tablets in blister  
Box with 4 tablets in blister  
Box with 10 tablets in blister  
Box with 20 tablets in blister  
Box with 50 tablets in blister  
Box with 100 tablets in blister

Not all pack sizes may be marketed

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

Elanco Milbemycin - Praziquantel 4 mg/10 mg film-coated tablets for small cats and kittens should not enter water courses as this may be dangerous for fish and other aquatic organisms.

**7. MARKETING AUTHORISATION HOLDER**

Elanco Europe Ltd  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

**8. MARKETING AUTHORISATION NUMBER**

Vm 00879/5008

**9. DATE OF FIRST AUTHORISATION**

29 December 2022

**10. DATE OF REVISION OF THE TEXT**

December 2022

A handwritten signature in black ink, consisting of several vertical strokes followed by a horizontal line that curves upwards and to the right.

Approved 29 December 2022