

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

EziWormer Plus Tablets for Dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains:

Active substances:

Praziquantel 50 mg

Pyrantel 50 mg (equivalent to 144 mg pyrantel embonate)

Febantel 150 mg

Excipients:

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablet

Pale yellow tablet with a cross breakline on one side.

The tablets can be divided into halves or quarters.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs.

#### **4.2 Indications for use, specifying the target species**

Treatment of the following gastrointestinal roundworms, tapeworms, hookworms and whipworms in dogs and puppies of the following species:

**Ascarids:** *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms).

**Hookworms:** *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

**Whipworms:** *Trichuris vulpis* (adults).

**Tapeworms:** *Echinococcus* species, (*E. granulosus*, *E. multilocularis*), *Taenia* species, (*T. hydatigena*, *T. pisiformis*, *T. taeniformis*), *Dipylidium caninum* (adult and immature forms).

#### **4.3 Contraindications**

Do not use simultaneously with piperazine compounds as piperazine may block the action of pyrantel embonate contained in EziWormer Plus Tablets for Dogs. Other worming products may contain piperazine.

Do not use in animals with a known sensitivity to the active ingredients or to any of the excipients.

#### **4.4 Special warnings for each target species**

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

Tapeworm infestation is unlikely in pups less than 6 weeks of age.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Dogs may become infected with worms by eating insects (including fleas and lice), birds, small rodents, rabbits or raw offal from affected sheep, goats and cattle. Dogs will continue to be re-infected unless the route of infection is controlled e.g. treating a flea infestation or preventing a dog from scavenging or hunting.

*Echinococcus multilocularis* is a tapeworm of foxes, dogs and cats which is common in parts of Europe. It can also infect humans potentially causing severe liver disease which can be fatal. To prevent *Echinococcus multilocularis* establishing in the UK and Ireland it is a requirement that all dogs must be treated with a product containing praziquantel before entering the country.

#### **4.5 Special precautions for use**

##### **i. Special precautions for use in animals**

Any part used tablet should be discarded.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Do not exceed the stated dose when treating pregnant bitches.

##### **ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals**

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards.

Echinococcosis represents a hazard for humans. As echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of humans, need to be obtained from the relevant competent authority.

#### **4.6 Adverse reactions (frequency and seriousness)**

In very rare cases slight and transient digestive tract disorders such as vomiting and /or diarrhoea may occur. In individual cases these signs can be accompanied by nonspecific signs such as lethargy, anorexia or hyperactivity.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals treated displaying adverse reaction(s))

- Common (more than 1 but less than 10 animals in 100 animals treated)
- Uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- Rare (more than 1 but less than 10 animals in 10,000 animals treated)
- Very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

#### 4.7 Use during pregnancy, lactation or lay

Consult a veterinary surgeon before treating pregnant animals. It is recommended that the product is not used in dogs during the first 4 weeks of pregnancy. Do not exceed the stated dose when treating pregnant bitches. Teratogenic effects attributed to high doses of febantel have been reported in sheep and rats. No studies have been performed in dogs during early pregnancy.

#### 4.8 Interaction with other medicinal products and other forms of interaction

Do not use simultaneously with piperazine compounds (see section 4.3 above). Concurrent use with other cholinergic compounds can lead to toxicity.

#### 4.9 Amounts to be administered and administration route

For oral administration only.

The recommended dose rates are: 15mg/kg bodyweight febantel, 5 mg/kg pyrantel (equivalent to 14.4 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg (22 lbs) bodyweight. It is important to follow the treatment recommendations as presented here. Do not deviate from the recommendations without the advice of your veterinary surgeon.

Dosage table:

	<b>Bodyweight (kg)</b>	<b>Tablets</b>
Puppies and small dogs	3-5	½
	6-10	1
Medium dogs	11-15	1½
	16-20	2
	21-25	2½
	26-30	3
Large dogs	31-35	3½
	36-40	4
	>40	1 tablet per 10 kg

For oral administration, the tablets can be given to the dog or disguised in food. No starvation is needed before, or after, treatment.

Puppies should be treated at 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 month intervals. It is advisable to treat the bitch at the same time as the puppies. Not for use in dogs weighing less than 3 kg.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every two weeks until weaning.

For routine worm control adult dogs should be treated every 3 months. For routine treatment a single dose is recommended.

In case of suspected heavy roundworm infestation, please contact your veterinary surgeon for diagnosis and treatment recommendation.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

The combination of praziquantel, pyrantel embonate and febantel is well tolerated in dogs. In safety studies, a single dose of 5 times the recommended dose or greater gave rise to occasional vomiting.

#### **4.11 Withdrawal period(s)**

Not applicable.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Anthelmintic, praziquantel combinations.  
ATC vet code: QP52AA51

#### **5.1 Pharmacodynamic properties**

This product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The product contains three active substances, as follows:

1. Febantel, a probenzimidazole
2. Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative
3. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative

In this fixed combination, pyrantel and febantel act against all relevant nematodes (ascarids, hookworms, and whipworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*.

This combination shows synergistic activity in the case of hookworms and febantel is effective against *T. vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular *Taenia* spp., *Dipylidium caninum*, *Echinococcus granulosus* and *Echinococcus multilocularis*. Praziquantel acts against all adult and immature forms of these parasites.

Praziquantel is very rapidly absorbed through the parasite's surface and distributed throughout the parasite. Both in vitro and in vivo studies have shown that praziquantel causes severe damage to the parasite integument, resulting in the contraction and paralysis of the parasites. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolization of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis of the nematodes and thereby allow removal from the gastrointestinal system by peristalsis.

Within the mammalian system, febantel undergoes ring closure, forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption of structures vital to the normal functioning of the helminth. Glucose uptake in particular is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2 – 3 days later.

## **5.2 Pharmacokinetic particulars**

Perorally administered praziquantel is absorbed almost completely from the intestinal tract. After absorption, the drug is distributed to all organs. Praziquantel is metabolized into inactive forms in the liver and secreted in bile. It is excreted within 24 hours to more than 95% of the administered dosage. Only traces of non-metabolised praziquantel are excreted.

Following administration of the product to dogs, peak plasma concentrations of praziquantel were achieved by approximately 2.5 hours.

The pamoate salt of pyrantel has low aqueous solubility, an attribute that reduces absorption from the gut and allows the drug to reach and be effective against parasites in the large intestine. Following absorption, pyrantel pamoate is quickly and almost completely metabolized into inactive metabolites that are excreted rapidly in the urine.

Febantel is absorbed relatively rapidly and metabolized to a number of metabolites including fenbendazole and oxfendazole, which have anthelmintic activity.

Following administration of the product to dogs, peak plasma concentrations of fenbendazole and oxfendazole were achieved by approximately 7-9 hours.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate,  
Microcrystalline cellulose,  
Magnesium stearate,  
Colloidal anhydrous silica,  
Croscarmellose sodium,  
Sodium laurilsulfate  
Pork flavour

### **6.2 Incompatibilities**

Not Applicable

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years  
Discard any unused divided tablets.

#### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

#### **6.5 Nature and composition of immediate packaging**

The product is presented in either:

Individual strips composed of aluminium foil 30 µm/30 gsm extruded polythene, containing 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20 tablets.

or

Individual blisters composed of 45 µm, soft temper aluminium foil and 25 µm hard temper aluminium foil, containing 2 or 8 tablets.

The strips or blisters are packed into cartons containing either 2,4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 70, 80, 84, 90, 98, 100, 104, 106, 120, 140, 150, 180, 200, 204, 206, 250, 280, 300, 500 or 1000 tablets.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

C&H Generics Ltd  
c/o Michael McEvoy and Co  
Seville House  
New Dock Street  
Galway  
Ireland

### **8. MARKETING AUTHORISATION NUMBER**

Vm 40162/4000


### **9. DATE OF FIRST AUTHORISATION**

15 March 2012

### **10. DATE OF REVISION OF THE TEXT**

August 2018

Approved: 17 August 2018

A handwritten signature in black ink that reads "D. Austin". The signature is written in a cursive style with a horizontal line extending to the right from the end of the name.