



Veterinary
Medicines
Directorate

United Kingdom
Veterinary Medicines Directorate
Woodham Lane
New Haw
Addlestone
Surrey KT15 3LS

DECENTRALISED PROCEDURE

**PUBLICLY AVAILABLE ASSESSMENT REPORT FOR A VETERINARY
MEDICINAL PRODUCT**

Qivitan LC 75 mg Intramammary Ointment for Lactating Cows

Date Created: July 2018

**PuAR correct as of July 2018, the RMS has now been transferred to IE.
Please contact the RMS for future updates**

MODULE 1

PRODUCT SUMMARY

EU Procedure number	UK/V/0641/001/DC
Name, strength and pharmaceutical form	Qivitan LC 75 Intramammary Ointment for Lactating Cattle
Applicant	Livisto Int'l S.L
Active substance(s)	Cefquinome 75 mg (as cefquinome sulfate 88.92 mg), in a syringe containing 8 g
ATC Vetcode	QJ51DE90
Target species	Cattle
Indication for use	For the treatment of clinical mastitis in the lactating cow caused by the following cefquinome-sensitive organisms: <i>Streptococcus uberis</i> , <i>Streptococcus dysgalactiae</i> , <i>Staphylococcus aureus</i> and <i>Escherichia coli</i> .

MODULE 2

The Summary of Product Characteristics (SPC) for this product is available on the Product Information Database of the Veterinary Medicines Directorate.

www.gov.uk/check-animal-medicine-licensed

MODULE 3

PUBLIC ASSESSMENT REPORT

Legal basis of original application	Generic 'hybrid' application in accordance with Article 13 (3) of Directive 2001/82/EC as amended.
Date of conclusion of the decentralised procedure	28 th March 2018.
Date product first authorised in the Reference Member State (MRP only)	Not applicable.
Concerned Member States for original procedure	Austria, Belgium, Croatia, Cyprus, France, Germany, Greece, Hungary, Ireland, Italy, The Netherlands, Poland, Portugal, Romania, Slovakia, Slovenia, Spain

I. SCIENTIFIC OVERVIEW

This was a generic 'hybrid' application, in accordance with Article 13 (3) of Directive 2001/82/EC, as amended. The product is locally acting, and therefore bioequivalence using plasma level of the active substance could not be demonstrated. A waiver from the requirement to conduct *in vivo* studies was permissible as outlined in the Guidance of conduct of efficacy studies for intramammary products for use in cattle (EMA/CVMP/EWP/14272/2011). The reference product is Cobacatan LC 75 mg Intramammary Ointment for Lactating Cows, marketed in the UK since March 1996.

The product is indicated for the treatment of clinical mastitis in the lactating cow caused by the cefquinome-sensitive organisms: *Streptococcus uberis*, *Streptococcus dysgalactiae*, *Staphylococcus aureus* and *Escherichia coli*.

The product is produced and controlled using validated methods and tests which ensure the consistency of the product released onto the market. It has been shown that the product can be safely used in the target species, any reactions observed are indicated in the SPC.¹ The product is safe for the user, the consumer of foodstuffs from treated animals and for the environment, when used as recommended. Suitable warnings and precautions are indicated in the SPC.

¹ SPC – Summary of product Characteristics.

The efficacy ² of the product was demonstrated according to the claims made in the SPC. The overall benefit/risk analysis is in favour of granting a marketing authorisation.

II. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE CONSTITUENTS

II.A. Composition

The product, (8 g in a syringe), contains 75 mg cefquinome (as 88.92 mg cefquinome sulfate). The excipients are white soft paraffin and liquid paraffin.

The container/closure system consists of the following:

Prefilled 8 g single-dose intramammary syringe consisting of white opaque LDPE barrel with white opaque LDPE plunger and white opaque LDPE cap.

Cleaning towels (smooth, white crepe paper impregnated with isopropyl alcohol/benzalkonium chloride) individually wrapped.

Cardboard boxes of 3 syringes and 3 cleaning towels.

Cardboard boxes of 12 syringes and 12 cleaning towels.

Cardboard boxes of 24 syringes and 24 cleaning towels.

Cardboard boxes of 36 syringes and 36 cleaning towels.

The choice of the formulation and the absence of preservative are justified. The product is an established pharmaceutical form and its development is adequately described in accordance with the relevant European guidelines.

II.B. Description of the Manufacturing Method

The product is manufactured fully in accordance with the principles of good manufacturing practice from a licensed manufacturing site. The manufacturing method consists of a simple addition, mixing and sterilisation process. The product is then packaged for sale.

Process validation data on the product have been presented in accordance with the relevant European guidelines.

II.C. Control of Starting Materials

The active substance is cefquinome, an established active for which the applicant provided specification. The active substance is manufactured in accordance with the principles of good manufacturing practice.

² Efficacy – The production of a desired or intended result.

The active substance specification is considered adequate to control the quality of the material. Batch analytical data demonstrating compliance with this specification have been provided. An acceptable Certificate of Analysis was provided.

Packaging complies with the monograph for the rubber stoppers as cited in the European Pharmacopoeia, and with Directive 1935/2004 (EC) for the tin pack. Excipients comply with monographs in the European Pharmacopoeia.

II.C.4. Substances of Biological Origin

No materials of animal origin are used during manufacture. A declaration of compliance with EMEA/410/01 has been provided.

II.D. Control Tests Carried Out at Intermediate Stages of the Manufacturing Process

Sterile micronised material is treated as intermediate product. Tests carried out on the immediate product are: appearance, identification of the active substance, colour and clarity of solution, pH, optical rotation, sulphated ash content, heavy metal content, water content, related substances, presence of endotoxins, sterility, assay of sulfate and particle size.

II.E. Control Tests on the Finished Product

The finished product specification controls the relevant parameters for the pharmaceutical form. The tests in the specification, and their limits, have been justified and are considered appropriate to adequately control the quality of the product. Satisfactory validation data for the analytical methods have been provided. Batch analytical data from the proposed production site have been provided demonstrating compliance with the specification. Control tests on the finished product include those for: appearance, particle size, deliverable mass, viscosity, identification of active substance, assay of active substance and related substances and sterility.

II.F. Stability

Stability data on the active substance and finished product have been provided in accordance with applicable European guidelines, demonstrating the stability of the active substance when stored under the approved conditions.

G. Other Information

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Do not store above 25°C.

III. SAFETY AND RESIDUES DOCUMENTATION (PHARMACO-TOXICOLOGICAL)

III.A Safety Documentation

Due to the legal basis of the application, there was no requirement to submit pharmacological and toxicological data. A user risk assessment (URA) and environmental risk assessment (ERA) were included. The proposed product and the reference product are identical with regards to composition, pharmaceutical form and strength.

User Safety

A user risk assessment was provided in compliance with the relevant guideline. The risk profile of the proposed product was considered to be the same as that of the reference product.

Warnings and precautions as listed on the product literature are adequate to ensure safety to users of the product. Therefore the following applicant's user recommendations are appropriate:

- When infusing the product, protective gloves should be worn to avoid skin contact.
- Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-sensitivity to cephalosporin and vice versa. Allergic reactions to these substances may occasionally be serious.
- Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.
- Handle this product with great care to avoid exposure, taking all recommended precautions.
- If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the Doctor this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.
- The cleaning towels provided with this product contain isopropyl alcohol and benzalkonium chloride, which may cause skin irritation in some people. It is recommended to wear protective gloves when using the towels.

Environmental Safety

The Environmental Risk Assessment (ERA) was carried out in accordance with VICH and CVMP guidelines.

The applicant has provided a Phase I assessment consisting of a description of the product, information on the pattern of use, and a VICH Phase I decision tree.

The $PEC_{soil\ initial}$ values of 1.7 $\mu\text{g}/\text{kg}$ (intensively reared dairy cattle), and 1.1 $\mu\text{g}/\text{kg}$ (pasture reared dairy cattle) are below the trigger value (100 $\mu\text{g}/\text{kg}$). As such, the assessment can end at Phase I (Question 17 of the VICH decision tree) as exposure of the environment is not extensive. The SPC and product literature carry suitable warnings.

III.B.2 Residues documentation

Residue Studies

The formulation for this product is sufficiently similar to that of the reference product such that residue depletion data are not required and the withdrawal periods of the reference product apply.

Pharmacologically active substance(s)	Marker residue	Animal species	MRLs ($\mu\text{g}/\text{kg}$)	Target tissues
Cefquinome	Cefquinome	Bovine	50 $\mu\text{g}/\text{kg}$ 50 $\mu\text{g}/\text{kg}$ 100 $\mu\text{g}/\text{kg}$ 200 $\mu\text{g}/\text{kg}$ 20 $\mu\text{g}/\text{kg}$	Muscle Fat Liver Kidney Milk

Withdrawal Periods

Meat and offal: 4 days
Milk: 5 days (120 hours).

IV CLINICAL DOCUMENTATION

IV.1. Pre-Clinical Studies

Pharmacology

Pharmacokinetics

No data required.

Pharmacokinetics

A biowaiver was established under Guideline EMA/CVMP/EWP/141272/2011, which states that:

'Efficacy and tolerance studies may be waived for generic products that are fully identical to the reference product.

Efficacy studies may also be waived if the following conditions are fulfilled: the proposed product has the same pharmaceutical form and contains qualitatively and quantitatively the same active substance(s) (same salts), the excipients of the generic are qualitatively and quantitatively very similar compared to the reference product, and the physicochemical properties (e.g. crystalline form, particle size distribution, viscosity, relative density, dissolution profile) of the generic product are similar to those of the reference product. Local tolerance data might be requested'.

Any differences between the formulations of the proposed and reference products were not expected to alter the disposition of the active substance. No further data were required.

Tolerance in the Target Species

No data were required due to the nature of the application.

Resistance

Adequate warnings and precautions appear on the product literature.

IV.II. Clinical Documentation

No data were required due to the nature of the application.

V OVERALL CONCLUSION AND BENEFIT– RISK ASSESSMENT

The data submitted in the dossier demonstrate that when the product is used in accordance with the Summary of Product Characteristics the benefit/risk profile of the product(s) is favourable.

MODULE 4

POST-AUTHORISATION ASSESSMENTS

The SPC and package leaflet may be updated to include new information on the quality, safety and efficacy of the veterinary medicinal product. The current SPC is available on the Product Information Database of the Veterinary Medicines Directorate website.

www.gov.uk/check-animal-medicine-licensed

The post-authorisation assessment (PAA) contains information on significant changes which have been made after the original procedure which are important for the quality, safety or efficacy of the product.

The PAA for this product is available on the Product Information Database of the Veterinary Medicines Directorate website.

www.gov.uk/check-animal-medicine-licensed