

ANNEX

Identification number of the additive	Name of the holder of authorisation	Additive (Trade name)	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content		Maximum content	Other provisions	End of period of authorisation	Provisional maximum residue limits (MRLs) in the relevant foodstuffs of animal origin
						mg of active substance/kg of complete feedstuff with a moisture content of 12 %	mg of active substance/kg of complete feedstuff with a moisture content of 12 %				
Coccidiostats and histomonostats											
5 1 701	Huvepharma NV Belgium	Monensin sodium (Coxidin)	<p><i>Active substance</i> $C_{38}H_{61}O_{11}Na$ Sodium salt of polyether monocarboxylic acid produced by <i>Streptomyces cinnamonensis</i>, 28682 LMG S-19095 in powder form.</p> <p><i>Factor composition:</i> Monensin A: not less than 90 % Monensin A + B: not less than 95 % Monensin C: 0,2-0,3 %</p> <p><i>Additive composition</i> Monensin sodium technical substance equivalent to monensin activity: 25 % Perlite: 15-20 % Wheat bran: 55-60 %</p> <p><i>Analytical method (1)</i> Method for determination of the active substance: high performance liquid chromatography (HPLC) with post-column derivatisation and UV detection ($\lambda = 520 \text{ nm}$)</p>	Chickens for fattening Turkeys	— 16 weeks	100 60	125 100	<p>1. Use prohibited at least one day before slaughter</p> <p>2. The additive shall be incorporated in compound feedstuffs in form of a premixture</p> <p>3. Maximum permitted dose of monensin sodium in complementary feedstuffs: — 625 mg/kg for chickens for fattening; — 500 mg/kg for turkeys</p> <p>4. Monensin sodium shall not be mixed with other coccidiostats</p> <p>5. Indicate in the instructions for use: 'Dangerous for equines. This feedstuff contains an ionophore: avoid simultaneous administration with tiamulin and monitor for possible adverse reactions when used concurrently with other medicinal substances'</p> <p>6. Wear suitable protective clothing, gloves and eye/face protection. In case of insufficient ventilation in the premise, wear suitable respiratory equipments</p>	6.2.2017	25 µg monensin sodium/kg of wet skin + fat 8 µg monensin sodium/kg of wet liver, wet kidney and wet muscle	

(1) Details of the analytical methods are available at the following address of the Community Reference Laboratory: www.irmm.jrc.be/crl-feed-additives