

## PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery+Parvo+Lepto suspension for injection

### 2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

50 ml, 100 ml and 250 ml presentations:

Each 2 ml dose contains inactivated strains of:

*Erysipelothrix rhusiopathiae*, serotype 2 (strain M2)  $\geq 1$  ppd; Porcine parvovirus (strain 014)  $\geq 130$  U; *Leptospira interrogans* serogroup Canicola serovar Portland-vero (strain Ca-12-000)  $\geq 2816$  U; *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni (strain Ic-02-001)  $\geq 210$  U; *L. interrogans* serogroup Australis serovar Bratislava (strain As-05-073)  $\geq 1310$  U; *L. kirschneri* serogroup Grippotyphosa serovar Dadas (strain Gr-01-005)  $\geq 648$  U; *L. interrogans* serogroup Pomona serovar Pomona (strain Po-01-000)  $\geq 166$  U; *L. santarosai* serogroup Tarassovi serovar Gatuni (strain S1148/02)  $\geq 276$  U

Adjuvant: dl- $\alpha$ -tocopheryl acetate 150 mg

20 ml presentation:

Each 2 ml dose contains inactivated strains of:

*Erysipelothrix rhusiopathiae*, serotype 2  $\geq 1$  ppd; Porcine parvovirus  $\geq 130$  U; *Leptospira interrogans* serogroup Canicola serovar Portland-vero  $\geq 2816$  U; *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni  $\geq 210$  U; *L. interrogans* serogroup Australis serovar Bratislava  $\geq 1310$  U; *L. kirschneri* serogroup Grippotyphosa serovar Dadas  $\geq 648$  U; *L. interrogans* serogroup Pomona serovar Pomona  $\geq 166$  U; *L. santarosai* serogroup Tarassovi serovar Gatuni  $\geq 276$  U

Adjuvant: dl- $\alpha$ -tocopheryl acetate 150 mg

### 3. PACKAGE SIZE

20 ml (10 doses)

10 x 20 ml (10 doses)

50 ml (25 doses)

10 x 50 ml (25 doses)

100 ml (50 doses)

250 ml (125 doses)

**4. TARGET SPECIES**

Pigs for reproduction.

**5. INDICATION(S)**

**6. ROUTES OF ADMINISTRATION**

Intramuscular use.

**7. WITHDRAWAL PERIODS**

Withdrawal period: Zero days.

**8. EXPIRY DATE**

EXP {mm/yyyy}

Once broached use within 10 hours.

**9. SPECIAL STORAGE PRECAUTIONS**

Store in a refrigerator.

Do not freeze.

Protect from light.

Keep the vial in the outer carton.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Ltd.  
Walton Manor, Walton  
Milton Keynes  
MK7 7AJ

**14. MARKETING AUTHORISATION NUMBERS**

Vm 01708/5084

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

Accidental injection is dangerous.

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-V To be supplied only on veterinary prescription.

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**PET vials (50, 100 and 250 ml)**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Porcilis Ery+Parvo+Lepto suspension for injection

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Inactivated *Erysipelothrix rhusiopathiae*, Porcine parvovirus and *Leptospira*

**3. TARGET SPECIES**

Pigs for reproduction.

**4. ROUTES OF ADMINISTRATION**

Intramuscular use.

**5. WITHDRAWAL PERIODS**

Withdrawal period: Zero days.

**6. EXPIRY DATE**

EXP {mm/yyyy}

Once broached use within 10 hours.

**7. SPECIAL STORAGE PRECAUTIONS**

Keep the vial in the outer carton.

Store in a refrigerator.

Do not freeze.

Protect from light.

**8. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Ltd.  
Walton Manor, Walton  
Milton Keynes  
MK7 7AJ, UK

**9. BATCH NUMBER**

Lot {number}

**10. PACKAGE SIZE**

50 ml (25 doses)  
100 ml (50 doses)  
250 ml (125 doses)

**11. INDICATION(S)**

**12. SPECIAL WARNING(S), IF NECESSARY**

Accidental injection is dangerous.

**13. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**14. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

For animal treatment only.  
To be supplied only on veterinary prescription.  
POM-V

**15. MARKETING AUTHORISATION NUMBER(S)**

Vm 01708/5084

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**PET vials (20 ml)**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Porcilis Ery+Parvo+Lepto



**2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCE(S)**

Inactivated *Erysipelothrix rhusiopathiae*, Porcine parvovirus and *Leptospira*

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

EXP {mm/yyyy}

Once broached use within 10 hours.

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

20 ml (10 doses)

**6. ROUTE(S) OF ADMINISTRATION**

IM use

**7. WITHDRAWAL PERIOD**

Withdrawal period: Zero days.

**8. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

POM-V

## PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery+Parvo+Lepto suspension for injection for pigs

### 2. COMPOSITION

Each 2 ml dose contains:

#### Active substances:

Inactivated strains of:

<i>Erysipelothrix rhusiopathiae</i> , serotype 2 (strain M2)	≥ 1 ppd <sup>1</sup>
Porcine parvovirus (strain 014)	≥ 130 U <sup>2</sup>
<i>Leptospira interrogans</i> serogroup Canicola serovar Portland-vere (strain Ca-12-000)	≥ 2816 U <sup>2</sup>
<i>Leptospira interrogans</i> serogroup Icterohaemorrhagiae serovar Copenhageni (strain Ic-02-001)	≥ 210 U <sup>2</sup>
<i>Leptospira interrogans</i> serogroup Australis serovar Bratislava (strain As-05-073)	≥ 1310 U <sup>2</sup>
<i>Leptospira kirschneri</i> serogroup Grippotyphosa serovar Dadas (strain Gr-01-005)	≥ 648 U <sup>2</sup>
<i>Leptospira interrogans</i> serogroup Pomona serovar Pomona (strain Po-01-000)	≥ 166 U <sup>2</sup>
<i>Leptospira santarosai</i> serogroup Tarassovi serovar Gatuni (strain S1148/02)	≥ 276 U <sup>2</sup>

#### Adjuvant:

dl-α-tocopheryl acetate	150 mg
-------------------------	--------

<sup>1</sup>Pig protective dose as compared to a reference preparation known to be protective in pigs.

<sup>2</sup>As determined in the *in vitro* antigenic mass ELISA potency test.

Homogenous white to nearly white suspension after shaking.

### 3. TARGET SPECIES

Pigs for reproduction.

### 4. INDICATIONS FOR USE

For the active immunisation of pigs:

- to reduce clinical signs (skin lesions and fever) of swine erysipelas caused by *Erysipelothrix rhusiopathiae*, serotype 1 and serotype 2.
- to reduce transplacental infection, viral load and fetal mortality caused by Porcine parvovirus.

- to reduce clinical signs (increase of body temperature and reduction in feed intake or activity), infection and bacterial excretion caused by *L. interrogans* serogroup Canicola serovar Canicola.
- to reduce clinical signs (increase of body temperature and reduction in feed intake or activity), severity of infection and foetal mortality caused by *L. interrogans* serogroup Pomona serovar Pomona.
- to reduce infection caused by *L. interrogans* serogroup Icterohaemorrhagiae serovars Copenhageni and Icterohaemorrhagiae, *L. interrogans* serogroup Australis serovar Bratislava, *L. kirschneri* serogroup Grippotyphosa serovars Grippotyphosa and Bananal/Liangguang, *L. weillii* serogroup Tarassovi serovar Vughia and *L. borgpetersenii* serogroup Tarassovi serovar Tarassovi.

Onset of immunity:

*E. rhusiopathiae*: 3 weeks

Porcine parvovirus: 10 weeks

*Leptospira* serogroups: 2 weeks

Duration of immunity:

*E. rhusiopathiae*: 6 months

Porcine parvovirus: 1 year

*Leptospira* serogroup Australis: 6 months

*Leptospira* serogroups Canicola, Icterohaemorrhagiae, Grippotyphosa, Pomona and Tarassovi: 1 year

## 5. CONTRAINDICATIONS

None.

## 6. SPECIAL WARNING(S)

For animal treatment only.

Special warnings:

Vaccinate healthy animals only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package insert or label to the physician.

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Interaction with other medicinal products and other forms of interactions:

No information is available on the safety and efficacy of this vaccine with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

#### Overdose:

No adverse events other than those mentioned in section “Adverse events” were observed after the administration of a double dose of vaccine.

#### Major incompatibilities:

Do not mix with any other veterinary medicinal product.

## **7. ADVERSE EVENTS**

Pigs for reproduction:

Very common (>1 animal / 10 animals treated):	Elevated temperature <sup>1</sup> Injection site swelling <sup>2</sup>
Uncommon (1 to 10 animals / 1,000 animals treated):	Decreased activity <sup>3</sup> , reduced food intake <sup>3</sup>
Rare (1 to 10 animals / 10,000 animals treated):	Vomiting <sup>4</sup> , reddening of the skin <sup>4</sup> , tachypnoea (rapid breathing) <sup>4</sup> , twitching <sup>4</sup>
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Hypersensitivity reaction.

<sup>1</sup>The observed mean increase was 0.5 °C (in individual cases the maximum increase was 1.5 °C) up until 2 days after vaccination.

<sup>2</sup>Local reactions, mostly consisting of red, mild to hard, non-painful swellings are a very common observation. In general, local reactions may have a diameter of ≤ 5 cm, in very rare cases local reactions in individual animals can be up to 20 cm in diameter. All local reactions disappear completely within approximately 2 weeks after vaccination.

<sup>3</sup>Feed intake and activity are completely restored within a week.

<sup>4</sup>Intermediate systemic reactions resolve in a few minutes.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system:

e-mail: [adverse.events@vmd.gov.uk](mailto:adverse.events@vmd.gov.uk)

<https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For intramuscular use. Administer a single dose of 2 ml in the neck region.

Basic vaccination scheme: Pigs which have not yet been vaccinated shall be given a primary injection 6 to 8 weeks before the expected date of insemination and a booster injection 4 weeks later.

Revaccination: A single revaccination with the veterinary medicinal product should be given once a year. Six months post each vaccination with the veterinary medicinal product, a single revaccination with an *Erysipelothrix rhusiopathiae* containing product should be given to maintain immunity against *Erysipelothrix rhusiopathiae*. In case of known infection pressure with *L. interrogans* serogroup Australis, a single revaccination with the veterinary medicinal product should be given every six months, as it is unknown if or for how long the duration of immunity for this serogroup persists beyond six months.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Before use allow the vaccine to reach room temperature.  
Shake well before use.  
Avoid introduction of contamination by multiple broaching.

## **10. WITHDRAWAL PERIOD(S)**

Zero days.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep the vial in the outer carton.  
Keep out of the sight and reach of children.

Store in a refrigerator (2 °C – 8 °C).  
Do not freeze.  
Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.  
Shelf life after first opening the immediate packaging: 10 hours.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater.  
Ask your veterinary surgeon how to dispose of medicines no longer required.  
These measures should help to protect the environment.

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

#### 14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 01708/5084

Pack sizes:

Cardboard box with 1 vial of 20 ml.  
Cardboard box with 10 vials of 20 ml.  
Cardboard box with 1 vial of 50 ml.  
Cardboard box with 10 vials of 50 ml.  
Cardboard box with 1 vial of 100 ml.  
Cardboard box with 1 vial of 250 ml.

Not all pack sizes may be marketed.

#### 15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED

October 2023

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk).

#### 16. CONTACT DETAILS

Marketing authorisation holder:

MSD Animal Health UK Ltd.  
Walton Manor, Walton  
Milton Keynes  
Buckinghamshire  
MK7 7AJ, UK

Manufacturer responsible for batch release:

Intervet International B.V.  
Wim de Körverstraat 35  
5831 AN Boxmeer  
The Netherlands

Contact details to report suspected adverse reactions:

MSD Animal Health UK Ltd.  
Tel.: +44 (0)1908 685685

#### 17. OTHER INFORMATION

☒ POM-V To be supplied only on veterinary prescription.

Approved 05 October 2023

