

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Deosect® 5% w/v Concentrate for Cutaneous Spray Solution

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Deosect 5% w/v Concentrate for Cutaneous Spray Solution contains cypermethrin technical (cis 50:trans 50) 5% w/v (0.1% w/v when diluted).

3. PHARMACEUTICAL FORM

Concentrate for Cutaneous Spray Solution

4. PACKAGE SIZE

250 ml
1 Litre

5. TARGET SPECIES

Horses

6. INDICATION(S)

Monthly general fly control

Topical spray to control flies and lice on horses

7. METHOD AND ROUTE(S) OF ADMINISTRATION

For use with sponge or handsprayer

Directions for use:

Deosect must be diluted and mixed with water to make a solution just before use. A knapsack, hand sprayer or sponge may be used to apply the solution.

The diluted solution should be used within 24 hours.

The following amounts should be used to give a 1:50 dilution.

Deosect	Water
10 ml	500 ml
20 ml	1 Litre
100 ml	5 Litres
200 ml	10 Litres

A medium fine spray is advisable to avoid fogging of the diluted Deosect.

Careful attention must be paid to total animal coverage, especially the face, legs, back and hindquarters.

The following amounts should be used.

Horses: 500 ml of diluted solution per animal.

Ponies: 125 ml of diluted solution per animal.

Fly Control: The application of the solution should commence at the onset of the fly season and be repeated monthly for general fly control and every two weeks for severe head and face fly control.

Lice Control: Apply when lice are observed and repeat as necessary at a minimum interval of 14 days.

8. WITHDRAWAL PERIOD

Not to be used in animals intended for human consumption. Treated horses may never be slaughtered for human consumption. The horse must have been declared as not intended for human consumption under national horse passport legislation.

9. SPECIAL WARNING(S), IF NECESSARY

For external use only.

Avoid spraying directly into animal's eyes. Do not apply to any lesions where the skin is broken.

Do not spray in or near milking parlours or where milking machinery may be contaminated.

This product belongs to the pyrethroid group. Possible undesirable effects can be irritation and allergic reactions on the skin.

Deosect can be safely used during pregnancy and lactation.

Operator warnings:

FLAMMABLE. Wear impermeable gloves, face shield and rubber boots when handling the concentrate, diluted liquid and when spraying animals.

Use in a well-ventilated area and avoid inhaling the spray mist. If splashed in the eye or on the skin, wash with plenty of clean, running water immediately and seek medical attention.

Remove contaminated clothing immediately and wash before re-use.

Do not smoke or eat while handling the product.

Wash hands and exposed skin after handling this product and before meals.

10. EXPIRY DATE

EXP: MM/YYYY

11. SPECIAL STORAGE CONDITIONS

Store in original container, tightly closed, in a safe place away from food, drink and animal feedstuffs.

Do not store above 25°C. Protect from frost.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal:

DANGEROUS to fish and crustaceans. Do not contaminate ponds, waterways or ditches with the product or used containers. Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

For animal treatment only.

POM-VPS

To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Zoetis UK Limited

1st Floor, Birchwood Building

Springfield Drive

Leatherhead

Surrey

KT22 7LP

16. MARKETING AUTHORISATION NUMBER(S)

Vm 42058/5230

17. MANUFACTURER'S BATCH NUMBER

LOT:

Gavin Hall
Approved: 09 December 2025