PARTICULARS TO APPEAR ON THE OUTER PACKAGE – Cardboard box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Vivitonin 100 mg tablets

2. STATEMENT OF ACTIVE SUBSTANCES

Each tablet contains:
Propentofylline 100 mg

3. PACKAGE SIZE

60 film-coated tablets

4. TARGET SPECIES

Dogs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Oral use.
Must be given at least 30 minutes before food.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Do not store above 25 °C.
Store in a dry place.
Keep the blister packs in the outer carton.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Intervet International B.V.

14. MARKETING AUTHORISATION NUMBER

Vm 06376/4078

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS Blister packs

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Vivitonin 100 mg tablets

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Propentofylline 100 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Vivitonin 100 mg tablets for dogs

2. Composition

Each tablet contains:

Active substance:
Propentofylline 100 mg

Ochre, biconvex and oblong film-coated tablets with a breaking line on one side and “K100” debossed on both sides of the breaking line.

3. Target species

Dogs.

4. Indications for use

For improvement in dullness, lethargy and overall demeanour in larger dogs. The veterinary medicinal product is particularly useful in older dogs, where it may increase willingness to exercise and exercise tolerance.
The active ingredient, propentofylline has been shown to increase blood flow, particularly of the heart and skeletal muscle. It also increases the blood flow of the brain and therefore its oxygen supply, without increasing the brain’s glucose demand. Propentofylline has a modest positive chronotropic effect and a marked positive inotropic effect. In addition, it has been shown to have an antiarrhythmic effect in dogs with myocardial ischemia and a bronchodilator action equivalent to that of aminophylline.
Propentofylline inhibits platelet aggregation and improves the flow properties of erythrocytes. It has a direct effect on the heart and reduces peripheral vascular resistance thereby lowering cardiac load.

5. Contraindications

Do not administer to pregnant bitches or breeding animals.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

6. Special warnings

None.

Special precautions for safe use in the target species:
Specific diseases (e.g. kidney disease) should be treated accordingly.
In the case of renal failure, the dose should be reduced.
Consideration should be given to rationalising the medication of dogs already receiving treatment for congestive heart failure or bronchial disease.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:
Care should be taken to avoid accidental ingestion.
Wash hands after use.

Pregnancy:
Do not use in pregnant bitches. The safety of the veterinary medicinal product has not been established during pregnancy.

Fertility:
Do not use in breeding animals.

Interaction with other medicinal products and other forms of interaction:
None known.

Overdose:
Symptoms of cardiac and cerebral overstimulation have been observed. In such cases, animals should be treated symptomatically.

Major incompatibilities:
None known.

7. Adverse events

Dogs:

Rare (1 to 10 animals / 10,000 animals treated):	Vomiting[1]; Allergic reaction (e.g., Urticaria)[2].

[1] Particularly at the commencement of therapy.
[2] Necessitates discontinuation of treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at: Website: https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine

e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, routes and method of administration

Oral use.

Half a tablet per 10 kg body weight twice a day.
The 100 mg tablets should not be quartered.

More accurate dosing may be achieved using a combination of Vivitonin 100 mg and Vivitonin 50 mg tablets. Dogs of less than 20 kg can be given Vivitonin 50 mg tablets.

Divide the tablets in halves with a knife or with a tablet splitter.

9. Advice on correct administration

The tablets can be administered directly onto the back of the dog's tongue or can be mixed in a small ball of food and should be administered at least 30 minutes before feeding.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Do not store above 25 °C.
Store in a dry place.
Keep the blister packs in the outer carton.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater.
Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.
Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

Vm 06376/4078

Pack size:
Cardboard box containing 60 tablets, presented in 6 blister strips of 10 tablets**.**

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder:
Intervet International B.V.
Wim de Körverstraat 35
5831 AN Boxmeer
Netherlands

Manufacturer responsible for batch release:
Intervet GesmbH
Siemensstrasse 107
1210 Vienna
Austria

Local representative:
MSD Animal Health UK Limited
Walton Manor, Walton
Milton Keynes, MK7 7AJ, United Kingdom

Contact details to report suspected adverse reactions:
UK(GB)
MSD Animal Health UK Ltd.
Tel.: +44 (0)1908 685685

UK(NI)
Intervet Ireland Ltd.
Tel.: +353 (0)1 2970220

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information

POM-V Veterinary medicinal product subject to prescription.

Gavin Hall
Approved: 15 October 2025