PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE - 2.2 L Bottle Front/Back Label

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Crovect 1.25% w/v Pour-On Solution for Sheep

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Contains 1.25% w/v cypermethrin (tech.)

(cis:trans/ 80:20).

3. PHARMACEUTICAL FORM

Pour-on

4. PACKAGE SIZE

0.8L
2.2L
5L

5. TARGET SPECIES

Sheep

6. INDICATION(S)

Blowfly Strike: Treatment and Prevention for 6-8 weeks
Ticks: Treatment of tick infestation with a persistence efficacy of 10 weeks
Headflies: Treatment and Control
Biting Lice: Treatment

<Front Label>
10 WEEK TICK CLAIM
Pictogram of Blowfly, Headfly, Tick and Lice

7. METHOD AND ROUTE(S) OF ADMINISTRATION

This product is administered by means of a Crovect Pour-On Gun.

FOR EXTERNAL USE ONLY.

Crovect Pour-On must be applied only with the Crovect Pour-On Gun as the product may have a detrimental effect on certain components of conventional dosing guns. Full operating and maintenance instructions, including details on the use of nozzles, are supplied with each Pour-On Gun. Please read carefully before use. Maintain gun carefully to ensure accurate dosage.

On completion of each treatment cycle of the product and before the dosing gun is stored, immediately expel any remaining product in the dosing gun.
Failure to use the product as directed may result in suboptimal parasite control.

Mothering up
Crovect Pour-On minimises mothering-up problems after treatment. However, care must be taken not to apply the product right to the lamb's tail, as a ewe recognises a young lamb partly by the smell of the tail area.
This product contains a blue dye to aid in the identification of treated animals. The colour is temporary and will fade from the fleece within 48 hours of application.

<Front Label>
Contains a blue dye to aid in the identification of treated sheep

Dosage:
BLOWFLIES
Prevention of blowfly strike

<table>

Bodyweight	Doses	Doses per pack
Up to 25 kg*	20 ml	110
25-40 kg	30 ml	73
Over 40 kg	40 ml	55

*Do not administer to animals of less than 12.5 kg bodyweight.

Apply as a fan-spray to the surface of the fleece on the back and hindquarters of the sheep using the Crovect Pour-On Gun fitted with the FAN-SPRAY NOZZLE. Half the dose should be applied to the shoulders, back and flanks, and half to the rump. The distance between the nozzle and the fleece should be approximately 20 cm. Each dose will require 2 or 3 sweeps to apply.
THE PRODUCT WILL PREVENT BLOWFLY STRIKE ONLY ON AREAS COVERED BY SPRAY. One application will give 6-8 weeks protection. Re-treat as required.

Treatment of blowfly strike
Apply directly to all the affected parts at the rate of 2.5 ml per 100-150 cm2 (roughly the size of a hand). Use the Crovect Pour-On Gun with the T-BAR NOZZLE. Most strikes will require 5-10 ml. Ensure that all affected parts are treated. A single application is sufficient to ensure that larvae are expelled and killed within a few hours.

<Diagram showing product application>

Further Information:
Blowflies are attracted by dirty, damp wool or open wounds. The most common site for fly strike is the rump due to soiling caused by diarrhoea and urine staining. The efficacy of Crovect our-On is reduced in the presence of soiled or dirty wool. Therefore, sheep must be crutched (dagged) regularly and appropriate worm control measures employed.

LICE
5 ml per 20 kg bodyweight up to a maximum of 20 ml.
<table>

Bodyweight	Doses	Doses per pack
Up to 20 kg	5 ml	440
21-40 kg	10 ml	220
41-60 kg	15 ml	146
Over 60 kg	20 ml	110

Using the STRAIGHT NOZZLE on the Crovect Pour-On Gun, apply as a pin stream from the shoulders to the rump along the middle of the backline. Sheep may be treated off-shears or at any time during the year.

<Diagram showing correct nozzle to use>

TICKS
Adult sheep and lambs over 10 kg: 10 ml per 20 kg bodyweight up to a maximum of 40 ml. One treatment will give 10 weeks cover.
Lambs under 10 kg: 5 ml followed 3 weeks later by a 10 ml application.

<table>

Bodyweight	Doses	Doses per pack
Up to 10 kg	5 ml	440
11-20 kg	10 ml	220
21-40 kg	20 ml	110
41-60 kg	30 ml	73
Over 60 kg	40 ml	55

Using the STRAIGHT NOZZLE on the Crovect Pour-On Gun, apply as a pin stream from the crown of the head to the top of the rump.

<Diagram showing correct nozzle to use>

HEADFLIES
5 ml per animal irrespective of size.

Apply to the top of the head between the ears using the Crovect Pour-On Gun with the T-BAR NOZZLE. Take care not to apply product in the sheep's eyes. One application of Crovect Pour-On before the start of the headfly season will give up to 4 weeks protection. Re-treat as required.

<Diagram showing correct nozzle to use>

8. WITHDRAWAL PERIOD

Meat and offal: - 8 days

Milk - 5 days.

9. SPECIAL WARNING(S), IF NECESSARY

Not to be used on lambs less than one week old.

Avoid treating very young lambs if the weather is unseasonably hot.

Take care not to apply product in the sheep's eyes.

This precaution should be observed particularly when treating breeds with little wool on their heads.

Prior to application, check the Pour-On gun for correct operation and calibration.

Overdose

Toxic signs in mammals are tremors, hyperexcitability, salivation, abnormal involuntary movements and paralysis. The signs disappear rapidly and the animals recover, generally within 1 week.

There is no specific antidote but symptomatic therapy can be given if considered necessary.

Operator Warnings

Make sure when attaching the Crovect Pour-On gun onto the container that the coiled retainer is secured onto the cap and the applicator. Wear eye protection, protective clothing, rubber gloves and boots when applying the product.

Wear a disposable face-mask when applying as a fan-spray for the prevention of blowfly-strike.

Remove contaminated clothing immediately and wash exposed skin with plenty of clean water. Wash splashes from skin and eyes immediately with plenty of clean water. If irritation persists, seek medical advice immediately.

Do not eat, drink or smoke whilst using the product.

Cypermethrin can be skin sensitizing and eye irritating. In a few instances operators may experience a mild tingling sensation of the skin when using the product. Wash splashes from skin immediately and thoroughly with plenty of clean water. If irritation persists seek medical advice immediately. If product splashes in eyes, flush thoroughly with water for 10 to 15 minutes and seek medical advice immediately

<Back Label - Diagram of bottle closure>
Attach tube to spouted cap provided. Replace original cap after use.

Environmental Warnings

EXTREMELY DANGEROUS to fish and other aquatic life. Do not contaminate ponds, water-ways or ditches with the product or empty container. As a precautionary measure sheep must be kept away from watercourses for at least one hour following treatment.

10. EXPIRY DATE

Exp:

11. SPECIAL STORAGE CONDITIONS

Pharmaceutical Precautions

Store in tightly closed original container. The product must be used within 3 months of opening the container. Discard unused material. Do not store above 25˚C. Protect from direct sunlight. Store away from food, drink and animal feedstuffs.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

[Distribution category]

FOR ANIMAL TREATMENT ONLY.
FOR EXTERNAL USE ONLY.

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION

POM-VPS products may only be supplied in accordance with a prescription from a Registered Qualified Person (RQP) as follows: (i) a registered veterinary surgeon; (ii) a registered pharmacist, or (iii) a registered suitably qualified person (SQP).

POM-VPS

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

KEEP OUT OF REACH AND SIGHT OF CHILDREN.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Elanco GmbH
Heinz-Lohmann Strasse 4
D-27472 Cuxhaven
Groden
Germany

16. MARKETING AUTHORISATION NUMBER(S)

Vm 52127/5079

17. MANUFACTURER’S BATCH NUMBER

Bn N˚.:

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE - 5 L Bottle Front/Back Label

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Crovect 1.25% w/v Pour-On Solution for Sheep

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Contains 1.25% w/v cypermethrin (tech.)

(cis:trans/ 80:20).

3. PHARMACEUTICAL FORM

Pour-on

4. PACKAGE SIZE

0.8L
2.2L
5L

5. TARGET SPECIES

Sheep

6. INDICATION(S)

Blowfly Strike: Treatment and Prevention for 6-8 weeks
Ticks: Treatment of tick infestation with a persistence efficacy of 10 weeks
Headflies: Treatment and Control
Biting Lice: Treatment

<Front Label>
10 WEEK TICK CLAIM
Pictogram of Blowfly, Headfly, Tick and Lice

7. METHOD AND ROUTE(S) OF ADMINISTRATION

This product is administered by means of a Crovect Pour-On Gun.

FOR EXTERNAL USE ONLY.

Crovect Pour-On must be applied only with the Crovect Pour-On Gun as the product may have a detrimental effect on certain components of conventional dosing guns. Full operating and maintenance instructions, including details on the use of nozzles, are supplied with each Pour-On Gun. Please read carefully before use. Maintain gun carefully to ensure accurate dosage.

On completion of each treatment cycle of the product and before the dosing gun is stored, immediately expel any remaining product in the dosing gun.
Failure to use the product as directed may result in suboptimal parasite control.

Mothering up
Crovect Pour-On minimises mothering-up problems after treatment. However, care must be taken not to apply the product right to the lamb's tail, as a ewe recognises a young lamb partly by the smell of the tail area.
This product contains a blue dye to aid in the identification of treated animals. The colour is temporary and will fade from the fleece within 48 hours of application.

<Front Label>
Contains a blue dye to aid in the identification of treated sheep

Dosage:
BLOWFLIES
Prevention of blowfly strike

<table>

Bodyweight	Doses	Doses per pack
Up to 25 kg*	20 ml	250
25-40 kg	30 ml	166
Over 40 kg	40 ml	125

*Do not administer to animals of less than 12.5 kg bodyweight.

Apply as a fan-spray to the surface of the fleece on the back and hindquarters of the sheep using the Crovect Pour-On Gun fitted with the FAN-SPRAY NOZZLE. Half the dose should be applied to the shoulders, back and flanks, and half to the rump. The distance between the nozzle and the fleece should be approximately 20 cm. Each dose will require 2 or 3 sweeps to apply.
THE PRODUCT WILL PREVENT BLOWFLY STRIKE ONLY ON AREAS COVERED BY SPRAY. One application will give 6-8 weeks protection. Re-treat as required.

Treatment of blowfly strike
Apply directly to all the affected parts at the rate of 2.5 ml per 100-150 cm2 (roughly the size of a hand). Use the Crovect Pour-On Gun with the T-BAR NOZZLE. Most strikes will require 5-10 ml. Ensure that all affected parts are treated. A single application is sufficient to ensure that larvae are expelled and killed within a few hours.

<Diagram showing product application>

Further Information:
Blowflies are attracted by dirty, damp wool or open wounds. The most common site for fly strike is the rump due to soiling caused by diarrhoea and urine staining. The efficacy of Crovect our-On is reduced in the presence of soiled or dirty wool. Therefore, sheep must be crutched (dagged) regularly and appropriate worm control measures employed.

LICE
5 ml per 20 kg bodyweight up to a maximum of 20 ml.
<table>

Bodyweight	Doses	Doses per pack
Up to 20 kg	5 ml	1000
21-40 kg	10 ml	500
41-60 kg	15 ml	333
Over 60 kg	20 ml	250

Using the STRAIGHT NOZZLE on the Crovect Pour-On Gun, apply as a pin stream from the shoulders to the rump along the middle of the backline. Sheep may be treated off-shears or at any time during the year.

<Diagram showing correct nozzle to use>

TICKS
Adult sheep and lambs over 10 kg: 10 ml per 20 kg bodyweight up to a maximum of 40 ml. One treatment will give 10 weeks cover.
Lambs under 10 kg: 5 ml followed 3 weeks later by a 10 ml application.

<table>

Bodyweight	Doses	Doses per pack
Up to 10 kg	5 ml	1000
11-20 kg	10 ml	500
21-40 kg	20 ml	250
41-60 kg	30 ml	166
Over 60 kg	40 ml	125

Using the STRAIGHT NOZZLE on the Crovect Pour-On Gun, apply as a pin stream from the crown of the head to the top of the rump.

<Diagram showing correct nozzle to use>

HEADFLIES
5 ml per animal irrespective of size.
Apply to the top of the head between the ears using the Crovect Pour-On Gun with the T-BAR NOZZLE. Take care not to apply product in the sheep's eyes. One application of Crovect Pour-On before the start of the headfly season will give up to 4 weeks protection. Re-treat as required.

<Diagram showing correct nozzle to use>

8. WITHDRAWAL PERIOD

Meat and offal - 8 days

Milk - 5 days.

9. SPECIAL WARNING(S), IF NECESSARY

Not to be used on lambs less than one week old.

Avoid treating very young lambs if the weather is unseasonably hot.

Take care not to apply product in the sheep's eyes.

This precaution should be observed particularly when treating breeds with little wool on their heads.

Prior to application, check the Pour-On gun for correct operation and calibration.

Overdose

Toxic signs in mammals are tremors, hyperexcitability, salivation, abnormal involuntary movements and paralysis. The signs disappear rapidly and the animals recover, generally within 1 week.

There is no specific antidote but symptomatic therapy can be given if considered necessary.

Operator Warnings

Make sure when attaching the Crovect Pour-On gun onto the container that the coiled retainer is secured onto the cap and the applicator. Wear eye protection, protective clothing, rubber gloves and boots when applying the product.

Wear a disposable face-mask when applying as a fan-spray for the prevention of blowfly-strike.

Remove contaminated clothing immediately and wash exposed skin with plenty of clean water. Wash splashes from skin and eyes immediately with plenty of clean water. If irritation persists, seek medical advice immediately.

Do not eat, drink or smoke whilst using the product.

Cypermethrin can be skin sensitizing and eye irritating. In a few instances operators may experience a mild tingling sensation of the skin when using the product. Wash splashes from skin immediately and thoroughly with plenty of clean water. If irritation persists seek medical advice immediately. If product splashes in eyes, flush thoroughly with water for 10 to 15 minutes and seek medical advice immediately.
<Back Label - Diagram of bottle closure>
Attach tube to spouted cap provided. Replace original cap after use.

Environmental Warnings

EXTREMELY DANGEROUS to fish and other aquatic life. Do not contaminate ponds, water-ways or ditches with the product or empty container. As a precautionary measure sheep must be kept away from watercourses for at least one hour following treatment.

10. EXPIRY DATE

Exp:

11. SPECIAL STORAGE CONDITIONS

Pharmaceutical Precautions

Store in tightly closed original container. The product must be used within 3 months of opening the container. Discard unused material. Do not store above 25˚C. Protect from direct sunlight. Store away from food, drink and animal feedstuffs.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

[Distribution category]

FOR ANIMAL TREATMENT ONLY.
FOR EXTERNAL USE ONLY.

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION

POM-VPS products may only be supplied in accordance with a prescription from a Registered Qualified Person (RQP) as follows: (i) a registered veterinary surgeon; (ii) a registered pharmacist, or (iii) a registered suitably qualified person (SQP).

POM-VPS

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

KEEP OUT OF REACH AND SIGHT OF CHILDREN.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Elanco GmbH
Heinz-Lohmann Strasse 4
D-27472 Cuxhaven
Groden
Germany

16. MARKETING AUTHORISATION NUMBER(S)

Vm 52127/5079

17. MANUFACTURER’S BATCH NUMBER

Bn N˚.:

MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO PACKAGE LEAFLET, E.g. Concertina Labels. 0.8L Front/Back

[The guidance contained below is national specific only and should be used in addition to EU QRD template guidance for both the Package Leaflet AND the Outer/Immediate package, available on the EMA website.]

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Elanco GmbH
Heinz-Lohmann Strasse 4
D-27472 Cuxhaven
Groden
Germany

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Crovect 1.25% w/v Pour-On Solution for Sheep

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

Contains 1.25% w/v cypermethrin (tech.)

(cis:trans/ 80:20).

4. PHARMACEUTICAL FORM

Pour-on

5. PACKAGE SIZE

0.8L
2.2L
5L

6. INDICATION(S)

Blowfly Strike: Treatment and Prevention for 6-8 weeks
Ticks: Treatment of tick infestation with a persistent efficacy of 10 weeks
Headflies: Treatment and Control
Biting Lice: Treatment

<Front Label>
10 WEEK TICK CLAIM
Pictogram of Blowfly, Headfly, Tick and Lice

7. CONTRAINDICATIONS

Not to be used on lambs less than one week old.

Avoid treating very young lambs if the weather is unseasonably hot.

Take care not to apply product in the sheep's eyes.

This precaution should be observed particularly when treating breeds with little wool on their heads.

Overdose

Toxic signs in mammals are tremors, hyperexcitability, salivation, abnormal involuntary movements and paralysis. The signs disappear rapidly and the animals recover, generally within 1 week.

There is no specific antidote but symptomatic therapy can be given if considered necessary.

8. ADVERSE REACTIONS

9. TARGET SPECIES

Sheep

10. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

FOR EXTERNAL USE ONLY.

Crovect Pour-On must be applied only with the Crovect Pour-On Gun as the product may have a detrimental effect on certain components of conventional dosing guns. Full operating and maintenance instructions, including details on the use of nozzles, are supplied with each Pour-On Gun. Please read carefully before use. Maintain gun carefully to ensure accurate dosage.
On completion of each treatment cycle of the product and before the dosing gun is stored, immediately expel any remaining product in the dosing gun.
Failure to use the product as directed may result in suboptimal parasite control.

Mothering up
Crovect Pour-On minimises mothering-up problems after treatment. However, care must be taken not to apply the product right to the lamb's tail, as a ewe recognises a young lamb partly by the smell of the tail area.
This product contains a blue dye to aid in the identification of treated animals. The colour is temporary and will fade from the fleece within 48 hours of application.

<Front Label>
Contains a blue dye to aid in the identification of treated sheep
Dosage:
BLOWFLIES
Prevention of blowfly strike
<table>

Bodyweight	Doses	Doses per pack
Up to 25 kg*	20 ml	40
25-40 kg	30 ml	26
Over 40 kg	40 ml	20

*Do not administer to animals of less than 12.5 kg bodyweight.

Apply as a fan-spray to the surface of the fleece on the back and hindquarters of the sheep using the Crovect Pour-On Gun fitted with the FAN-SPRAY NOZZLE. Half the dose should be applied to the shoulders, back and flanks, and half to the rump. The distance between the nozzle and the fleece should be approximately 20 cm. Each dose will require 2 or 3 sweeps to apply.
THE PRODUCT WILL PREVENT BLOWFLY STRIKE ONLY ON AREAS COVERED BY SPRAY. One application will give 6-8 weeks protection. Re-treat as required.

Treatment of blowfly strike
Apply directly to all the affected parts at the rate of 2.5 ml per 100-150 cm2 (roughly the size of a hand). Use the Crovect Pour-On Gun with the T-BAR NOZZLE. Most strikes will require 5-10 ml. Ensure that all affected parts are treated. A single application is sufficient to ensure that larvae are expelled and killed within a few hours.

<Diagram showing product application>

Further Information:
Blowflies are attracted by dirty, damp wool or open wounds. The most common site for fly strike is the rump due to soiling caused by diarrhoea and urine staining. The efficacy of Crovect our-On is reduced in the presence of soiled or dirty wool. Therefore, sheep must be crutched (dagged) regularly and appropriate worm control measures employed.

LICE
5 ml per 20 kg bodyweight up to a maximum of 20 ml.
<table>

Bodyweight	Doses	Doses per pack
Up to 20 kg	5 ml	160
21-40 kg	10 ml	80
41-60 kg	15 ml	53

Over 60 kg	20 ml	40

Using the STRAIGHT NOZZLE on the Crovect Pour-On Gun, apply as a pin stream from the shoulders to the rump along the middle of the backline. Sheep may be treated off-shears or at any time during the year.

<Diagram showing correct nozzle to use>

TICKS
Adult sheep and lambs over 10 kg: 10 ml per 20 kg bodyweight up to a maximum of 40 ml. One treatment will give 10 weeks cover.
Lambs under 10 kg: 5 ml followed 3 weeks later by a 10 ml application.

<table>

Bodyweight	Doses	Doses per pack
Up to 10 kg	5 ml	160
11-20 kg	10 ml	80
21-40 kg	20 ml	40
41-60 kg	30 ml	26
Over 60 kg	40 ml	20

Using the STRAIGHT NOZZLE on the Crovect Pour-On Gun, apply as a pin stream from the crown of the head to the top of the rump.

<Diagram showing correct nozzle to use>

HEADFLIES
5 ml per animal irrespective of size.
Apply to the top of the head between the ears using the Crovect Pour-On Gun with the T-BAR NOZZLE. Take care not to apply product in the sheep's eyes. One application of Crovect Pour-On before the start of the headfly season will give up to 4 weeks protection. Re-treat as required.

<Diagram showing correct nozzle to use>

11. ADVICE ON CORRECT ADMINISTRATION

The product is administered by means of a Crovect Pour-On Gun.

12. WITHDRAWAL PERIOD

Meat and offal - 8 days

Milk - 5 days.

13. SPECIAL STORAGE PRECAUTIONS

Pharmaceutical Precautions

Store in tightly closed original container. The product must be used within 3 months of opening the container. Discard unused material. Do not store above 25˚C. Protect from direct sunlight. Store away from food, drink and animal feedstuffs.

Store

14. SPECIAL WARNING(S)<User Warnings>

Make sure when attaching the Crovect Pour-On gun onto the container that the coiled retainer is secured onto the cap and the applicator. Wear eye protection, protective clothing, rubber gloves and boots when applying the product.

Wear a disposable face-mask when applying as a fan-spray for the prevention of blowfly-strike.

Remove contaminated clothing immediately and wash exposed skin with plenty of clean water. Wash splashes from skin and eyes immediately with plenty of clean water. If irritation persists, seek medical advice immediately.

Do not eat, drink or smoke whilst using the product.

Cypermethrin can be skin sensitizing and eye irritating. In a few instances operators may experience a mild tingling sensation of the skin when using the product. Wash splashes from skin immediately and thoroughly with plenty of clean water. If irritation persists seek medical advice immediately. If product splashes in eyes, flush thoroughly with water for 10 to 15 minutes and seek medical advice immediately.

<Back Label - Diagram of bottle closure>
Attach tube to spouted cap provided. Replace original cap after use.

15. EXPIRY DATE

Use Before:

16. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

17. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

18. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

FOR ANIMAL TREATMENT ONLY.
FOR EXTERNAL USE ONLY.

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION

POM-VPS products may only be supplied in accordance with a prescription from a Registered Qualified Person (RQP) as follows: (i) a registered veterinary surgeon; (ii) a registered pharmacist, or (iii) a registered suitably qualified person (SQP).

POM-VPS

19. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

KEEP OUT OF REACH AND SIGHT OF CHILDREN.

20. MARKETING AUTHORISATION NUMBER(S)

Vm 52127/5079

21. MANUFACTURER'S BATCH NUMBER

Bn N˚.:

22. OTHER INFORMATION

Environmental Warnings

EXTREMELY DANGEROUS to fish and other aquatic life. Do not contaminate ponds, water-ways or ditches with the product or empty container. As a precautionary measure sheep must be kept away from watercourses for at least one hour following treatment.

<Back Label - Diagram of bottle closure>
Attach tube to spouted cap provided. Replace original cap after use.

Approved 29 April 2025
Gavin Hall