

LABELLING AND PACKAGE LEAFLET

A. LABELLING

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Facing page of Booklet Label (bag 20 vaginal delivery systems)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

CIDR OVIS 0.35 g Vaginal Delivery System for Sheep
Progesterone

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Each vaginal delivery system contains: Progesterone 0.35 g.

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

20 vaginal delivery systems

4. ROUTE(S) OF ADMINISTRATION

Vaginal use.

5. WITHDRAWAL PERIOD(S)

Meat and offal: zero days.
Milk: zero hours.

6. BATCH NUMBER

Lot {number}

7. EXPIRY DATE

<EXP {month/year}>

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Label face physically stuck to the bag (20 vaginal delivery systems)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

CIDR OVIS 0.35 g Vaginal Delivery System for Sheep
Progesterone

2. STATEMENT OF ACTIVE SUBSTANCES

Each vaginal delivery system contains: Progesterone 0.35 g.

3. PHARMACEUTICAL FORM

Vaginal delivery system.

4. PACKAGE SIZE

20 vaginal delivery systems

5. TARGET SPECIES

Sheep

6. INDICATION(S)

Read the package leaflet before use.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

8. WITHDRAWAL PERIOD(S)

Meat and offal: zero days.
Milk: zero hours.

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}

11. SPECIAL STORAGE CONDITIONS

Do not store above 30°C

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.
To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Zoetis UK Limited
1st Floor, Birchwood Building
Springfield Drive
Leatherhead
Surrey
KT22 7LP

16. MARKETING AUTHORISATION NUMBER(S)

Vm 42058/4206

17. MANUFACTURER’S BATCH NUMBER

<Batch> {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET FOR:

CIDR OVIS 0.35 g Vaginal Delivery System for Sheep

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Zoetis UK Limited
1st Floor, Birchwood Building
Springfield Drive
Leatherhead
Surrey
KT22 7LP

Manufacturer responsible for batch release:

Zoetis Belgium SA
Rue Laid Burniat 1
1348 Louvain-la-Neuve
Belgium

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

CIDR OVIS 0.35 g Vaginal Delivery System for Sheep
Progesterone

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each vaginal delivery system contains:

Active substance(s):

Progesterone 0.35 g

A "T" shape device consisting of progesterone impregnated silicone rubber elastomer skin moulded over an inert nylon spine.

4. INDICATION(S)

For the induction and synchronisation of oestrus and ovulation in non-cycling ewes during seasonal anoestrus.

For the induction and synchronization of oestrus and ovulation in cycling and in non-cycling ewes for advancing the breeding season.

To be used in combination with eCG (equine chorionic gonadotrophin).

5. CONTRAINDICATIONS

Do not use in pregnant ewes.

Do not use in:

- sexually immature ewes or in females with abnormal or immature genital tracts.

- animals presenting with infectious or non-infectious diseases of the genital tract..

6. ADVERSE REACTIONS

Local irritation and discharge of cloudy/yellow mucus are common and discharge of dark red/brown mucus or mucus with fresh blood is uncommon. However, these signs typically resolve within 2 days of removal of the device without the need for treatment.

The frequency of adverse reactions is defined using the following convention:- very common (more than 1 in 10 animals treated displaying adverse reaction(s))

- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

Alternatively you can report via your national reporting system *{national system details}*.

7. TARGET SPECIES

Sheep (ewes)

8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

Vaginal use.

0.35 g of progesterone (1 device) per animal. The vaginal insert should be left in position for 12 days followed by an injection of eCG (PMSG) administered at device removal. The onset of oestrus occurs within 1-2 days after removal of the insert.

In a study of 11 Lacaune breed ewes, ovulation occurred between 42 and 58 hours following eCG injection, with the majority (73%) ovulating between 50 and 54 hours. In the case that artificial insemination and advanced breeding techniques (e.g. embryo transfer) are applied, the timing of ovulation should be taken into consideration for the selected technique for optimal results.

9. ADVICE ON CORRECT ADMINISTRATION

Administration

A device applicator should be used for administration, following the procedure described below:

1. Ensure that the applicator is clean and dipped in a non-irritant antiseptic solution before use.
2. Wearing sterile disposable plastic gloves, fold the arms of the device and load into the applicator. The arms of the device should protrude slightly from the end of the applicator. Care should be taken to avoid unnecessary or prolonged handling of the product to minimise transfer of the active substance to the operator's gloves.

3. Apply a small quantity of obstetrical lubricant to the end of the loaded applicator.
4. Lift the tail and clean the vulva and perineum.
5. Gently insert the applicator into the vagina, first in a vertical direction and then horizontally until some resistance is encountered.
6. Make sure the removal string is free, press the handle of the applicator and allow the barrel to move back towards the handle. This releases the arms of the device, which will then retain the device in the anterior vagina.
7. With the device correctly positioned, withdraw the applicator, leaving the removal string exiting from the vulva.

The applicator should be cleaned and disinfected before being used on another animal.

Removal

The device may be removed by gently pulling on the tail. On occasions the tail of the device may not be visible from the outside of the animal, in such cases it may be located in the posterior vagina using a gloved finger. Approximately 1 in 10 devices may be lost by an animal. Withdrawal of the device should not require force. If any resistance is encountered a gloved finger should be used to ease removal.

If there is any difficulty in removal from the animal beyond that itemised above veterinary advice must be sought.

The device is intended for single use only.

10. WITHDRAWAL PERIOD(S)

Meat and offal: zero days.

Milk: zero hours.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not store above 30°C.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

12. SPECIAL WARNING(S)

Special precautions for use in animals

The efficacy and safety of the veterinary medicinal product has not been evaluated in ewes which are unwell, which have a BCS < 2 or ≥ 4, which have had complications during previous pregnancies or lambings, or which have lambed within the last 45 days.

Use only according to the benefit/risk assessment by the responsible veterinarian.

Animals in poor condition, whether from illness, inadequate nutrition, or other factors, may respond poorly to treatment.

Do not use in pregnant ewes. The safety of the veterinary medicinal product has not been established during lactation. During lactation use only according to the benefit/risk assessment by the responsible veterinarian.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Progesterone is a potent steroid hormone and may cause adverse effects on the reproductive system in cases of high or prolonged exposure.

As adverse effects on unborn children cannot be ruled out, pregnant women should avoid using this product

The product may cause skin and eye irritation, as well as allergic skin rashes.

Those administering the product should avoid contact with the silicone section; pregnant women should avoid using the product completely.

Wear gloves when administering and disposing of the product; insert the device using the applicator.

Wash hands and exposed skin with soap and water after use.

Do not smoke, eat or drink while handling the product.

Pregnancy:

The safety of the veterinary medicinal product has not been established in pregnant ewes and the use is not recommended during gestation

Interaction with other medicinal products and other forms of interaction:

None known.

Overdose (symptoms, emergency procedures, antidotes):

Not applicable.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED


March 2022

15. OTHER INFORMATION

For animal treatment only.

Pack sizes:

Bag with 20 vaginal delivery systems



Approved: 18 May 2022