PARTICULARS TO APPEAR ON <THE OUTER PACKAGE - Box label no leaflet

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Armitage Pet-Care Protect Flea and Tick Collar for Dogs 15% w/w

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

30 g collar 42g collar

Active ingredient:

3. PHARMACEUTICAL FORM

Collar.

4. PACKAGE SIZE

Collar.

5. TARGET SPECIES

Dogs over 12 weeks of age.

6. INDICATION(S)

Control of fleas and ticks on dogs.

Protection will be provided for up to 300 days against flea infestation and for up to 200 days against tick infestation.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Remove collar from blister and fasten collar around the dog's neck. When fitted to the growing dog, ensure the collar is loosened as the animal grows. In the fully-grown dog, correct fitting will allow two fingers to be inserted between the collar and the neck. Any excess length of collar extending beyond the buckles should be cut off and disposed of with the domestic refuse. The collar does not need to be removed when the animal is being bathed, as insecticidal activity remains unaffected by water. Leave in place until the collar effective life is exhausted.

Replace after 7 months if re-infestation is a risk.

8. WITHDRAWAL PERIOD

Not applicable.

9. SPECIAL WARNING(S), IF NECESSARY

CONTRAINDICATIONS:

Do not use in cats. No other insecticide should be used with the collar or within 7 days of removal. Do not use on nursing bitches or puppies under 3 months of age. REMOVE COLLAR FROM DOG IMMEDIATELY SHOULD ANY SIGNS OF DROWSINESS, LISTLESSNESS OR RESPIRATORY DIFFICULTIES OCCUR SOON AFTER FIXING OF THE COLLAR. Remove collar and consult your veterinary surgeon if any sign of skin irritation or adverse reactions such as salivation or nervous signs occur. Do not allow animals to chew the collar.

OPERATOR WARNINGS:

Wash hands thoroughly after handling the collar.

Dimpylate is an organosphosphorus compound; **do not handle** if under medical advice not to work with ANTICHOLINESTERASES. Do not allow children to play with the collar.

Do not allow children to suck or chew the collar.

Do not sleep with pets wearing collars.

Do not smoke, eat or drink while handling collar.

If you feel unwell after handling the collar, seek medical advice.

GENERAL PRECAUTIONS:

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated with a suitable insecticide and vacuumed regularly. Remove fleas from youg puppies and nursing bitches by careful use of a flea comb. If signs of distress or disease persist or appear consult your veterinary surgeon.

10. EXPIRY DATE

EXP:

11. SPECIAL STORAGE CONDITIONS

Do not store above 25 °C. Keep away from direct heat and sunlight. Do not remove the collar from pack until ready to use.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of empty packaging, excess collar length, used and unused collars in the household refuse.

EXTREMELY DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the collar or empty packaging.

13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]

AVM-GSL

For animal treatment only.

14. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of the of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Virbac S.A.

1ère avenue - 2065 m - L.I.D.

06516 Carros Cedex

France

16. MARKETING AUTHORISATION NUMBER(S)

Vm 05653/4088

17. MANUFACTURER'S BATCH NUMBER

Batch:

PACKAGE LEAFLET FOR:

Armitage Pet-Care Protect Flea and Tick Collar for Dogs 15% w/w

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Virbac S.A.

1ère avenue - 2065 m - L.I.D.

06516 Carros Cedex

France

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Armitage Pet-Care Protect Flea and Tick Collar for Dogs 15% w/w

3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS

30 g collar 42g collar

Active ingredient:

4. INDICATION(S)

Control of fleas and ticks on dogs.

Protection will be provided for up to 300 days against flea infestation and for up to 200 days against tick infestation.

5. CONTRAINDICATIONS

Do not use on cats.

6. ADVERSE REACTIONS

Occasional skin irritation and alopecia.

Remove collar and consult your veterinary surgeon if any sign of skin irritation or adverse reactions such as salivation or nervous signs occur.

7. TARGET SPECIES

Dogs over 12 weeks of age.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Remove collar from blister and fasten collar around the dog's neck. When fitted to the growing dog, ensure the collar is loosened as the animal grows. In the fully-grown dog, correct fitting will allow two fingers to be inserted between the collar and the neck. Any excess length of collar extending beyond the buckles should be cut off and disposed of with the domestic refuse. The collar does not need to be removed when the animal is being bathed, as insecticidal activity remains unaffected by water. Leave in place until the collar effective life is exhausted.

Replace after 7 months if re-infestation is a risk.

9. ADVICE ON CORRECT ADMINISTRATION

Not applicable.

10. WITHDRAWAL PERIOD(S)

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Do not store above 25 °C. Keep away from direct heat and sunlight.

12. SPECIAL WARNING(S)

Special warnings for each target species

Do not allow the animals to chew the collar.

Fleas from pets often infest animals' basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated with a suitable insecticide and vacuumed regularly. Remove fleas from young puppies and nursing bitches by careful use of a flea comb.

Special precautions for use in animals

Do not use on nursing bitches or puppies under 3 months of age.

REMOVE COLLAR FROM DOG IMMEDIATELY SHOULD ANY SIGNS OF DROWSINESS, LISTLESSNESS, HYPERSALIVATION, TREMOR OR RESPIRATORY DIFFICULTIES OCCUR SOON AFTER FIXING OF COLLAR. For external use only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands thoroughly after handling the collar.

Dimpylate is an organosphorus compound; **do not handle** if under medical advice not to work with ANTICHOLINESTERASES.

Do not allow children to play with the collar.

Do not allow children to suck or chew the collar.

Do not sleep with pets wearing collars.

Do not smoke, eat or drink while handling collar.

If you feel unwell after handling the collar, seek medical advice.

Use during pregnancy, lactation or lay

Not for use on nursing or lactating bitches.

Interaction with other medicinal products and other forms of interaction

Do not use any other insecticides while the collar is being worn or within 7 days of removal.

For Animal Treatment Only

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

EXTREMELY DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the collar or empty packaging.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

15. OTHER INFORMATION>

Vm 05653/4088

AVM-GSL

Approved: 08/09/2017