

**MINIMUM PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING WHERE  
THERE IS NO PACKAGE LEAFLET, i.e. Combined label and package leaflet  
HDPE bottle/can**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Pulmovet 250 mg/ml solution for use in drinking water/milk for cattle, pigs, chickens and turkeys

**2. COMPOSITION**

**Active substance:**

Tilmicosin (as tilmicosin phosphate): 250 mg/ml

**Excipients:**

Propyl gallate (E310): 0.2 mg/ml

Disodium edetate: 2.0 mg/ml

Clear yellow to brown solution.

**3. PACKAGE SIZE**

960 ml

5040 ml

**4. TARGET SPECIES**

Cattle (pre-ruminant), pigs, chickens (except hens producing eggs for human consumption) and turkeys.

**5. INDICATIONS FOR USE**

Calves: For the treatment and metaphylaxis of bovine respiratory disease, associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis* and *M. dispar* susceptible to tilmicosin.

Pigs: For the treatment and metaphylaxis of respiratory disease associated with *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *Mycoplasma hyopneumoniae* susceptible to tilmicosin.

Chickens: For the treatment and metaphylaxis of respiratory disease associated with *Mycoplasma gallisepticum* and *M. synoviae* susceptible to tilmicosin.

Turkeys: For the treatment and metaphylaxis of respiratory disease associated with *Mycoplasma gallisepticum* and *M. synoviae* susceptible to tilmicosin.

The presence of the disease in the group/flock must be established before the veterinary medicinal product is used.

## 6. CONTRAINDICATIONS

Do not allow horses and other equines access to drinking water containing tilmicosin. Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

## 7. SPECIAL WARNINGS

### Special warnings:

Important: Must be diluted before administration to animals.

Pigs, chickens and turkeys: Water consumption should be monitored in order to guarantee adequate dosing. In case water consumption does not match quantities for which recommended concentrations were calculated, the concentration of the veterinary medicinal product has to be adapted in a way that the recommended dosage will be taken up by the animals or different medication should be considered.

The uptake of medication by animals can be altered as a consequence of illness. In case of insufficient uptake of water or milk replacer, animals should be treated parenterally using an appropriate injectable veterinary medicinal product.

Repeated use of the veterinary medicinal product should be avoided by improving management practices and thorough cleansing and disinfection.

Use of the veterinary medicinal product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with other macrolides, lincosamides and streptogramin B due to the potential for cross-resistance.

### Special precautions for safe use in the target species:

For oral use only. Contains disodium edetate; do not inject.

Use of the veterinary medicinal product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria. Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water or milk replacer, wear overalls, safety glasses, and impervious gloves. Do not eat, drink or smoke when handling this veterinary medicinal product. Wash hands after use.

In the case of accidental ingestion, wash out mouth immediately with water, seek medical advice and show the package leaflet or the label to the physician. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the veterinary medicinal product if you are allergic to ingredients in the veterinary medicinal product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Pregnancy, lactation and lay:

The safety of the veterinary medicinal product has not been established during pregnancy, lactation or lay. Use only according to the benefit/risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

Tilmicosin may lessen the antibacterial activity of beta-lactam antibiotics.  
Do not use simultaneously with bacteriostatic antimicrobial agents.

Overdose:

No symptoms of overdose, with exception of a slight decrease in the milk consumption, have been seen in calves given twice daily doses 5 times the maximum recommended dose or for twice the maximum recommended duration of treatment.

When pigs are offered drinking water containing 300 or 400 mg tilmicosin/litre (equivalent to 22.5-40 mg tilmicosin/kg body weight or 1.5-2 times the recommended concentration) commonly animals exhibit a reduced water intake. Although this has a self-limiting effect on tilmicosin intake, it could, in extreme circumstances, result in dehydration. This can be corrected by the removal of the medicated drinking water and replacement with fresh unmedicated water.

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 75-100 mg tilmicosin/kg body weight or 5 times the recommended dose) for 5 days; daily treatment with 75 mg tilmicosin/litre (equivalent to the maximum recommended dose) for 10 days resulted in a reduction in faecal consistency.

No symptoms of overdose have been seen in turkeys given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 50-135 mg tilmicosin/kg body weight or 5 times the recommended dose) for 3 days; daily treatment with 75 mg tilmicosin/litre (equivalent to the maximum recommended dose) for 6 days also produced no symptoms of overdose.

## 8. ADVERSE EVENTS

Cattle, pigs, chickens and turkeys:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Decreased drinking
--	--------------------

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system.

## 9. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

For oral use only. The veterinary medicinal product must be diluted in drinking water (pigs, chickens, turkeys) or milk replacer (calves) before administration.

Calves: 12.5 mg tilmicosin/kg body weight (i.e. 1 ml of veterinary medicinal product per 20 kg BW), twice daily for 3-5 consecutive days.

Pigs: 15-20 mg tilmicosin/kg body weight/day (i.e. 6-8 ml of veterinary medicinal product per 100 kg of BW per day), for 5 consecutive days, which may be achieved by the inclusion of 200 mg of tilmicosin per litre (80 ml of veterinary medicinal product per 100 litres).

Chickens: 15-20 mg tilmicosin/kg body weight/day (i.e. 6-8 ml of veterinary medicinal product per 100 kg of BW per day), for 3 consecutive days, which may be achieved by the inclusion of 75 mg of tilmicosin per litre (30 ml of veterinary medicinal product per 100 litres).

Turkeys: 10-27 mg tilmicosin/kg body weight/day (i.e. 4-11 ml of veterinary medicinal product per 100 kg of BW per day), for 3 consecutive days, which may be achieved by the inclusion of 75 mg of tilmicosin per litre (30 ml of veterinary medicinal product per 100 litres).

Based on the recommended dose and the number and weight of animals to be treated, the exact daily amount of the veterinary medicinal product required should be calculated according to the following formula:

$$\text{ml veterinary medicinal product per litre drinking water} = \frac{[\text{ml veterinary medicinal product per kg body weight per day} \times \text{average body weight (kg) of animals to be treated}]}{\text{average daily water intake (l/animal)}}$$

One 960 ml bottle of veterinary medicinal product is sufficient to medicate 1200 litres of drinking water for pigs or 3200 litres of drinking water for chickens or turkeys. One 5040 ml can is sufficient to medicate 6300 litres of drinking water for pigs or 16800 litres of drinking water for chickens or turkeys.

One 960 ml bottle and 5040 ml can of veterinary medicinal product are sufficient to medicate in milk replacer respectively 48 to 80 and 252 to 420 veal calves each of 40 kg body weight depending on the duration of treatment.

## 10. ADVICE ON CORRECT ADMINISTRATION

To ensure a correct dosage, body weight should be determined as accurately as possible.

The required dose should be measured using suitably calibrated measuring equipment. Only sufficient medicated drinking water should be prepared to cover the daily requirements.

The medicated water should be the only source of drinking water for the animals for the entire duration of the treatment period.

Water intake should be monitored at frequent intervals during medication.

After the end of the medication period, the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

Medicated drinking water should be prepared fresh every 24 hours. Medicated milk replacer should be prepared fresh every 6 hours.

The veterinary medicinal product should be diluted with water or milk replacer prior to its use and the concentration of the prediluted solution should not exceed 200 ml of veterinary medicinal product/litre (i.e. 1 in 5). The lowest concentrations of the veterinary medicinal product at which stability can be assured are: 0.3 ml veterinary medicinal product/litre drinking water and 0.7 ml veterinary medicinal product/litre milk replacer.

The intake of medicated water/milk replacer depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of tilmicosin may need to be adjusted accordingly.

## **11. WITHDRAWAL PERIODS**

Withdrawal periods:

Calves: Meat and offal: 42 days.  
Pigs: Meat and offal: 14 days.  
Chickens: Meat and offal: 12 days.  
Turkeys: Meat and offal: 19 days.

Not authorised for use in animals producing milk for human consumption.

Not for use in birds producing eggs for human consumption. Do not use within 2 weeks of the start of the laying period.

## **12. SPECIAL STORAGE PRECAUTIONS**

Store below 25°C. Do not refrigerate or freeze. Protect from frost. Protect from light.

## **13. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via waste water. Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

POM-V ('Veterinary medicinal product subject to prescription')
--

## **15. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

Vm 28365/5006

### **Pack sizes**

- Bottle of 960 ml
- Can of 5040 ml

Not all pack sizes may be marketed.

## 16. DATE ON WHICH THE LABEL WAS LAST REVISED

September 2023

Find more product information by searching for the Product Information Database 'PID' on [www.gov.uk](http://www.gov.uk).

## 17. CONTACT DETAILS

Marketing authorisation holder and contact details to report suspected adverse reactions:

Dopharma Research B.V.

Zalmweg 24

NL-4941 VX Raamsdonksveer

Tel: +31-162-582000

[pharmacovigilance@dopharma.com](mailto:pharmacovigilance@dopharma.com)

Manufacturer responsible for batch release:

Dopharma B.V.

Zalmweg 24

NL-4941 VX Raamsdonksveer

## 18. OTHER INFORMATION

## 19. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

## 20. EXPIRY DATE

Exp. {mm/yyyy}

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Once opened use by...

Shelf life after first opening the container: 3 months

Shelf life after dilution in drinking water according to instructions: 24 hours


Shelf life after dilution in milk replacer according to instructions: 6 hours

## 21. BATCH NUMBER

Lot {number}

## 22. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of the sight and reach of children.



Approved: 14 February 2024