

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**{CARDBOARD BOX}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Furosivet 20 mg tablets for dogs and cats  
Furosemide

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains Furosemide 20 mg

**3. PHARMACEUTICAL FORM**

Tablet

**4. PACKAGE SIZE**

10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 250 or 500 tablets

**5. TARGET SPECIES**

Dogs and cats

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

The cap of the container must be securely engaged after use. If smaller quantities are dispensed from the pack, they must be supplied in a container with a

child-resistant closure. If appropriate containers are not available, the product must be supplied in the original container.

Wear impervious gloves during handling and administration of the product and wash hands thoroughly afterwards

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Store in the original container in order to protect from light and moisture. The silica gel bag should be kept inside the container. Half tablets should be replaced back into the original container and should be given at the next administration.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Millpledge Ltd  
Whinleys Estate  
Clarborough  
Retford  
Notts.  
DN22 9NA  
United Kingdom

**16. MARKETING AUTHORISATION NUMBER**

Vm 04409/4007

**17. MANUFACTURER'S BATCH NUMBER**

Batch><Lot> {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**{CONCERTINA FRONT FACE OF THE CONTAINER LABEL, WHICH IS ALSO  
DUPLICATED AND ATTACHED TO THE CONTAINER }**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Furosivet 20 mg tablets for dogs and cats  
Furosemide

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains Furosemide 20 mg

**3. PHARMACEUTICAL FORM**

Tablet

**4. PACKAGE SIZE**

250 tablets

**5. TARGET SPECIES**

Dogs and cats

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

The cap of the container must be securely engaged after use. If smaller quantities are dispensed from the pack, they must be supplied in a container with a child-resistant closure. If appropriate containers are not available, the product must be supplied in the original container.

Wear impervious gloves during handling and administration of the product and wash hands thoroughly afterwards

#### **10. EXPIRY DATE**

EXP {month/year}

#### **11. SPECIAL STORAGE CONDITIONS**

Store in the original container in order to protect from light and moisture. The silica gel bag should be kept inside the container. Half tablets should be replaced back into the original container and should be given at the next administration.

#### **12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

#### **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

#### **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

#### **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Millpledge Ltd  
Whinleys Estate  
Clarbrough  
Retford  
Notts.  
DN22 9NA  
United Kingdom

**16. MARKETING AUTHORISATION NUMBER**

Vm 04409/4007

**17. MANUFACTURER'S BATCH NUMBER**

Batch><Lot> {number}



**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**{BLISTER}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Furosivet 20 mg tablets for dogs and cats  
Furosemide

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Millpledge

**3. EXPIRY DATE**

EXP {month/year}

**4. BATCH NUMBER**

Batch><Lot> {number}

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:  
Furosivet 20 mg tablets for dogs and cats**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Millpledge Ltd  
Whinleys Estate  
Clarborough  
Retford  
Nottinghamshire  
DN22 9NA  
United Kingdom

Manufacturer responsible for batch release:

Millpledge Europe BVBA  
38 Verrekijker  
8750 Wingene  
Belgium

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Furosivet 20 mg tablets for dogs and cats  
Furosemide

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

Each tablet contains Furosemide 20 mg

White circular, flat faced tablet with bevelled edges. Embossed F20 with break-line on one face and plain on reverse.

The tablets can be divided into two equal parts.

**4. INDICATION(S)**

For the treatment of oedema and ascites, particularly resulting from cardiac insufficiency, renal dysfunction, or of a traumatic origin.

**5. CONTRAINDICATIONS**

Do not use in acute glomerular nephritis.

Do not use in patients that have received excessive doses of cardiac glycosides.

Do not use in combination with other loop diuretics.

Do not use the product in animals suffering from hypovolaemia, hypotension or dehydration.

Do not use in cases of renal failure with anuria.  
Do not use in cases of electrolyte deficiency.  
Do not use in cases of known hypersensitivity to furosemide, sulphonamides or any of the excipients.

## **6. ADVERSE REACTIONS**

Due to the diuretic action of furosemide, there may be haemoconcentration and impairment of the circulation. In cases of prolonged treatment electrolyte deficiency (including hypokalaemia, hyponatraemia) and dehydration may occur.  
If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dogs and cats

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Oral route

The initial dosage is 5 mg furosemide per kg bodyweight per day, given in a single dose or in two divided daily doses. This corresponds to 1 tablet per 4 kg bodyweight per day or ½ tablet per 4 kg bodyweight twice a day. Depending on the severity of the oedema or ascites or in refractory cases, the initial daily dosage may be doubled. For maintenance, daily dosage should be adapted to the lowest effective dose by the veterinarian depending on the clinical response of the dog/cat to the therapy.

## **9. ADVICE ON CORRECT ADMINISTRATION**

## **10. WITHDRAWAL PERIOD(S)**

N/A

## **11. SPECIAL STORAGE PRECAUTIONS**

Store in the original container in order to protect from light and moisture. The silica gel bag should be kept inside the container. Half tablets should be replaced back into the original container and should be given at the next administration.

Do not use this veterinary medicinal product after the expiry date which is stated on the container.

The expiry date refers to the last day of that month.

## **12. SPECIAL WARNING(S)**

Special warnings for each target species:

The patient may increase its water intake to compensate for the diuresis. Consideration should be given to restricting water intake to physiologically normal amounts, if the patient's condition makes such a course appropriate.

Special precautions for use in animals:

Furosemide should be used with caution in case of pre-existing electrolyte and/or water imbalance, impaired hepatic function (may precipitate hepatic coma) and diabetes mellitus. In case of prolonged treatment, hydration status and serum electrolytes should be monitored frequently. Prolonged dosage may on occasions justify potassium supplementation, especially if the product is used in conjunction with cardiac glycosides.

1-2 days before and after commencement of treatment with diuretics and ACE inhibitors renal function and hydration status should be monitored.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Furosemide has possible genotoxic properties and there is evidence of carcinogenicity in mice. Although there is inconclusive evidence relating to these effects in humans, skin contact with or accidental ingestion of the product should be avoided. Wear impervious gloves during handling and administration of the product and wash hands thoroughly afterwards.

For the same reasons, the product is supplied in a container with a child-resistant closure. The cap of the container must be securely engaged after use. If smaller quantities are dispensed from the pack, they must be supplied in a container with a child-resistant closure. If appropriate containers are not available, the product must be supplied in the original container.

Unused tablets or half tablets should be placed back into the container, the child-resistant closure replaced, and the product stored safely, out of the sight and reach of children.

In case of accidental ingestion seek medical attention and show product label and/or pack insert to the doctor.

People with known hypersensitivity to furosemide should avoid contact with the veterinary medicinal product.

Do not handle this product if you know you are sensitive to sulphonamides as hypersensitivity to sulphonamides may lead to hypersensitivity to furosemide.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Pregnancy and lactation:

Laboratory studies have shown evidence of teratogenic effects.

The safety of the veterinary medicinal product in the target species has not been established during pregnancy or lactation.

Use only according to the benefit-risk assessment by the responsible veterinarian. A deleterious effect on lactation is to be expected, particularly if drinking water is restricted. Furosemide passes into milk, but not to a great extent.

Interaction with other medicinal products and other forms of interaction:

Concurrent use with drugs affecting electrolyte balance (corticosteroids, other diuretics, amphotericin B, cardiac glycosides) requires careful monitoring.

Furosemide may increase the risk of sulphonamide allergy.

Furosemide may alter insulin requirements in diabetic animals.

Furosemide may reduce the excretion of NSAIDs.

The dose regimen may need to be modified for long term treatment in combination with ACE inhibitors, depending upon the animal's response to therapy.

Concomitant use with aminoglycosides or cephalosporins may increase the risk of nephrotoxicity.

Overdose (symptoms, emergency procedures, antidotes):

Dehydration and electrolyte depletion may occur. Monitor and correct as necessary.

Dosage higher than that which is recommended, may cause transitory deafness and CNS effects (lethargy, coma, seizures). Cardiovascular side effects (e.g.

hypotension, heart rhythm disorders, collapse) may be observed in weak and old patients following overdose.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

May 2023

**15. OTHER INFORMATION**

Pack sizes:

Bottle containing 250 tablets.

Cardboard box of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 25 or 50 Aluminium-PVC foil/PVC-PVDC blisters of 10 tablets each corresponding to 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 150, 250 or 500 tablets per box.

Not all pack sizes may be marketed.

For animal treatment only. To be supplied only on veterinary prescription.

Keep out of the sight and reach of children.



Approved: 13 May 2024