

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Carton**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Alphafluben 44 mg/ml oral gel for dogs  
Flubendazole

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each ml contains:

**Active substance:**

Flubendazole: 44 mg

**3. PHARMACEUTICAL FORM**

Oral gel

**4. PACKAGE SIZE**

7.5 ml

**5. TARGET SPECIES**

Dogs

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Not applicable.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP: {month/year}  
Once opened use within 90 days.

Once opened, use by.....

**11. SPECIAL STORAGE CONDITIONS**

Store below 25°C.  
Do not refrigerate or freeze.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of waste material in accordance with local requirements.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.  
To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Alpha-Vet Allatgyogyaszati Kft  
Hofherr Albert UTCA 42  
Budapest HU-1194  
Hungary

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 51645/4001

**17. MANUFACTURER’S BATCH NUMBER**

Batch {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**Label**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Alphafluben 44 mg/ml oral gel for dogs  
Flubendazole

**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

Flubendazole 44 mg/ml

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

7.5 ml

**4. ROUTE(S) OF ADMINISTRATION**

For oral use.

**5. WITHDRAWAL PERIOD(S)**

Not applicable.

**6. BATCH NUMBER**

Batch {number}

**7. EXPIRY DATE**

EXP: {month/year}  
Once opened use within 90 days.  
Once opened, use by.....

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:**  
**Alphafluben 44 mg/ml oral gel for dogs**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Alpha-Vet Allatgyogyaszati Kft  
Hofherr Albert UTCA 42  
Budapest HU-1194  
Hungary

Manufacturer responsible for batch release:

Alpha-Vet Kft., Bábolna, Köves János utca 13, H-2943 Bábolna, Hungary

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Alphafluben 44 mg/ml oral gel for dogs  
Flubendazole

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each ml contains:

**Active substance:**

Flubendazole: 44 mg

**Excipients:**

Methyl parahydroxybenzoate (E218) 1.8 mg  
Propyl parahydroxybenzoate 0.2 mg

White or almost white, odourless suspension gel

**4. INDICATION(S)**

Anthelmintic for the treatment of dogs infected with roundworms, hookworms and whipworms

Roundworms: *Toxocara canis*, *Toxascaris leonina*

Hookworms: *Ancylostoma caninum*

Whipworms: *Trichuris vulpis*

**5. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to the active substance(s), to the adjuvant(s) or to any of the excipient(s).

## 6. ADVERSE REACTIONS

Transient vomiting has been observed in dogs very rarely.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon. Alternatively you can report via your national reporting system {national system details}.

## 7. TARGET SPECIES

Dogs

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Recommended dose

22 mg flubendazole per kg bodyweight, one 7.5 ml syringe contains 330 mg flubendazole.

Administration


1 ml gel/2 kg bodyweight, once a day for three consecutive days.  
One syringe is for a dog of up to 15 kg.

Administration route

The gel can be administered as follows:

- the exact dosage should be administered directly onto the tongue of the dog,
- the exact dosage should be mixed into the feed of the dog (recommended in case of aggressive dogs that are difficult to treat).

## 9. ADVICE ON CORRECT ADMINISTRATION



Remove the safety cap (3). Turn the ring (2) counterclockwise until it is at the mark on the dosing piston (1), which corresponds to the body weight of the animal in kg. Give the animal the dose. At the next treatment, add the animal's body weight to the number that the ring (2) was previously set to; then turn the ring to this new mark and administer the appropriate dose.



Example: For a dog of 3 kg body weight the ring for the first treatment is set to the 3 kg mark, of 6 kg body weight for the second and of 9 kg body weight for the third treatment.

#### Recommended treatment

##### Dogs:

- Puppies: at 1-2 weeks of age
- Young dogs (under 12 months of age): every 2-3 months
- Breeding bitches: during oestrous cycle, 10 days before and 10 days after parturition
- Adult dogs: every 3-4 months considering the local regulations
- All dogs: prior to vaccination

To ensure a correct dosage, body weight should be determined as accurately as possible.

#### **10. WITHDRAWAL PERIOD(S)**

Not applicable.

#### **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store below 25°C.

Do not refrigerate or freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Shelf life after first opening the container: 90 days.

#### **12. SPECIAL WARNING(S)**

##### Special warnings for each target species:

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of body weight or misadministration of the product'

The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infection based on its epidemiological features, for each individual animal'

##### Special precautions for use in animals:

Parasite resistance may develop to a particular class of anthelmintics after frequent and repeated administration of that anthelmintic class.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product may cause allergic reactions. People with known hypersensitivity to flubendazole or to the excipients methyl parahydroxybenzoate and propyl parahydroxybenzoate should avoid contact with the veterinary medicinal product.

The product might be mildly irritant to eyes and skin. Direct contact of the product with the skin and eyes must be avoided. In the event of accidental skin or eye contact with the product, rinse the area immediately with plenty of clean water.

Additionally, pregnant women and women of child bearing potential should be careful to avoid accidental exposure.

This veterinary medicinal product may be harmful when ingested, particularly for children. Avoid accidental ingestion of the product. Do not leave a syringe in the sight or reach of children. In order to prevent children from getting access to used syringes, keep the syringe in the original packaging after use. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Additional warnings when administering the product into the feed:

Avoid the access of children to the dog's medicated food. In order to prevent children from getting access to the dog's medicated food, pour it over a part of the feed and wait until the animal has completely consumed the medicated feed, then administer the rest of the feed. Give the treatment out of the sight and reach of children. Any uneaten medicated food must be removed immediately and the bowl washed thoroughly; wash hands when handling the product and cleaning the contaminated food bowl.

Wash hands thoroughly after use.

Pregnancy and lactation:

Laboratory studies in rats have shown evidence of teratogenic and foetotoxic effects at high doses.

Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

Overdose (symptoms, emergency procedures, antidotes):

This veterinary medicinal product has a wide therapeutic margin. Five time overdose does not cause adverse reactions.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

**15. OTHER INFORMATION**

Package size:

7.5 ml gel in a linear low density polyethylene (LLDPE) plastic oral syringe, with polystyrene plunger packaged in carton.

When the container is opened for the first time, the discard date should be calculated using the in-use shelf life specified on this package leaflet and the date recorded in the available space on the carton.

For any information about this veterinary medicinal product, please contact the local representative of the marketing auth

Approved 07 February 2022

