

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE  
POLYETHYLENE BACK PACK CONTAINER**

**1 litre container**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

CLiKZiN 12.5 mg/ml Pour-On Suspension for Sheep.

**2. STATEMENT OF ACTIVE SUBSTANCES**

Dicyclanil 12.5 mg/ml

**3. PACKAGE SIZE**

0.8 litres.

2.2 litres.

5 litres.

**4. TARGET SPECIES**

Sheep.

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Pour on

**7. WITHDRAWAL PERIODS**

Meat and offal: 7 days.

Not authorised for use in animals producing milk for human consumption.

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once opened use within 1 year

Once opened use by...

## **9. SPECIAL STORAGE PRECAUTIONS**

Protect from frost.  
Store in the original container.  
Keep the container tightly closed, away from food, drink and animal feedstuffs.  
Protect from direct sunlight.

## **10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

## **11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

## **13. NAME OF THE MARKETING AUTHORISATION HOLDER**

Elanco Europe Ltd

## **14. MARKETING AUTHORISATION NUMBERS**

Vm 00879/3009

## **15. BATCH NUMBER**

Lot {number}

## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET**  
**1 litre container**

**1. Name of the veterinary medicinal product**

CLiKZiN 12.5 mg/ml Pour-On Suspension for Sheep.

**2. Composition**

Each ml contains:

Active substance:  
Dicyclanil 12.5 mg

**Excipients:**

Quinoline yellow (E104)	0.05 mg
Patent blue V (E131)	0.05 mg
Methyl Parahydroxybenzoate (E218)	1.50 mg
Propyl Parahydroxybenzoate	3.00 mg
Butylated Hydroxytoluene (E321)	0.50 mg

Green coloured pour-on suspension.

**3. Target species**

Sheep.

**4. Indications for use**

Prevention of blowfly strike on sheep due to *Lucilia sericata*.

**5. Contraindications**

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

**6. Special warnings**

Special warnings:

None.

Special precautions for safe use in the target species:

The veterinary medicinal product is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm. Established strikes may require a separate treatment with a knockdown insecticide.

It is recommended that animals with dirty back ends are dagged or crutched prior to application. If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.

Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Redness and irritation may develop after skin or eye contact with the veterinary medicinal product.

Contact with skin and eyes should be avoided.

Personal protective equipment consisting of synthetic rubber gloves and PVC trousers should be worn when handling the veterinary medicinal product.

In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.

In case of eye contact, wash immediately with clean water.

Always wash hands and exposed skin with soap and water after work.

Do not eat, drink or smoke whilst using the veterinary medicinal product.

It is good agricultural practice to minimise handling of sheep after treatment. If you need to handle sheep within 2 months after treatment, wear synthetic rubber gloves and long trousers or coveralls. If sheep are wet wear waterproof trousers.

Special precautions for the protection of the environment:

Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is a **serious** risk to aquatic life if this advice is not followed.

The use of the veterinary medicinal product has harmful effects on dung flies and beetles.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation.

Use only according to the benefit-risk assessment by the responsible veterinarian.

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects.

Interaction with other medicinal products and other forms of interaction:

None known.

### Overdose:

An overdose of up to at least 20 times the recommended dose does not lead to any signs of local or systemic intolerance.  
No antidote is known.

### Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## **7. Adverse events**

Target species: Sheep

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system.

## **8. Dosage for each species, routes and method of administration**

For external use only.

The veterinary medicinal product is applied according to the following recommendations:

<b>Bodyweight (kg)</b>	<b>Dose Volume (ml)</b>
10 - 20	20
21 – 30	24
31 - 50	30
>50	36

(Guide dose volumes correspond to 0.7–2 ml [7.5–25 mg dicyclanil] per kg bodyweight.)

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.


If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

Recommended for treatment in sheep 3 weeks post shearing. Shake the container well before use.

The veterinary medicinal product must be applied with a manual or automatic dosing gun, fitted with a spray nozzle, which guarantees the correct spreading of the



product on the fleece. Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray using a 4 stroke method as shown in the pictogram along the spine of the animal in bands at least 10 cm wide from the middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.



The veterinary medicinal product should be administered before or at the start of predicted fly activity but is also suitable for use during the fly season. The veterinary medicinal product will protect against fly strike for 8 weeks. It is good agricultural practice to check animals regularly for fly strike.

## **9. Advice on correct administration**

Please refer to section on “Special Warnings” for further advice on administration.

## **10. Withdrawal periods**

Meat and offal: 7 days.

Not authorised for use in animals producing milk for human consumption.

## **11. Special storage precautions**

Keep out of the sight and reach of children.

Protect from frost.

Store in the original container.

Keep the container tightly closed, away from food, drink and animal feedstuffs.

Protect from direct sunlight.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the container: 1 year.

## **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater.

The veterinary medicinal product should not enter water courses as dicyclanil is dangerous for fish and aquatic organisms.

Do not contaminate ponds or other waterways with product of empty containers.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.

## **14. Marketing authorisation numbers and pack sizes**

Vm 00879/3009

The pack is composed of a pigmented white opaque polyethylene back pack container with blue polypropylene screw cap, containing 0.8, 2.2 or 5 litres of finished product.

Not all pack sizes may be marketed.

## **15. Date on which the package leaflet was last revised**

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

## **16. Contact details**

Marketing authorisation holder and contact details to report suspected adverse reactions:

Manufacturer responsible for batch release:

Elanco France S.A.S  
26 Rue de la Chapelle  
68330 Huningue  
France

## **17. Other information**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE - COMBINED LABEL AND PACKAGE LEAFLET**

**POLYETHYLENE BACK PACK CONTAINER**

**2.5 and 5 litre pack sizes**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

CLiKZiN 12.5 mg/ml Pour-On Suspension for Sheep.

**2. COMPOSITION**

Each ml contains:

Active substance:  
Dicyclanil 12.5 mg

**Excipients:**

Quinoline yellow (E104)	0.05 mg
Patent blue V (E131)	0.05 mg
Methyl Parahydroxybenzoate (E218)	1.50 mg
Propyl Parahydroxybenzoate	3.00 mg
Butylated Hydroxytoluene (E321)	0.50 mg

Green coloured pour on suspension.

**3. PACKAGE SIZE**

0.8 litres.  
2.2 litres.  
5 litres.

**4. TARGET SPECIES**

Sheep.

**5. INDICATIONS FOR USE**

**Indications for use**

Prevention of blowfly strike on sheep due to *Lucilia sericata*.

## 6. CONTRAINDICATIONS

### Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

## 7. SPECIAL WARNINGS

### Special warnings

#### Special warnings:

None.

#### Special precautions for safe use in the target species:

The veterinary medicinal product is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm. Established strikes may require a separate treatment with a knockdown insecticide.

It is recommended that animals with dirty back ends are dagged or crutched prior to application. If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.

Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Redness and irritation may develop after skin or eye contact with the veterinary medicinal product.

Contact with skin and eyes should be avoided.

Personal protective equipment consisting of synthetic rubber gloves and PVC trousers should be worn when handling the veterinary medicinal product.

In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.

In case of eye contact, wash immediately with clean water.

Always wash hands and exposed skin with soap and water after work.

Do not eat, drink or smoke whilst using the veterinary medicinal product.

It is good agricultural practice to minimise handling of sheep after treatment. If you need to handle sheep within 2 months after treatment, wear synthetic rubber gloves and long trousers or coveralls. If sheep are wet wear waterproof trousers.

#### Special precautions for the protection of the environment:

Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is a **serious** risk to aquatic life if this advice is not followed.

The use of the veterinary medicinal product has harmful effects on dung flies and beetles.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation.

Use only according to the benefit-risk assessment by the responsible veterinarian. Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects.

Interactions with other medicinal products and other forms of interaction:

None known.

Overdose:

An overdose of up to at least 20 times the recommended dose does not lead to any signs of local or systemic intolerance.

No antidote is known.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## **8. ADVERSE EVENTS**

### **Adverse events**

Target species: Sheep.

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system.

## **9. DOSAGE FOR EACH TARGET SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

### **Dosage for each species, routes and method of administration**

For external use only.

The veterinary medicinal product is applied according to the following recommendations:

<b>Bodyweight (kg)</b>	<b>Dose Volume (ml)</b>
10 - 20	20
21 – 30	24
31 - 50	30
>50	36


(Guide dose volumes correspond to 0.7–2 ml [7.5–25 mg dicyclanil] per kg bodyweight.)

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

Recommended for treatment in sheep 3 weeks post shearing. Shake the container well before use.

The veterinary medicinal product must be applied with a manual or automatic dosing gun, fitted with a spray nozzle, which guarantees the correct spreading of the product on the fleece. Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray using a 4 stroke method as shown in the pictogram along the spine of the animal in bands at least 10 cm wide from the middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.



The veterinary medicinal product should be administered before or at the start of predicted fly activity but is also suitable for use during the fly season.

The veterinary medicinal product will protect against fly strike for 8 weeks. It is good agricultural practice to check animals regularly for fly strike.

## 10. ADVICE ON CORRECT ADMINISTRATION

### Advice on correct administration

Please refer to section on “Special Warnings” for further advice on administration.

## **11. WITHDRAWAL PERIODS**

### **Withdrawal periods**

Meat and offal: 7 days.

Not authorised for use in animals producing milk for human consumption.

## **12. SPECIAL STORAGE PRECAUTIONS**

### **Special storage precautions**

Keep out of the sight and reach of children.

Protect from frost.

Store in the original container.

Keep the container tightly closed, away from food, drink and animal feedstuffs.

Protect from direct sunlight.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

## **13. SPECIAL PRECAUTIONS FOR DISPOSAL**

### **Special precautions for disposal**

Medicines should not be disposed of via wastewater.

The veterinary medicinal product should not enter water courses as dicyclanil is dangerous for fish and aquatic organisms.

Do not contaminate ponds or other waterways with product or empty containers.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

### **Classification of veterinary medicinal products**

Veterinary medicinal product subject to prescription.



## 15. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 00879/3009

### Pack sizes

The pack is composed of a pigmented white opaque polyethylene back pack container with blue polypropylene screw cap, containing 0.8, 2.2 or 5 litres of finished product.

Not all pack sizes may be marketed.

## 16. DATE ON WHICH THE LABEL WAS LAST REVISED

### Date on which the label was last revised

Detailed information on this veterinary medicinal product is available in the Union Product Database.

## 17. CONTACT DETAILS

### Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Manufacturer responsible for batch release:

Elanco France S.A.S  
26 Rue de la Chapelle  
68330 Huningue  
France

## 18. OTHER INFORMATION

## 19. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

## 20. EXPIRY DATE

Exp {mm/yyyy}

Shelf life after first opening the container: 1 year.  
Once opened use by...

<b>21. BATCH NUMBER</b>
-------------------------

Lot {number}

Approved 16 May 2023

A handwritten signature in black ink, appearing to read "J. Hunter.", is positioned below the approval date.