

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARDBOARD BOX (24, 60 Syringes) and PLASTIC BUCKET (120 or 144 syringes)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Orbeseal dry cow 2.6 g intramammary suspension

2. STATEMENT OF ACTIVE SUBSTANCES

Each intramammary syringe contains 4 g intramammary suspension containing 2.6 g Bismuth subnitrate, heavy.

3. PACKAGE SIZE

24 intramammary syringes
60 intramammary syringes
120 intramammary syringes
144 intramammary syringes

4. TARGET SPECIES

Cattle (dairy cow at drying-off)

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Intramammary use only.

Dosage:


One intramammary syringe into each udder quarter immediately after the last milking of the lactation (at drying off). Do not massage the teat or udder after infusion.

Administration:

Aseptic intramammary infusion must be employed when administering this veterinary medicinal product.

Under cold conditions the veterinary medicinal product may be warmed to room temperature in a warm environment, to aid syringeability.

See images for correct administration.



7. WITHDRAWAL PERIODS

Withdrawal periods:
Meat and offal: zero days.
Milk: zero hours.

8. EXPIRY DATE

Exp. {month/year}

9. SPECIAL STORAGE PRECAUTIONS

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Zoetis Belgium S.A.

14. MARKETING AUTHORISATION NUMBERS

Vm 60021/3063

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
SYRINGE LABEL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Orbeseal dry cow 2.6 g

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

2.6 g Bismuth subnitrate, heavy

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Orbeseal Dry cow 2.6 g intramammary suspension for cattle

2. Composition

Each 4 g intramammary syringe contains:

Active substance:

Bismuth subnitrate, heavy 2.6 g
(equivalent to Bismuth, heavy 1.858 g)

Excipients:

Liquid paraffin
Aluminium Di Tri stearate
Silica, colloidal anhydrous

Greyish white, smooth, unctuous intramammary suspension.

3. Target species

Cattle (dairy cow at drying-off).

4. Indications for use

The veterinary medicinal product is indicated for the prevention of new intramammary infections throughout the dry period.

The veterinary medicinal product prevents new intramammary infections by producing a physical barrier against the entry of bacteria.

In cows considered likely to be free of subclinical mastitis, the veterinary medicinal product can be used on its own in dry cow management and mastitis control.

5. Contraindications

Do not use alone in cows with subclinical mastitis at drying off.

Do not use in cows with clinical mastitis at drying off.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients
See section "Special warnings"; "Pregnancy" and "Lactation".

6. Special warnings

Special warnings:

Selection of cows for treatment with the veterinary medicinal product should be based on veterinary clinical judgement. Selection criteria may be based on the mastitis and cell count

history of individual cows or recognised tests for the detection of subclinical mastitis or bacteriology sampling.

Special precautions for safe use in the target species:

In cows that may have subclinical mastitis, the veterinary medicinal product may be used following administration of a suitable dry cow antibiotic treatment to the infected quarter. As with all dry cow intramammary treatments, it is good practice to observe dry cows regularly for signs of clinical mastitis. If a sealed quarter develops clinical mastitis the affected quarter should be stripped out manually before appropriate antibiotic therapy is instituted. To reduce the risk of contamination, do not immerse syringes in water.

Use syringe only once.

Since the veterinary medicinal product does not have antimicrobial activity, in order to minimize the risk of acute mastitis due to poor infusion technique and lack of hygiene (see section "Adverse events"), it is crucial to follow the aseptic technique of administration described in section "Dosage for each species, routes and method of administration".

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product may cause skin and eye irritation.

Avoid contact with skin or eyes.

Should skin or eye contact occur, wash the affected area thoroughly with water.

If irritation persists, seek medical advice and show this label to the doctor.

People with known hypersensitivity to bismuth salts should avoid contact with the veterinary medicinal product.

Wash hands after use.

Pregnancy:

The veterinary medicinal product is not absorbed following intramammary infusion.

Can be used during pregnancy. At calving, the seal may be ingested by the calf. Ingestion of the veterinary medicinal product by the calf is safe and produces no adverse effects.

Lactation:

Do not use during lactation. If accidentally used in a lactating cow, a small (up to 2-fold) transient rise in somatic cell count may be observed. In such an event, strip out the seal manually, no additional precautions are necessary.

Interaction with other medicinal products and other forms of interaction:

In clinical trials, the compatibility of the veterinary medicinal product has only been shown with a cloxacillin-containing dry cow preparation.

Do not administer any other intramammary product following the administration of the veterinary medicinal product.

Overdose:

Twice the recommended dose has been administered to cows with no clinical adverse effects.

7. Adverse events

Cattle (dairy cow at drying-off).

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Acute mastitis ¹ .
---	-------------------------------

¹Primarily due to poor infusion technique and lack of hygiene. Please refer to sections “Dosage for each species, routes and method of administration” and “Special warnings” regarding the importance of aseptic technique.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.


8. Dosage for each species, routes and method of administration

Intramammary use only.

Dosage:

One intramammary syringe into each udder quarter immediately after the last milking of the lactation (at drying off). Do not massage the teat or udder after infusion.

Administration:



Care must be taken not to introduce pathogens into the teat. It is essential that strict aseptic techniques are used for the infusion of the veterinary medicinal product as it possesses no antimicrobial activity. Failure to follow these recommendations can lead to serious cases of post-infusion mastitis and even death.

1. All teats need to be thoroughly cleansed and disinfected prior to infusion of the veterinary medicinal product. Ensure sufficient time is allocated to treat each animal and do not combine this with other husbandry activities.
2. Ensure animals are appropriately restrained in hygienic conditions. Keep syringes clean and DO NOT immerse in water.
3. A separate pair of clean disposable gloves should be worn for the treatment of each cow.
4. Start with a visibly clean, dry teat and udder. If teats are obviously dirty then clean off dirt from teats only, with

moistened disposable paper towels and dry thoroughly. Dip teats in a rapid-acting pre-dip, leave for 30 seconds, then wipe each teat completely dry with separate disposable paper towels. Strip fore milk into a strip cup and discard.

5. Thoroughly disinfect the whole surface of the teat with a disposable spirit/alcohol soaked swab. Studies indicate that the most effective means of teat cleaning involves the use of swabs freshly prepared from clean dry cotton wool soaked in surgical spirit (or the equivalent). If this is not available, then the supplied cleaning swabs can be used. Clean the teats furthest away from you first, to avoid contaminating clean teats. **See image I.**
6. Gently scrub each teat end with new individual, disposable, spirit/alcohol swabs, until both teat end and swab are visibly clean.
7. Remove the cap from the intramammary tube, being careful not to touch the nozzle.
8. Grip the teat base firmly between your fingers at the junction with the udder. Turn the teat to a slight angle. Infuse the contents of the syringe into the bottom portion of the teat below where you are pinching the teat avoiding contaminating the teat end. Infuse teats in the opposite order to cleaning i.e. treat the quarters closest to you first. **See image II.** Do not massage the veterinary medicinal product into the udder.
9. Apply a post-milking teat disinfectant and confine the treated cows to a yard where they should stand for at least 30 minutes to allow the teat canal to close. **See image III.**

9. Advice on correct administration

It is important that you read the instructions before using this veterinary medicinal product. Great care should be taken in maintaining cleanliness when administering the veterinary medicinal product in order to reduce the risk of potentially fatal post-infusion mastitis. Full advice on teat cleaning technique prior to tubing is included in the instructions and should be followed.

Under cold conditions the veterinary medicinal product may be warmed to room temperature in a warm environment, to aid syringeability.

10. Withdrawal periods

Meat and offal: zero days.

Milk: zero hours.

11. Special storage precautions

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and label after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater <or household waste>.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

To be completed nationally

14. Marketing authorisation numbers and pack sizes

Vm 60021/3063

Available in cardboard boxes of 24, 60 and plastic bucket of 120 or 144 syringes.
Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Zoetis Belgium S.A.
2nd Floor, Building 10
Cherrywood Business Park
Loughlinstown
Dublin 18
D18 T3Y1
Ireland

Manufacturers responsible for batch release:

Cross Vetpharm Group Ltd,
Dublin 24
IRELAND

Haupt Pharma Latina S.r.l
Strada Statale 156 Dei Monti Lepini Km 47600
Latina
04100
ITALY

<Local representatives < and contact details to report suspected adverse reactions*>:>
To be completed nationally (if needed)*

17. Other information

Most of the seal comes out at the first stripping or suckling after calving, but small amounts may occasionally be seen for a few days as flecks on the filter. The veterinary medicinal product can be differentiated from mastitis by its texture.

After calving, the following steps are recommended for the effective removal of the veterinary medicinal product to minimise residual veterinary medicinal product entering the milking machine. The milking machine should not be used to remove the veterinary medicinal product from the teat.

1. Pinch the teat at the top and strip quarter 10-12 times prior to first milking.
2. Strip foremilk and check for residual veterinary medicinal product for first few milkings.
3. Inspect mastitis filters and milk sock for evidence of residual veterinary medicinal product after every milking.

To be completed nationally

Approved 20 March 2025
Gavin Hall