

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Box containing 1, 2, 3, 4, 6, 8, 12, 24, 30, 60, 90 or 150 pipettes

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

ALFAMED FIPRONIL 50 mg spot-on solution for cats
Fipronil

2. STATEMENT OF ACTIVE SUBSTANCE

One 0.5 ml pipette contains:

Active substance:

Fipronil 50 mg

Excipients:

Butylhydroxyanisole E320 0.1 mg

Butylhydroxytoluene E321 0.05 mg

3. PHARMACEUTICAL FORM

Spot-on solution.

4. PACKAGE SIZE

1, 2, 3, 4, 6, 8, 12, 24, 30, 60, 90 or 150 pipettes

5. TARGET SPECIES

Cats.

6. INDICATIONS

Treatment of flea (*Ctenocephalides* spp.) and tick (*Dermacentor reticulatus*) infestations.

The product has a persistent insecticidal efficacy for up to 5 weeks against fleas (*Ctenocephalides felis*).

The product has a persistent acaricidal efficacy for up to 2 weeks against ticks (*Rhipicephalus sanguineus*, *Ixodes ricinus*, *Dermacentor reticulatus*). If ticks of some species (*Rhipicephalus sanguineus* and *Ixodes ricinus*) are present when the product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) where this has been previously diagnosed by a veterinary surgeon.


7. METHOD AND ROUTE OF ADMINISTRATION

External use only.

Administer by topical application to the skin 1 pipette of 0.5 ml per animal.

Hold the pipette upright. Tap the narrow part of the pipette to ensure that the contents are within the main body of the pipette. Break the snap-off top of the spot-on pipette along the scored line.

Part the pet's coat until its skin is visible. Place the tip of the pipette directly against the bared skin and squeeze gently several times to empty its contents Repeat this procedure at one or two different points along the cat's back, preferably at the base of the head and between the shoulders.



It is important to make sure that the product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment.

Care should be taken to avoid excessive wetting of the hair with the product since this will cause a sticky appearance of hairs at the treatment spot. However, should this occur, it will disappear within 24 hours post application.

White deposits may also be seen at the site for up to 48 hours after application.

8. WITHDRAWAL PERIOD

Not applicable.

9. SPECIAL WARNINGS

User warnings: please read the package leaflet before use.

10. EXPIRY DATE

EXP:{month/year}

11. SPECIAL STORAGE CONDITIONS

Store below 30°C. Store in a dry place. Store in the original package.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Read the package leaflet before use.

13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only.

14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”

Keep out of the reach and sight of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

ALFAMED
13ème rue – LID
06517 Carros Cedex
FRANCE

16. MARKETING AUTHORISATION NUMBER

Vm 17902/4056

17. MANUFACTURER’S BATCH NUMBER

Batch: {number}

MINIMUM PARTICULARS TO APPEAR ON BLISTERS

1 blister

1. NAME OF THE VETERINARY MEDICINAL PRODUCT



ALFAMED FIPRONIL

2. CONTENT BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSE

50 mg

3. ROUTE(S) OF ADMINISTRATION

(reference is made to the cat logo and drop in section 1)

4. EXPIRY DATE

EXP: {month/year}

5. BATCH NUMBER

Batch: {number}

6. MARKETING AUTHORISATION HOLDER

ALFAMED

MINIMUM PARTICULARS TO APPEAR ON PIPETTES

1 pipette

1. NAME OF THE VETERINARY MEDICINAL PRODUCT



ALFAMED FIPRONIL

2. CONTENT BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSE

50 mg

3. ROUTE(S) OF ADMINISTRATION

(reference is made to the cat logo and drop in section 1)

4. EXPIRY DATE

EXP: {month/year}

5. BATCH NUMBER

Batch: {number}

6. MARKETING AUTHORISATION HOLDER

ALFAMED

PACKAGE LEAFLET

ALFAMED FIPRONIL 50 mg spot-on solution for cats

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder and batch release site:

ALFAMED
13ème rue – LID
06517 Carros Cedex
FRANCE

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

ALFAMED FIPRONIL 50 mg spot-on solution for cats
Fipronil

3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS

One 0.5 ml pipette contains:

Active substance:

Fipronil 50 mg

Excipients:

Butylhydroxyanisole E320 0.1 mg

Butylhydroxytoluene E321 0.05 mg

4. INDICATIONS

Treatment of flea (*Ctenocephalides* spp.) and tick (*Dermacentor reticulatus*) infestations.

The product has a persistent insecticidal efficacy for up to 5 weeks against fleas (*Ctenocephalides felis*).

The product has a persistent acaricidal efficacy for up to 2 weeks against ticks (*Rhipicephalus sanguineus*, *Ixodes ricinus*, *Dermacentor reticulatus*). If ticks of some species (*Rhipicephalus sanguineus* and *Ixodes ricinus*) are present when the product is applied, all the ticks may not be killed within the first 48 hours but they may be killed within a week.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) where this has been previously diagnosed by a veterinary surgeon.

5. CONTRAINDICATIONS

In the absence of available data, the product should not be used on kittens less than 2 months old and/or weighing less than 1 kg.

Do not use on sick (systemic diseases, fever...) or convalescent animals.

Do not use in rabbits, as adverse drug reactions and even death could occur.

Do not use in cases of hypersensitivity to the active substance or to any of excipients.

6. ADVERSE REACTIONS

If licking occurs, a brief period of hypersalivation may be observed due mainly to the nature of the carrier.

Among the extremely rare suspected adverse reactions, transient cutaneous reactions at the application site (squamosis, local alopecia, pruritus, erythema) and general pruritus or alopecia have been reported after use. Exceptionally, hypersalivation, reversible neurologic symptoms (hyperesthesia, depression, nervous symptoms) or vomiting have been observed after use.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Cats.

8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

Route of administration and dosage:

External use only.


Administer by topical application to the skin 1 pipette of 0.5 ml per animal.

Method of administration:

Thermoformed pipettes:

Hold the pipette upright. Tap the narrow part of the pipette to ensure that the contents are within the main body of the pipette. Break the snap-off top of the spot-on pipette along the scored line.


Part the pet's coat until its skin is visible. Place the tip of the pipette directly against the bared skin and squeeze gently several times to empty its contents. Repeat this procedure at one or two different points along the cat's back, preferably at the base of the head and between the shoulders.



Polypropylene pipettes:

Remove the pipette from the blister packaging. Hold the pipette in an upright position, twist and pull the cap off. Turn the cap around and place the other end of the cap back on the pipette. Twist the cap to break the seal, then remove the cap from the pipette.

Part the pet's coat until its skin is visible. Place the tip of the pipette directly against the bared skin and squeeze gently several times to empty its contents. Repeat this procedure at one or two points along the cat's back, preferably at the base of the head and between the shoulders.



It is important to make sure that the product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment.

Care should be taken to avoid excessive wetting of the hair with the product since this will cause a sticky appearance of hairs at the treatment spot. However, should this occur, it will disappear within 24 hours post application.

White deposits may also be seen at the site for up to 48 hours after application.

(Note: There will be 2 insert leaflets, one for each type of pipette. However, for practical reasons both are stated on 1 insert leaflet.)

9. ADVICE ON CORRECT ADMINISTRATION

Treatment schedule:

For optimal control of flea and/or tick infestation the treatment schedule can be based on the local epidemiological situation.

In the absence of safety studies, the minimum treatment interval is 4 weeks.

10. WITHDRAWAL PERIOD

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.

Store below 30°C. Store in a dry place. Store in the original package.

Do not use after the expiry date stated on the pipette.

Do not remove from blister until required for use.

12. SPECIAL WARNINGS

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.

The product does not prevent ticks from attaching to the animals. If the animal has been treated prior to exposure to the ticks, the ticks will be killed in the first 24-48 hours after attachment. This will usually be prior to engorgement, minimising but not excluding the risk of transmission of diseases. Once dead, ticks will often drop off the animal, but any remaining ticks may be removed with a gentle pull.

It is advisable to avoid frequent bathing or shampooing because the maintenance of effectiveness of the product in these cases has not been tested.

When used as part of a strategy for the treatment of Flea Allergy Dermatitis, monthly applications to the allergic patient and to other cats in the household are recommended.

For optimum control of flea problems in a multi-pet household, all dogs and cats in the household should be treated with a suitable insecticide.

Special precautions for use in animals

Avoid contact with the animal's eyes. In the case of accidental eye contact, immediately and thoroughly flush the eyes with water.

Do not apply the product on wounds or damaged skin.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product can cause mucous membrane and eye irritation. Therefore, contact between the product and the mouth or eyes should be avoided.

In the case of accidental eye contact, immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the physician.

Avoid contents coming into contact with the fingers. If this occurs, wash hands with soap and water. Wash hands after use.

Do not smoke, drink or eat during application.

People with known hypersensitivity to fipronil or excipients should avoid contact with the veterinary medicinal product.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Other precautions

The alcohol carrier may have adverse effects on painted, varnished or other household surfaces or furnishings.

Use during pregnancy and lactation

Laboratory studies using fipronil have not shown any evidence of teratogenic or embryotoxic effects. Studies have not been carried out with this product in pregnant and lactating queens. Use in pregnancy and lactation only in accordance with professional veterinary advice and a benefit/risk assessment.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

The product should not enter water courses as this may be dangerous for fish or other aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

Pack sizes:

Revised: August 2017
AN. 00407/2017

Thermoformed pipettes: boxes of 1, 2, 3, 4, 6, 8, 12, 24, 30, 60, 90 or 150 pipettes.

Polypropylene pipettes: blister cards or boxes of 1, 2, 3, 4, 6, 8, 12, 24, 30, 60, 90 or 150 pipettes.

Not all pack sizes may be marketed.

A handwritten signature in black ink, consisting of several stylized, overlapping loops and a long, sweeping tail that extends downwards and to the right.

Approved 03 August 2017