

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARDBOARD OR PLASTIC BOX containing vials of lyophilisate and solvent (1 dose presentations)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Nobivac KC nasal drops, lyophilisate and solvent for suspension

2. STATEMENT OF ACTIVE SUBSTANCES

Each dose (0.4 ml) of reconstituted vaccine contains:
Live *Bordetella bronchiseptica* bacteria strain B-C2: $\geq 10^{8.0}$ and $\leq 10^{9.7}$ cfu
Live canine parainfluenza virus strain Cornell: $\geq 10^{3.0}$ and $\leq 10^{5.8}$ TCID₅₀

3. PACKAGE SIZE

5 x 1 dose
25 x 1 dose

4. TARGET SPECIES

Dogs

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Nasal use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}
Once reconstituted use within 1 hour.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.
Do not freeze.
Protect from light.

10. THE WORDS "READ THE PACKAGE LEAFLET BEFORE USE"

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Limited

14. MARKETING AUTHORISATION NUMBERS

Vm 01708/3022

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

VIAL LABEL – Lyophilisate (vial with 1 dose)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Nobivac KC



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Per dose:

B. bronchiseptica: $\geq 10^{8.0}$ cfu

CPi: $\geq 10^{3.0}$ TCID₅₀

1 dose

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Once reconstituted use within 1 hour.

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

VIAL LABEL – Solvent (vial with 1 dose)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Sterile Solvent
Water for injection



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

1 dose

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Nobivac KC nasal drops, lyophilisate and solvent for suspension for dogs

2. Composition

Each dose (0.4 ml) of reconstituted vaccine contains:

Active substances:

Live *Bordetella bronchiseptica* bacteria strain B-C2: $\geq 10^{8.0}$ and $\leq 10^{9.7}$ cfu¹
Live canine parainfluenza virus strain Cornell: $\geq 10^{3.0}$ and $\leq 10^{5.8}$ TCID₅₀²

¹ colony forming units

² Tissue Culture Infective Dose 50%

Lyophilisate: off-white or cream-coloured pellet

Solvent: clear colourless solution

3. Target species

Dogs.

4. Indications for use

Active immunisation of dogs against *Bordetella bronchiseptica* and canine parainfluenza virus for periods of increased risk to reduce clinical signs induced by *B. bronchiseptica* and canine parainfluenza virus and to reduce shedding of canine parainfluenza virus.

Onset of immunity: *Bordetella bronchiseptica*: 72 hours after vaccination;
Canine parainfluenza virus: 3 weeks after vaccination.

Duration of immunity: 1 year.

5. Contraindications

None.

6. Special warnings

Special warnings:

Vaccinate healthy animals only.

Special precautions for safe use in the target species:

Vaccinated dogs may excrete the *Bordetella bronchiseptica* vaccine strain up to 6 weeks and the canine parainfluenza vaccine strain up to a few days following vaccination. During this time, the contact of immunosuppressed and unvaccinated dogs with vaccinated dogs should be avoided.

Immunosuppressive medication may impair the development of active immunity and may increase the chance of adverse effects caused by the live vaccine strains.

Cats, pigs and unvaccinated dogs may react to the vaccine strains with mild and transient respiratory signs. Other animals, like rabbits and small rodents have not been tested.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Immunocompromised persons are advised to avoid contact with the vaccine and vaccinated dogs up to 6 weeks after vaccination.

Disinfect hands and equipment after use.

Pregnancy:

Can be used during pregnancy.

Interaction with other medicinal products and other forms of interaction:

Do not administer in conjunction with other intranasal treatments or during antibiotic treatment.

Safety and efficacy data are available which demonstrate that this vaccine can be administered on the same day, but not mixed, with the live vaccines of the Nobivac series against canine distemper, canine contagious hepatitis caused by canine adenovirus type 1, canine parvovirus disease (based on strain 154) and respiratory disease caused by canine adenovirus type 2, where authorised, and inactivated vaccines of the Nobivac series against canine leptospirosis caused by all or some of the following serovars: *L. interrogans* serogroup Canicola serovar Canicola, *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni, *L. interrogans* serogroup Australis serovar Bratislava, and *L. kirschneri* serogroup Grippotyphosa serovar Bananal/Liangguang.

Safety data are available which demonstrate that this vaccine can be administered on the same day but not mixed with the bivalent puppy vaccine of the Nobivac series that contains canine parvovirus strain 630a. Efficacy of this vaccine after concurrent use has not been tested. Therefore, while safety of concurrent use has been demonstrated, the veterinarian should take this into account when deciding to administer the products at the same time.

In very rare cases a transient acute hypersensitivity reaction may occur when this product is used with other vaccines.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the products mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

In case antibiotics are administered within one week after vaccination, the vaccination should be repeated after the antibiotic treatment is finished.

Overdose:

Particularly in very young puppies, signs of upper respiratory tract disease have been observed after a 10-fold overdose of the vaccine, including ocular and nasal

discharges, pharyngitis, sneezing and coughing. The signs started the day after vaccination and have been seen for up to 4 weeks after vaccination.

Major incompatibilities:

Do not mix with any other veterinary medicinal product, except the solvent recommended for use with the veterinary medicinal product.

7. Adverse events

Dogs:

Very common (>1 animal / 10 animals treated):	Nasal discharge ¹ . Ocular discharge ¹ .
Common (1 to 10 animals / 100 animals reacted):	Sneezing ¹ , cough ¹ .
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Wheezing ¹ . Lethargy. Vomiting. Hypersensitivity reaction, anaphylactic-type (allergic) reaction ² . Immune mediated haemolytic anaemia (low amounts of red blood cells), immune mediated thrombocytopenia (decreased platelet count), immune mediated polyarthritis (inflammation of the joints).

¹ Particularly observed in very young susceptible puppies. Signs are generally mild and transient, but in occasional cases may persist for up to four weeks. In animals, which show more severe signs, appropriate antibiotic treatment may be indicated.

² Such reaction may evolve to a more severe condition, which may be life-threatening. If such reaction occurs, appropriate treatment should be administered without delay.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: <{national system details}>.

8. Dosage for each species, routes and method of administration

Nasal use.

Administer one dose of 0.4 ml per animal.

Reconstitute 1 vial of vaccine with 1 vial of solvent.

Vaccination scheme:

Dogs should be at least 3 weeks of age. When this vaccine is concurrently administered (i.e. not mixed) with another vaccine of the Nobivac series as indicated in section 'Special warnings', dogs should not be younger than the minimum age recommended for the other Nobivac vaccine.

Unvaccinated dogs should receive one dose at least 3 weeks prior to the period of anticipated risk, e.g. temporary kennelling, in order to get protection for both vaccine agents. In order to get protection for *Bordetella bronchiseptica* unvaccinated dogs should receive one dose at least 72 hours prior to the period of anticipated risk (see also section 'Special warnings').

Revaccinate annually.

9. Advice on correct administration

Allow the sterile solvent provided to reach room temperature (15 °C – 25 °C). Aseptically reconstitute the lyophilisate with the solvent. Shake the vial well after addition of the solvent. Withdraw the vaccine into the syringe, remove the needle and administer 0.4 ml directly from the tip of the syringe into one nostril. The reconstituted vaccine is an off-white or yellowish coloured suspension.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.
Store and transport refrigerated (2 °C – 8 °C).
Do not freeze. Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after

Exp. The expiry date refers to the last day of that month.

Shelf life after reconstitution according to directions: 1 hour.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste. Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

Vm 01708/3022

Pack sizes:

- Cardboard or plastic boxes with
- 5 x 1 dose of vaccine and solvent
 - 25 x 1 dose of vaccine and solvent

Not all pack sizes may be marketed.

15. Date on which the package leaflet was last revised

November 2023

Detailed information on this veterinary medicinal product is available in the Union Product Database (<https://medicines.health.europa.eu/veterinary>).

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release and contact details to report suspected adverse reactions:

Manufacturer responsible for batch release:

Intervet International B.V.
Wim de Körverstraat 35
5831 AN Boxmeer
The Netherlands

Local representative and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information



Approved: 03 April 2024