PARTICULARS TO APPEAR ON THE OUTER PACKAGE CARTON

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Flypor 40 mg/ml Pour-on Solution

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Permethrin technical (cis:trans 80:20) 40 mg

3. PACKAGE SIZE

250/ 500 ml

1/ 2/ 2.5/ 3/ 4/ 5 litre(s)

4. TARGET SPECIES

Cattle.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

For external use only.

7. WITHDRAWAL PERIODS

Withdrawal periods:

Meat and offal: 3 days.

Milk: Cattle producing milk for human consumption should be treated immediately after milking to allow as long an interval as possible (at least 6 hours) before the next milking. There is no need to withhold milk from subsequent milkings.

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Do not store above 25 °C.
Store in the original container.
Keep the container tightly closed.
Store away from food, drink and animal feedstuffs.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Elanco GmbH

14. MARKETING AUTHORISATION NUMBERS

Vm 52127/3002

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING WHERE THERE IS NO PACKAGE LEAFLET, i.e. Combined label and package leaflet
BOTTLE/FLEXI-PACK

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Flypor 40 mg/ml Pour-on Solution

2. COMPOSITION

Each ml contains:

Permethrin technical (cis:trans 80:20) 40 mg

Clear, straw-coloured liquid.

3. PACKAGE SIZE

250/ 500 ml

1/ 2/ 2.5/ 3/ 4/ 5 litre(s)

4. TARGET SPECIES

Cattle

5. INDICATIONS FOR USE

Indications for use

For the control of flies and lice, and for the treatment of mange on cattle.

6. CONTRAINDICATIONS

Contraindications

Do not treat calves under one week old.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

7. SPECIAL WARNINGS

Special warnings

As with all insecticides, the degree of fly-killing and repellent activity afforded by the veterinary medicinal product is influenced by prevailing weather conditions and the species of flies involved.

Special precautions for safe use in the target species:

Avoid the eyes when applying to the head.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Personal protective equipment consisting of waterproof apron, impervious gloves and boots should be worn when handling the veterinary medicinal product and recently treated animals.

Avoid contact with the eyes. If splashed in the eye or on the skin, wash with plenty of clean, running water immediately.

Do not eat, drink or smoke whilst using the veterinary medicinal product.

Wash hands and exposed skin before eating, drinking or smoking and after work.

Use in a well-ventilated area. Do not inhale the vapour.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

Care should be taken when applying this veterinary medicinal product, as it may have an adverse effect on certain plastics.

Pregnancy:

The safety of the veterinary medicinal product has not been established during pregnancy.

Use only according to the benefit/risk assessment of the responsible veterinary surgeon.

Interactions with other medicinal products and other forms of interaction:

No data available.

Overdose:

Toxic signs in mammals are tremors, hyperexcitability, salivation, chorea, athetosis and paralysis. The signs disappear rapidly and the animals recover, generally within a week.

There is no specific antidote but symptomatic therapy can be given if considered necessary.

Major incompatibilities:

None known.

8. ADVERSE EVENTS

Adverse events

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system at:

Website: https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine

e-mail: adverse.events@vmd.gov.uk

9. DOSAGE FOR EACH TARGET SPECIES, ROUTES AND METHOD OF ADMINISTRATION

Dosage for each species, routes and method of administration

FOR EXTERNAL USE ONLY.

Dosage: Flies, Lice and Mange:
Apply at the rate of 1ml per 10kg bodyweight (up to a maximum of 40ml).

The veterinary medicinal product should be applied using the dispensing chamber incorporated into the special applicator bottle. The chamber is calibrated up to 40ml.

Directions for using the applicator bottle: Leave cap on bottle neck in position. Remove cap and foil seal from dispensing chamber. Gently squeeze bottle until required amount of product has entered the dispensing chamber. Pour measured liquid along animal's head, neck and back. Replace cap on dispensing chamber after use. **Do not squeeze the bottle while pouring.**

FLIES: Apply approximately one quarter of the calculated dose to the top of the head; the remainder should be applied along the neck and mid-backline. A single dose is normally sufficient to control flies for up to 8 weeks. The first treatment should be carried out when the number of flies starts to increase.
In cases of heavy infestation, the dose rate may be repeated at 4 week intervals.

As with all insecticides, the degree of fly-killing and repellent activity afforded by the veterinary medicinal product is influenced by prevailing weather conditions and the

species of flies involved.

LICE: Apply in the same way as the fly dosage. A single dose is normally sufficient to control all lice. In certain cases of heavy infestation the dose may be repeated after 4 weeks.

MANGE: The calculated dose should be applied directly to all visibly affected areas; any remaining should be poured along the back line in the normal way. The dose should be repeated after 4 weeks.

Bodyweight	Dosage	Doses per pack
Up to 100kg	10ml	100
101 to 200kg	20ml	50
201 to 300kg	30ml	33
Over 300kg	40ml	25

10. ADVICE ON CORRECT ADMINISTRATION

Advice on correct administration

The veterinary medicinal product must be applied only with the Elanco Cattle Pour-On Gun as the veterinary medicinal product may have a detrimental effect on certain components of conventional dosing guns.

11. WITHDRAWAL PERIODS

Withdrawal periods

Meat and offal: 3 days.

Milk: Cattle producing milk for human consumption should be treated immediately after milking to allow as long an interval as possible (at least 6 hours) before the next milking. There is no need to withhold milk from subsequent milkings.

12. SPECIAL STORAGE PRECAUTIONS

Special storage precautions

Keep out of the sight and reach of children.

Do not store above 25 °C.
Store in the original container.
Keep the container tightly closed.

Store away from food, drink and animal feedstuffs.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and container after Exp. The expiry date refers to the last day of that month.

13. SPECIAL PRECAUTIONS FOR DISPOSAL

Special precautions for disposal

Medicines should not be disposed of via wastewater.

This veterinary medicinal product should not enter water courses as permethrin may be dangerous for fish and other aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

15. MARKETING AUTHORISATION NUMBER AND PACK SIZES

Vm 52127/3002

Pack sizes

250/ 500 ml bottle

1/ 2/ 2.5/ 3/ 4/ 5 litres flexi-pack

The 3-litre flexi-pack is supplied in a cardboard carton.

Not all pack sizes may be marketed.

16. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the ‘Product Information Database’ on www.gov.uk.

17. CONTACT DETAILS

Contact details

Marketing authorisation holder and contact details to report suspected adverse reactions:

Elanco GmbH
Heinz-Lohmann Strasse 4
Groden
27472 Cuxhaven
Germany

Tel: +44 3308221732

PV.XXI@elancoah.com

Manufacturer responsible for batch release:

Elanco France S.A.S.
26 Rue de la Chapelle,
Huningue, 68330
France

18. OTHER INFORMATION

Other information

POM-VPS

19. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

20. EXPIRY DATE

Exp {mm/yyyy}

21. BATCH NUMBER

Lot {number}

Gavin Hall

Approved: 28 August 2026