

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

{CARDBOARD BOX}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Pet Shield Worm Screen 96 mg/24 mg Spot-on Solution

**2. STATEMENT OF ACTIVE SUBSTANCES**

**WHAT'S IN THIS PRODUCT?**

Each 1.12 ml pipette contains:

**Active substances:**

Praziquantel	96 mg
Emodepside	24 mg

**3. PACKAGE SIZE**

1 x 1.12 ml  
2 x 1.12 ml  
3 x 1.12 ml  
6 x 1.12 ml

**4. TARGET SPECIES**

LARGE CATS  
> 5 – 8 kg



**5. INDICATIONS**

**WHAT DOES THIS PRODUCT DO?**

Treatment of mixed infections by roundworms and tapeworms.


**6. ROUTES OF ADMINISTRATION**

**IS THIS PRODUCT SUITABLE FOR MY CAT?**

Only use PetShield Worm Screen Spot On solution if your cat is aged over 8 weeks and weighs at least 0.5 kg. Read product leaflet for full guidance and warnings.

Spot on use for external application to the skin.

How to use.



**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

Exp. {mm/yyyy}

**9. SPECIAL STORAGE PRECAUTIONS**

**HOW SHOULD I STORE THE PRODUCT?**

Store in the original packaging in order to protect from moisture.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

KRKA

**14. MARKETING AUTHORISATION NUMBERS**

Vm 01656/5103  
Vm 01656/3103

**15. BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

{BAG}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Pet Shield Worm Screen



> 5 – 8 kg

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

praziquantel/emodepside  
96 mg/24 mg


**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

KRKA



**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

{PIPETTE}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Pet Shield Worm Screen



**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

> 5 – 8 kg

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

KRKA

**PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

**PACKAGE LEAFLET**

**1. Name of the veterinary medicinal product**

Pet Shield Worm Screen 30 mg/7.5 mg Spot-on Solution for Small Cats  
Pet Shield Worm Screen 60 mg/15 mg Spot-on Solution for Medium Cats  
Pet Shield Worm Screen 96 mg/24 mg Spot-on Solution for Large Cats

**2. Composition**

Each unit dose (pipette) contains:

**Active substances:**

Pet Shield worm screen spot-on solution	Pipette (ml)	Praziquantel	Emodepside
Small cats (≥ 0.5 – 2.5 kg)	0.35	30 mg	7.5 mg
Medium cats (> 2.5 – 5 kg)	0.70	60 mg	15 mg
Large cats (> 5 – 8 kg)	1.12	96 mg	24 mg

**Excipients:**

Butylhydroxyanisole (E320) 5.4 mg/ml

Clear, colourless to yellow or to brown solution.

**3. Target species**

Small cats (Equal to/greater than 0.5 – 2.5 kg).

Medium cats (Greater than 2.5 – 5 kg).

Large cats (Greater than 5 – 8 kg).



**4. Indications for use**

For the treatment of mixed parasitic infections in cats caused by roundworms and tapeworms of the following species:

Roundworms (Nematodes):

*Toxocara cati* (mature adult, immature adult, L4 and L3)

*Toxocara cati* (L3 larvae) – treatment of queens during late pregnancy to prevent lactogenic transmission to the offspring

*Toxascaris leonina* (mature adult, immature adult and L4)

*Ancylostoma tubaeforme* (mature adult, immature adult and L4)

Tapeworms (Cestodes):

*Dipylidium caninum* (mature adult and immature adult)

*Taenia taeniaeformis* (adult)

*Echinococcus multilocularis* (adult)

The veterinary medicinal product is only indicated when use against nematodes and cestodes is indicated at the same time.

## **5. Contraindications**

Do not use in kittens under 8 weeks of age or weighing less than 0.5 kg.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

## **6. Special warnings**

Special warnings:

Shampooing or immersion of the animal in water directly after treatment may reduce the efficacy of the veterinary medicinal product. Treated animals therefore should not be bathed until the solution has dried.

In the absence of the risk of co-infection with nematodes and cestodes, a narrow spectrum product should be used.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Special precautions for safe use in the target species:

Apply only to the skin surface and on intact skin. Do not administer orally or parenterally.

Avoid the treated cat or other cats in the household licking the site of application while it is wet.

There is limited experience on the use of the veterinary medicinal product in sick and debilitated animals. Therefore, the product should not be administered to these animals.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can be irritating to the skin and eyes.

Care should be taken to avoid the contents of the pipette coming into contact with the skin, eyes and mouth, including hand-to-mouth and hand-to-eye contact.

Avoid direct contact with application area while it is wet. Keep children away from treated animals during that time.

In case of accidental contact with the skin or eyes, wash off any skin contamination immediately with soap and water.

Rinse the affected eyes thoroughly with clean, fresh water.

People with known hypersensitivity to praziquantel should avoid contact with the veterinary medicinal product.

If skin or eye symptoms persist, or in case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician. Care should be taken not to allow children to have prolonged intensive contact (for example, by sleeping) with treated cats during the first 24 hours after application of the veterinary medicinal product.

Do not smoke, eat or drink during application.

Wash hands after use.

Other precautions:

The solvent in this veterinary medicinal product may stain certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the WOAHA, specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

Pregnancy and lactation:

Can be used during pregnancy and lactation. See section »Dosage for each species, routes and method of administration.«

Interaction with other medicinal products and other forms of interaction:

Emodepside is a substrate for P-glycoprotein. Co-treatment with other drugs that are P-glycoprotein substrates/inhibitors (for example, ivermectin and other antiparasitic macrocyclic lactones, erythromycin, prednisolone and cyclosporine) could give rise to pharmacokinetic drug interactions. The potential clinical consequences of such interactions have not been investigated. If your cat is receiving any medications, please contact your vet to discuss this before applying the product. Similarly, please inform your vet that you are using this product if s/he provides your cat with any medication.

Overdose:

Salivation, vomiting and neurological signs (tremor) were observed occasionally when the veterinary medicinal product was administered at up to 10 times the recommended dose in adult cats and up to 5 times the recommended dose in kittens. These symptoms were thought to occur as a result of the cat licking the application site. The symptoms were completely reversible.

There is no known specific antidote.

Major incompatibilities:

None known.

**7. Adverse events**

Cats:

Very rare	Systemic disorders (anorexia, lethargy)
-----------	-----------------------------------------

(<1 animal / 10,000 animals treated, including isolated reports):	<p>Digestive tract disorders (salivation (drooling), vomiting, diarrhoea)<sup>1</sup></p> <p>Neurological disorders (e.g. ataxia (incoordination), tremor)<sup>1,2,3</sup></p> <p>Behavioural disorders (e.g. hyperactivity, anxiety, vocalisation)</p> <p>Application site disorders (alopecia (hair loss), pruritus (itchiness) and/or inflammation)<sup>3</sup></p>
-------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<sup>1</sup>As a result of the cat licking the application site immediately after treatment.

<sup>2</sup>Mild signs.

<sup>3</sup>Transient signs.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: [adverse.events@vmd.gov.uk](mailto:adverse.events@vmd.gov.uk)

## 8. Dosage for each species, routes and method of administration

Spot-on use for external application to the skin.

### Dosage and Treatment Schedule

The recommended minimum doses are 12 mg praziquantel / kg body weight and 3 mg emodepside / kg body weight, equivalent to 0.14 ml of the veterinary medicinal product / kg body weight.

Body weight of the cat (kg)	Pipette size/volume (ml) to be used	Praziquantel (mg/kg body weight)	Emodepside (mg/kg body weight)
Equal to/greater than 0.5-2.5	0.35	12 – 60	3 – 15
Greater than 2.5-5	0.70	12 – 24	3 – 6
Greater than 5-8	1.12	12 – 19.2	3 – 4.8
Greater than 8	Use an appropriate combination of pipettes.		

To ensure a correct dosage, body weight should be determined as accurately as possible. Underdosing could result in ineffective use and may favour resistance development.


For the treatment of roundworms and tapeworms a single administration per treatment is effective.

For the treatment of queens to prevent transmission of *Toxocara cati* (L3 larvae) through the milk to the offspring, a single administration per treatment approximately seven days prior to expected parturition (birthing) is effective.

The need for and frequency of re-treatments should be based on professional advice and should take into account individual risk factors (the local epidemiological situation and the animal's lifestyle).

## 9. Advice on correct administration

1. Remove the pipette from its packaging. Hold the pipette in an upright position, twist and pull the cap off.
2. Turn the cap around and place the other end of the cap back on the pipette. Push and twist the cap to break the seal, then remove the cap from the pipette.
3. Part the coat on the animal's neck at the base of the skull until the skin is visible.
4. Place the tip of the pipette onto the skin and squeeze the pipette several times to empty its contents completely and directly onto the skin in one spot. Avoid contact between the veterinary medicinal product and your fingers.



Application at the base of the skull will minimise the opportunity for the animal to lick the veterinary medicinal product off.

## 10. Withdrawal periods

Not applicable.

## 11. Special storage precautions

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special temperature storage conditions.

Store in the original packaging (PP pipettes inserted into laminated aluminium bags) in order to protect from moisture.

Do not use this veterinary medicinal product after the expiry date which is stated on the labels and carton after Exp. The expiry date refers to the last day of that month.

## **12. Special precautions for disposal**

Medicines should not be disposed of via wastewater.

This veterinary medicinal product should not enter water courses as emodepside may be dangerous for fish and other aquatic organisms.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## **13. Classification of veterinary medicinal products**

Veterinary medicinal product not subject to prescription.

## **14. Marketing authorisation numbers and pack sizes**

Vm 01656/5102

Vm 01656/3102

Vm 01656/5101

Vm 01656/3101

Vm 01656/5103

Vm 01656/3103

0.35 ml, 0.70 ml and 1.12 ml per pipette; cardboard box containing 1, 2, 3 or 6 pipettes.

Not all pack sizes may be marketed.

## **15. PID LINK (Do not print heading)**

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

## **16. Contact details**

Marketing authorisation holder and manufacturer responsible for batch release:  
KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

Local representatives and contact details to report suspected adverse reactions:

KRKA UK Ltd  
United Kingdom  
Tel: 02071 646 156  
[info.uk@krka.biz](mailto:info.uk@krka.biz)

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**17. Other information**

NFA-VPS

For animal treatment only.

Approved 26 November 2025

*Gavin Hall*