AN: 00206/2014 Revised: July 2014

## <PARTICULARS TO APPEAR ON THE OUTER PACKAGE><PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE>

{ High-density polyethylene 5, 10, 20, 60 or 200 litre cans }

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Bovidip 2% w/v Concentrate for Teat Dip or Spray Solution for cows Active substance: 2% w/v available iodine

## 2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Active ingredient: 2% w/v iodine Emollients: 10% w/v glycerol

#### 3. PHARMACEUTICAL FORM

Concentrate for teat dip or spray solution.

## 4. PACKAGE SIZE

5, 10, 20, 60 or 200 litre cans

#### 5. TARGET SPECIES

(Dairy) cattle.

### 6. INDICATION(S)

Teat disinfection as an aid in the prevention of mastitis in (dairy) cattle.

## 7. METHOD AND ROUTE(S) OF ADMINISTRATION

Prior to milking, wash teats with an udder wash solution and dry with a disposable paper towel. Dilute one part of Bovidip 2% w/v Concentrate for Teat Dip or Spray Solution for cows with three parts of clean water prior to use. Use 5 ml per cow per application of the diluted product immediately after milking. Spraying: Spray the entire surface of the teats. Dipping: Dip the full length of the teats and replenish the dip cup as necessary. Diluted product should be used within one day. Contra-indication: Do not use in cases of known hypersensitivity to iodine.

#### 8. WITHDRAWAL PERIOD

Meat : zero days / milk : zero hours.

#### 9. SPECIAL WARNING(S), IF NECESSARY

Do not mix with other chemicals. In absence of compatibility studies, the product should not be mixed with other medicinal products. Do not use on injured teats; discontinue treatment until lesions have resolved. If signs of disease persist or appear, consult your veterinary surgeon. Allow product to dry before exposing the cows to wet (rainy), cold or windy conditions. Discard any product that becomes contaminated. Always use a clean spray or dip cup. The use of gloves during milking and dipping or

AN: 00206/2014 Revised: July 2014

spraying is recommended to protect the skin and for hygienic milk collection. The product can be used during lactation and pregnancy.

#### User warnings:

For external use only. Care should be taken to avoid eye contact. In case of eye contact, flush the eyes with copious amounts of water and seek medical advice. In case of ingestion, drink large quantities of water and obtain medical attention as soon as possible. When used as a spray, avoid working in a spray mist. Wash hands after use. Persons with iodine allergy should wear gloves and mask.

#### 10. EXPIRY DATE

Batch number and expiry date: see label top of can.

EXP {month/year}

Opened cans should be used within 6 months.

#### 11. SPECIAL STORAGE CONDITIONS

Store upright in the tightly closed original container. Do not store above 25°C. Protect from light. Protect from frost. If the product has frozen, thaw in a warm place and shake well before use. For the larger pack sizes, the product should be rolled sufficiently to mix the solution. Under no circumstances should an attempt be made to shake the 60 or 200 litre packs. The 200 L container should not be returned for re-filling.

# 12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

# 13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only.

## 14. THE WORDS "KEEP OUT OF THE REACH AND SIGHT OF CHILDREN"

Keep out of the reach and sight of children.

## 15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

DeLaval NV, Industriepark-Drongen 10, B-9031 Gent

#### 16. MARKETING AUTHORISATION NUMBER(S)

UK Marketing authorization N°: Vm 17140/4008 - UK authorised veterinary medicinal product AVM-GSL

IE Marketing authorization N°: 10827/003/001 - Licensed Merchant LM

## 17. MANUFACTURER'S BATCH NUMBER

Batch number and expiry date: see label top of can.