

<PARTICULARS TO APPEAR ON THE OUTER PACKAGE>

{NATURE/TYPE}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

PRILBEN® vet 5 mg film-coated tablet for dogs and cats
Benazepril hydrochloride

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Benazepril4.6 mg
(equivalent to Benazepril Hydrochloride 5 mg)

3. PHARMACEUTICAL FORM

Film-coated tablets

4. PACKAGE SIZE

14 tablets
140 tablets

5. TARGET SPECIES

Dogs and cats.

6. INDICATION(S)

For treatment of congestive heart failure in dogs.
For reduction of proteinuria associated with chronic kidney disease in cats.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Not applicable

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP

11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C. Store in a dry place.
Return any halved tablet to the blister pack and use within 1 day. The blister pack should be inserted back into the cardboard box.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only.
To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”

Keep out of the reach and sight of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

CP-Pharma Handelsgesellschaft mbH
Ostlandring 13
31303 Burgdorf
Germany

16. MARKETING AUTHORISATION NUMBER

Vm 20916/4042

17. MANUFACTURER'S BATCH NUMBER

Batch

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

{NATURE/TYPE}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

PRILBEN vet 5 mg film-coated tablet for dogs and cats
Benazepril hydrochloride

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Chemo Ibérica, S.A.

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PACKAGE LEAFLET

PRILBEN® vet 5 mg film-coated tablet for dogs and cats Benazepril hydrochloride

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder

CP-Pharma Handelsgesellschaft mbH
Ostlandring 13
31303 Burgdorf
Germany

Manufacturer for the batch release

LABORATORIUM SANITATIS, S.L.
C/Leonardo da Vinci, 11 (Parque Tecnológico de Álava) Miñano
01510 Álava
SPAIN

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

PRILBEN® vet 5 mg film-coated tablet for dogs and cats
Benazepril hydrochloride

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each divisible tablet contains:

Benazepril4.6 mg
(equivalent to Benazepril Hydrochloride 5 mg)

Excipients :

Titanium dioxide (E171).....1.929 mg
Iron oxide yellow (E172)0.117 mg
Iron oxide red (E172).....0.014 mg
Iron oxide black (E172).....0.004 mg

4. INDICATION(S)

Prilben vet 5 mg belongs to a group of medicines called Angiotensin Converting Enzyme (ACE) inhibitors. It is prescribed by the veterinary surgeon for the treatment of congestive heart failure in dogs and for reduction of proteinuria associated with chronic kidney disease in cats.

5. CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substance benazepril hydrochloride or to any ingredient of the tablets.

Do not use in cases of hypotension (low blood pressure), hypovolemia (low blood volume) or acute renal failure.

Do not use in cases of cardiac output failure due to aortic or pulmonary stenosis.

Do not use in pregnant or lactating dogs or cats because the safety of benazepril hydrochloride has not been established during pregnancy or lactation in these species.

6. ADVERSE REACTIONS

Some dogs with congestive heart failure may exhibit vomiting or fatigue during treatment.

In dogs and cats with chronic kidney disease there may be a moderate increase in levels of creatinine, an indicator of kidney function, in the blood. This is likely due to the effect of the medication in reducing the blood pressure within the kidney and is therefore not necessarily a reason for treatment to be stopped, unless the animal is showing other adverse reactions.

The product may increase food consumption and body weight in cats. Vomiting, poor appetite, dehydration, lethargy and diarrhoea have been reported on rare occasions in cats.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Dogs and cats

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

The product should be given orally once daily, with or without food. The duration of treatment is unlimited.

In dogs the product should be administered orally at a minimum dose of 0.25 mg (range 0.25-0.5) benazepril hydrochloride/kg body weight once daily, according to the following table:

Weight of dog (kg)	Prilben Vet 5 mg Film-Coated Tablets	
	Standard dose	Double dose
> 5 - 10	0.5 tablet	1 tablet
>10 - 20	1 tablet	2 tablets

In dogs with congestive heart failure, the dose may be doubled, still administered once daily, to a minimum dose of 0.5 mg (range 0.5-1.0) benazepril hydrochloride/kg body weight if judged necessary and advised by the veterinary surgeon. Always follow the dosing instructions given by the veterinary surgeon.

In cats the product should be administered orally at a minimum dose of 0.5 mg (range 0.5-1.0) benazepril hydrochloride/kg body weight once daily according to the following table:

Weight of cat (kg)	Prilben Vet 5 mg Film-Coated Tablets
2.5 – 5	0.5 tablet
> 5 - 10	1 tablet

9. ADVICE ON CORRECT ADMINISTRATION

None

10. WITHDRAWAL PERIOD

Not applicable

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children

Do not store above 25°C. Store in a dry place.

Return any halved tablet to the blister pack and use within 1 day. The blister pack should be inserted back into the cardboard box.

Do not use after the expiry date stated on the carton after EXP

12. SPECIAL WARNING(S)

Special warnings for dogs and cats

The efficacy and safety of the product has not been established in dogs and cats below 2.5 kg body weight.

Special precautions for use in animals

In cases of chronic kidney disease, your veterinarian will check the hydration status of your pet before starting therapy, and may recommend that regular blood tests are carried out during therapy in order to monitor plasma creatinine concentrations and blood erythrocyte counts.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

In case of accidental oral ingestion, seek medical advice immediately and show the label or the package leaflet to the physician.

Pregnant women should take special care to avoid accidental oral exposure because ACE inhibitors have been found to affect the unborn child during pregnancy in humans.

Use during pregnancy, lactation

Do not use during pregnancy or lactation. The safety of the product has not been established in breeding, pregnant or lactating dogs or cats.

Interactions

Inform the veterinary surgeon if the animal is taking, or has recently taken, any other medicines.

In dogs with congestive heart failure, the veterinary medicinal product has been given in combination with digoxin, diuretics, pimobendan and anti-arrhythmic products without evidence of associated adverse reactions.

In humans, the combination of ACE inhibitors and NSAIDs (Non-Steroidal Anti-Inflammatory Drugs) can lead to reduced anti-hypertensive efficacy or impaired kidney function. The combination of the product and other anti-hypertensive agents (e.g. calcium channel blockers, β -blockers or diuretics), anaesthetics or sedatives may lead to additive hypotensive effects. Therefore, concurrent use of NSAIDs or other medications with a hypotensive effect should be considered with care.

Your veterinary surgeon may recommend to closely monitor kidney function and for signs of hypotension (lethargy, weakness etc) and treat these if necessary.

Interactions with potassium-preserving diuretics like spironolactone, triamterene or amiloride cannot be ruled out. Your veterinary surgeon may recommend to monitor plasma potassium concentrations when using the product in combination with a potassium-sparing diuretic because of the risk of hyperkalaemia (high blood potassium).

Overdose

Transient reversible hypotension (low blood pressure) may occur in cases of accidental overdose. Therapy should consist of intravenous infusion of warm isotonic saline.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

March 2024

15. OTHER INFORMATION

Benazepril hydrochloride is a prodrug hydrolysed in vivo to its active metabolite, benazeprilat.

Benazeprilat is a highly potent and selective inhibitor of the angiotensin converting enzyme (ACE), thus preventing the conversion of inactive angiotensin I to active angiotensin II and thereby also reducing synthesis of aldosterone. Therefore, it blocks effects mediated by angiotensin II and aldosterone, including vasoconstriction of both arteries and veins, retention of sodium and water by the kidney and remodeling effects (including pathological cardiac hypertrophy and degenerative renal changes).

The product causes long-lasting inhibition of plasma ACE activity in dogs and cats, with more than 95% inhibition at peak effect and significant activity (>80% in dogs and >90% in cats) persisting 24 hours after dosing.

The product reduces the blood pressure and volume load on the heart in dogs with congestive heart failure.


In cats with experimental renal insufficiency, the product normalized the elevated glomerular capillary pressure and reduced the systemic blood pressure. Reduction in glomerular hypertension may retard the progression of kidney disease by inhibition of further damage to the kidneys. In a clinical trial in cats with chronic kidney disease, The product significantly reduced protein loss in the urine; this effect is probably mediated via reduced glomerular hypertension and beneficial effects on the glomerular basement membrane. The product also increased the appetite of the cats, particularly in more advanced cases.

In contrast with other ACE inhibitors, benazeprilat is excreted equally by both biliary and urinary routes in dogs and 85% via the biliary and 15% via the urinary route in cats, and therefore no adjustment of the dose of the product is necessary in the treatment of cases with renal insufficiency.

Box with 14 or 140 tablets.

Not all pack size may be marketed.

For animal treatment only – to be supplied only on veterinary prescription.

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 06 March 2024