

## **LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

{CARDBOARD BOX}

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Scalibor Protectorband 0.76 g medicated collar for small and medium dogs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each collar contains:

Active substance: Deltamethrin 0.76 g

Excipient: Titanium dioxide (E171) 0.285 g

**3. PACKAGE SIZE**

One collar

**4. TARGET SPECIES**

Dogs

**5. INDICATION(S)**

For OTC products

Control of infestations with ticks (*Ixodes Ricinus*, *Rhipicephalus sanguineus*) for 5 to 6 months.

Control of blood sucking by phlebotomine sandflies (*Phlebotomus perniciosus*) for a period of 5 to 6 months.

Anti-feeding effect on adult mosquitoes of the species *Culex pipiens pipiens* for 6 months.

**6. ROUTES OF ADMINISTRATION**

For external use.

**7. WITHDRAWAL PERIODS**

**8. EXPIRY DATE**

EXP {mm/yyyy}

**9. SPECIAL STORAGE PRECAUTIONS**

Keep the foil sachet in the outer carton.  
Store below 25 °C.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Ltd.  
Walton Manor, Walton  
Milton Keynes  
MK7 7AJ

**14. MARKETING AUTHORISATION NUMBERS**

Vm 01708/5078

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater.  
Do not contaminate ponds, waterways or ditches with the product or packaging as this may be dangerous for fish and other aquatic organisms.  
Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

NFA-VPS

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**{SACHET}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Scalibor Protectorband 0.76 g medicated collar



**2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCE(S)**

0.76 g Deltamethrin

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

EXP {mm/yyyy}

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

One collar.

**6. ROUTE(S) OF ADMINISTRATION**

For external use.

**7. WITHDRAWAL PERIOD**

**8. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Scalibor Protectorband 0.76 g medicated collar for small and medium dogs

### 2. COMPOSITION

One white collar of 48 cm length contains:

**Active substance:**

Deltamethrin 0.76 g

**Excipients:**

Titanium dioxide (E171) 0.285 g

White collar of smooth consistency with a plastic buckle at one extremity.

### 3. TARGET SPECIES

Dogs

### 4. INDICATIONS FOR USE

Control of infestations with ticks (*Ixodes ricinus*; *Rhipicephalus sanguineus*) for 5 to 6 months.

Control of blood sucking by phlebotomine sandflies (*Phlebotomus perniciosus*) for a period of 5 to 6 months.

Anti-feeding effect on adult mosquitoes of the species *Culex pipiens pipiens* for 6 months.

### 5. CONTRAINDICATIONS

Do not use in puppies less than 7 weeks of age.

Do not use on dogs with skin lesions.

Do not use in animals known to be hypersensitive to pyrethroids.

Do not use on cats.



DO NOT USE  
ON CATS

## 6. SPECIAL WARNING(S)

For animal treatment only.

### Special warnings:

As the collar exerts its full effect after one week, the collar should be preferably applied 1 week before animals are likely to become exposed to infestation.

In rare cases attachment of ticks can occur while wearing the collar. Under unfavourable conditions the transmission of infectious diseases through ticks or sandflies can therefore not be ruled out entirely.

### Special precautions for safe use in the target species:

In case of skin lesions remove the collar until symptoms have resolved.

The influence of shampooing on the duration of efficacy has not been investigated.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash hands with soap and cold water after fitting the collar.

People with known hypersensitivity to triphenyl phosphate should avoid contact with the veterinary medicinal product and the treated animal. Seek medical advice in case of hypersensitivity reactions.

This veterinary medicinal product contains deltamethrin which may cause transient tingling, itchiness, and blotchy redness on exposed skin.

Avoid letting children, in particular those under 2 years old, touch the collar, play with it or put it into their mouth.

Care should be taken not to allow young children to have prolonged intensive contact, e.g. sleeping with a pet wearing a collar.

Keep the sachet with the collar in the outer carton until use.

### Special precautions for the protection of the environment:

While occasional contact with water does not reduce the effectiveness of the collar, it should always be removed before swimming and bathing the dog to avoid adverse effects on aquatic organisms. Dogs must be prevented from swimming in water for the first five days of wearing the collar.

### Pregnancy:

Can be used during pregnancy.

### Lactation:

Can be used during lactation.

### Interaction with other medicinal products and other forms of interaction:

Do not use with other ectoparasiticides containing organophosphates.

### Overdose:

In the unlikely event of the dog eating the collar the following symptoms may occur: Uncoordinated movements, tremor, drooling of saliva, vomiting, rigidity of the hindquarters.

These symptoms usually subside within 48 hours.

Diazepam can be used for symptomatic treatment if necessary.

Major incompatibilities:

None known.

## 7. ADVERSE EVENTS

Dogs

Rare (1 to 10 animals / 10,000 animals treated):	Localised skin reaction (pruritus/ scratching, erythema/ rash, hair loss) <sup>1</sup> Hypersensitivity reaction <sup>1</sup>
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Behavioural disorders (lethargy, hyperactivity) <sup>2</sup> Digestive tract disorders (vomiting, diarrhoea, hypersalivation) Neurological disorders (ataxia, muscle tremor) <sup>3</sup>

<sup>1</sup> involving the neck or the skin in general, which might indicate a local or general hypersensitivity reaction

<sup>2</sup> often associated with skin irritation

<sup>3</sup> subsides within 48 hours after removal of the collar

If any of these symptoms occur, the collar should be removed. Treatment should be symptomatic as no specific antidote is known.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

The 48cm long collar is to be used on small and medium dogs.


For fastening around the neck.

One collar per dog.

For external use only.

## 9. ADVICE ON CORRECT ADMINISTRATION

Remove the collar from the sealed protective sachet. Adjust the collar around the animal's neck without tightening it too tight. Two fingers side-by-side should fit between the band and the dog's neck. Slide the end through the buckle and cut off any excess length extending beyond 5 cm.



## 10. WITHDRAWAL PERIOD(S)

Not applicable.

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store below 25 °C.

Keep the foil sachet in the outer carton.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and sachet after EXP. The expiry date refers to the last day of that month.

## 12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Do not contaminate ponds, waterways or ditches with the product or packaging as this may be dangerous for fish and other aquatic organisms.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

## 13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

NFA-VPS

Veterinary medicinal product not subject to prescription.

## 14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 01708/5078.

One collar is packed into a sachet made of paper-aluminium-polyethylene or paper-aluminium-polyester-polyethylene and secured in the outer carton.

## 15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED

July 2023

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk).

## 16. CONTACT DETAILS

Marketing authorisation holder:

MSD Animal Health UK Ltd.  
Walton Manor, Walton  
Milton Keynes  
Buckinghamshire  
MK7 7AJ

Manufacturer responsible for batch release:

Intervet International B.V.  
Wim de Korverstraat 35  
5831 AN Boxmeer  
Netherlands

or

Intervet Productions S.A.  
Rue de Lyons  
27460 Igoville  
France

Contact details to report suspected adverse reactions:

MSD Animal Health UK Ltd.  
Tel: +44 (0)1908 685685

## 17. OTHER INFORMATION



Approved: 21 July 2023