

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

ZACTRAN 150 mg/ml solution for injection for cattle, sheep and pigs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One ml contains:

**Active substance:**

Gamithromycin 150 mg

**Excipient(s):**

Monothioglycerol 1 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.

Colourless to pale yellow solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cattle, sheep and pigs.

#### **4.2 Indications for use, specifying the target species**

Cattle:

Treatment and metaphylaxis of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*. The presence of the disease in the herd should be established before metaphylactic use.

Pigs:

Treatment of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Haemophilus parasuis* and *Bordetella bronchiseptica*.

Sheep:

Treatment of infectious pododermatitis (foot rot) associated with virulent *Dichelobacter nodosus* and *Fusobacterium necrophorum* requiring systemic treatment.

### **4.3 Contraindications**

Do not use in cases of hypersensitivity to macrolide antibiotics or to any of the excipients.

Do not use this veterinary medicinal product simultaneously with other macrolides or lincosamides (see section 4.8).

### **4.4 Special warnings for each target species**

Cattle and pigs:  
None.

Sheep:

The efficacy of antimicrobial treatment of foot rot might be reduced by other factors, such as wet environmental conditions, as well as inappropriate farm management. Treatment of foot rot should therefore be undertaken along with other flock management tools, for example providing dry environment. Antibiotic treatment of benign foot rot is not considered appropriate.

### **4.5 Special precautions for use**

#### Special precautions for use in animals

Use of the veterinary medicinal product should be based on susceptibility testing and take into account official and local policies on the use of antimicrobials in farm animals.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to the macrolide class should avoid contact with the veterinary medicinal product. Gamithromycin may cause irritation to eyes and/or skin.

Avoid contact with skin or eyes. If eye exposure occurs, flush eyes immediately with clean water. If skin exposure occurs, wash the affected area immediately with clean water.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands after use.

### **4.6 Adverse reactions (frequency and seriousness)**

During clinical trials transient injection site swellings were observed.

- Visible injection site swellings associated with occasional slight pain may develop very commonly in cattle for one day. The swellings typically resolve within 3 to 14 days but may persist in some animals for up to 35 days after treatment.
- Mild to moderate injection site swelling has been reported commonly in sheep and pigs in clinical trials, with occasional slight pain evident for one day in sheep. These local reactions are transient, and typically resolve within 2 (pigs) to 4 (sheep) days.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- Common (more than 1 but less than 10 animals in 100 animals treated)
- Uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- Rare (more than 1 but less than 10 animals in 10,000 animals treated)
- Very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

Based on laboratory animal data, gamithromycin has not produced any evidence of specific developmental or reproductive effects. The safety of gamithromycin during pregnancy and lactation has not been evaluated in cattle, sheep and pigs. Use only according to the risk/benefit assessment by the responsible veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Cross resistance may occur with other macrolides.

Avoid simultaneous administration of antimicrobials with a similar mode of action such as other macrolides or lincosamides.

#### **4.9 Amounts to be administered and administration route**

A single dose of 6 mg gamithromycin/kg body weight (equivalent to 1 ml/25 kg body weight) into the neck (cattle and pigs) or anterior to the shoulder (sheep). To ensure correct dose, body weight should be determined as accurately as possible to avoid underdosing.

##### Cattle and sheep

Subcutaneous injection. For treatment of cattle over 250 kg and sheep over 125 kg body weight, divide the dose so that no more than 10 ml (cattle) or 5 ml (sheep) are injected at a single site.

##### Pigs

Intramuscular injection. The injection volume should not exceed 5 ml per injection site.

The cap may be safely punctured up to 50 times with a 16G needle and up to 80 times with a 18G needle. For multiple vial entry, an automatic dosing device is recommended to avoid excessive broaching of the stopper.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Clinical studies have demonstrated the wide margin of safety for gamithromycin injection in the target species. In young adult cattle, sheep and pig studies, gamithromycin was administered by injection at 6, 18, and 30 mg/kg (1, 3, and 5 times the recommended dose) and repeated three times at 0, 5 and 10 days (three times the recommended duration of use). Injection site reactions were noted in a dose related manner.

#### 4.11 Withdrawal period(s)

Meat and offal:  
Cattle: 64 days.  
Sheep: 29 days.  
Pigs: 16 days.

Not authorised for use in lactating animals producing milk for human consumption. Do not use in pregnant animals which are intended to produce milk for human consumption within 2 months (cows, heifers) or 1 month (ewes) of expected parturition.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antibacterials for systemic use, macrolides  
ATC vet code: QJ01FA95.

#### 5.1 Pharmacodynamic properties

Gamithromycin is an azalide, 15-membered semisynthetic macrolide class antibiotic with uniquely positioned alkylated nitrogen at 7a-position of the lactone ring. This special chemistry facilitates rapid absorption at physiological pH and a long duration of action at the target tissues, the lung and the skin.

Macrolides in general have both bacteriostatic and bactericidal action mediated through disruption of bacterial protein synthesis. Macrolides inhibit bacterial protein biosynthesis by binding to the 50S ribosomal subunit and by preventing peptide chain elongation. The *in vitro* data show that gamithromycin acts in a bactericidal manner. The broad spectrum antimicrobial activity of gamithromycin includes *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, *Actinobacillus pleuropneumoniae*, *Haemophilus parasuis* and *Bordetella bronchiseptica*, the bacterial pathogens most commonly associated with BRD and SRD, and also *Fusobacterium necrophorum* and *Dichelobacter nodosus*. The MIC and MBC data (cattle and pig) are reported from a representative sample of isolates from field materials within different EU geographic areas.

Cattle	MIC <sub>90s</sub>	MBC <sub>90s</sub>
	µg/ml	
<i>Mannheimia haemolytica</i>	0.5	1
<i>Pasteurella multocida</i>	1	2
<i>Histophilus somni</i>	1	2
Pigs	MIC <sub>90s</sub>	MBC <sub>90s</sub>
	µg/ml	
<i>Actinobacillus pleuropneumoniae</i>	4	4
<i>Pasteurella multocida</i>	1	2
<i>Haemophilus parasuis</i>	0.5	0.5
<i>Bordetella bronchiseptica</i>	2	4
Sheep	MIC	

	µg/ml
<i>Fusobacterium necrophorum</i>	MIC <sub>90</sub> : 32
<i>Dichelobacter nodosus</i>	0.008 – 0.016

Three mechanisms are generally considered responsible for resistance to the macrolide class of compounds. This is often referred to as MLS<sub>B</sub> resistance as it affects macrolides, lincosamides and streptogramins. The mechanisms involve the alteration of the ribosomal target site, the utilization of active efflux mechanism and the production of inactivating enzymes.

## 5.2 Pharmacokinetic particulars

### Cattle

Gamithromycin administered subcutaneously into the neck of cattle at a single dose of 6 mg/kg body weight, resulted in rapid absorption with peak plasma concentrations observed after 30 to 60 min with a long plasma half-life (> 2 days). The bioavailability of the compound was > 98% with no gender differences. The volume of distribution at steady-state was 25 l/kg. Gamithromycin levels in lung reached a maximum in less than 24 hr, with lung-to-plasma ratio of > 264 indicating that gamithromycin was absorbed rapidly into the target tissue for BRD.

*In vitro* plasma protein binding studies determined that the mean concentration of the free active substance was 74%. Biliary excretion of the unchanged drug substance was the major route of elimination.

### Pigs

Gamithromycin administered intramuscularly in pigs at single dose of 6 mg/kg body weight, resulted in rapid absorption with peak plasma concentrations observed after 5 to 15 min, with a long plasma half-life (about 4 days). The bioavailability of gamithromycin was > 92%. The compound is absorbed rapidly into the target tissue for SRD. Accumulation of gamithromycin in the lung has been demonstrated by high and sustained concentrations in the lung and bronchial fluid which far exceed those in blood plasma. The volume of distribution at steady-state was approximately 39 l/kg. *In vitro* plasma protein binding studies determined that the mean concentration of the free active drug was 77%. Biliary excretion of the unchanged drug was the major route of elimination.

### Sheep

Gamithromycin administered subcutaneously into the neck of sheep at a single dose of 6 mg/kg body weight is rapidly absorbed, and maximum plasma concentrations were observed between 15 minutes and 6 hours after dosing (2.30 hours on average) with high absolute bioavailability of 89%. Gamithromycin skin concentrations were much higher than the plasma concentrations resulting in skin/plasma concentration ratios of approximately 21, 58, and 138 at two, five and ten days post-dosing, respectively, demonstrating extensive distribution and accumulation in skin tissue.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Monothioglycerol  
Succinic Acid  
Glycerol Formal

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening the immediate packaging: 28 days.

### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

Type 1 glass vial of 50, 100, 250 or 500 ml with a chlorobutyl rubber stopper, a polypropylene cap and an aluminium crimp seal.

Polypropylene vial of 100, 250 or 500 ml with a chlorobutyl rubber stopper, a polypropylene cap and an aluminium crimp seal.

Cardboard box containing 1 vial of 50, 100, 250 or 500 ml.

The 500 ml vial is for cattle and pigs only.

Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Vetmedica GmbH  
55216 Ingelheim/Rhein  
GERMANY

**8. MARKETING AUTHORISATION NUMBER**

Vm 04491/5061


**9. DATE OF FIRST AUTHORISATION**

24 July 2008

**10. DATE OF REVISION OF THE TEXT**

March 2021

Approved: 01 March 2021

A handwritten signature in black ink, appearing to read 'J. Long', is positioned below the approval date.