

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

FORTEKOR PLUS 1.25 mg/2.5 mg tablets for dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains:

#### **Active substances:**

	pimobendan	benazepril hydrochloride
FORTEKOR PLUS 1.25 mg/2.5 mg tablets	1.25 mg	2.5 mg

#### **Excipients:**

	iron oxide brown (E172)
FORTEKOR PLUS 1.25 mg/2.5 mg tablets	0.5 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablets.

White and light brown oval bilayer tablets with a score line on both sides. The tablets can be divided into equal halves.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs.

#### **4.2 Indications for use, specifying the target species**

For the treatment of congestive heart failure due to atrioventricular valve insufficiency or dilated cardiomyopathy in dogs. FORTEKOR PLUS is a fixed dose combination and should only be used in patients whose clinical signs are successfully controlled by administration of the same doses of the individual components (pimobendan and benazepril hydrochloride) given concurrently.

### 4.3 Contraindications

Do not use in cases of hypertrophic cardiomyopathies or clinical conditions where an augmentation of cardiac output is not possible for functional or anatomical reasons (e.g. aortic or pulmonary stenosis).

Do not use in cases of hypotension, hypovolaemia, hyponatremia or acute renal failure. Do not use during pregnancy and lactation (see section 4.7).

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

### 4.4 Special warnings for each target species

None.

### 4.5 Special precautions for use

#### Special precautions for use in animals

In cases of chronic kidney disease, it is recommended to check the dog's hydration status before starting therapy, and to monitor its plasma creatinine and blood erythrocyte counts during therapy.

As pimobendan is metabolised in the liver, the product should not be administered to dogs with severe hepatic insufficiency.

The efficacy and safety of the product has not been established in dogs below 2.5 kg body weight or under 4 months of age.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

People with known hypersensitivity to pimobendan or benazepril hydrochloride should avoid contact with the veterinary medicinal product.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnant women should take special care to avoid accidental oral exposure because angiotensin converting enzyme (ACE) inhibitors have been found to affect the unborn child during pregnancy in humans.

### 4.6 Adverse reactions (frequency and seriousness)

#### Pimobendan

A moderate positive chronotropic effect and vomiting may occur in rare cases.

However, these effects are dose-dependent and can be avoided by reducing the dose in those cases.

Transient diarrhoea, anorexia or lethargy have been observed in rare cases.

### Benazepril hydrochloride

Transient vomiting, incoordination or signs of fatigue have been reported in dogs very rarely, during post authorization experience. In dogs with chronic kidney disease, benazepril may increase plasma creatinine concentrations at the start of therapy very rarely. A moderate increase in plasma creatinine concentrations following administration of ACE inhibitors is compatible with the reduction in glomerular hypertension induced by these agents, and is therefore not necessarily a reason to stop therapy in the absence of other signs.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

Do not use during pregnancy or lactation.

The safety of the veterinary medicinal product has not been established in breeding, pregnant or lactating dogs.

Laboratory studies in rats and rabbits with pimobendan have shown foetotoxic effects at maternotoxic doses. Laboratory studies in rats and rabbits with pimobendan have not shown any effect on fertility. Laboratory studies in rats have shown that pimobendan is excreted into milk.

Laboratory studies in rats with benazepril have shown foetotoxic effects (foetal urinary tract malformation) at maternally non-toxic doses. It is not known if benazepril is secreted into the milk of lactating bitches.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

In dogs with congestive heart failure, benazepril hydrochloride and pimobendan have been given in combination with digoxin and diuretics without demonstrable adverse interactions.

In pharmacological studies no interaction between the cardiac glycoside ouabain and pimobendan was detected. The pimobendan-induced increase in contractility of the heart is attenuated in the presence of the calcium antagonist verapamil and the  $\beta$ -antagonist propranolol.

In man, the combination of angiotensin converting enzyme (ACE) inhibitors and non-steroidal anti-inflammatory drugs (NSAIDs) can lead to reduced anti-hypertensive efficacy or impaired renal function. Therefore, the concurrent use of FORTEKOR PLUS with NSAIDs or any other medications with a hypotensive effect should be considered carefully before using such combinations.

The combination of FORTEKOR PLUS and other anti-hypertensive agents (e.g. calcium channel blockers,  $\beta$ -blockers or diuretics), anaesthetics or sedatives may lead to additive hypotensive effects. Renal function and signs of hypotension (lethargy, weakness, etc) should be monitored closely and treated as necessary.

Interactions with potassium-sparing diuretics such as spironolactone, triamterene or amiloride cannot be ruled out. It is therefore recommended to monitor plasma potassium levels when using FORTEKOR PLUS in combination with a potassium sparing diuretic because of the risk of hyperkalemia.

#### 4.9 Amounts to be administered and administration route

Oral use.

##### Dose and treatment schedule:

FORTEKOR PLUS is a fixed combination product which should only be used in dogs which require both active substances to be administered concomitantly at this fixed dose.

The recommended dose range for FORTEKOR PLUS is 0.25–0.5 mg pimobendan per kg body weight and 0.5–1 mg benazepril hydrochloride per kg body weight divided into two daily doses. FORTEKOR PLUS tablets should be administered orally, twice daily 12 hours apart (morning and evening) and approximately 1 hour before feeding.

The tablets are breakable along the score line. The table below may be used for guidance.

Body weight (kg) of dog	Strength and number of tablets to be administered	
	<b>FORTEKOR PLUS 1.25 mg/2.5 mg tablets</b>	
	Morning	Evening
2.5 – 5	0.5	0.5
5 – 10	1	1

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

In case of overdose the dog should be treated symptomatically. Transient reversible hypotension may occur in accidental overdose. Therapy should consist of intravenous infusion(s) of warm isotonic saline as required.

#### 4.11 Withdrawal period(s)

Not applicable.

## 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: ACE inhibitors, combinations. ATCvet code: QC09BX90

## 5.1 Pharmacodynamic properties

Benazepril hydrochloride is a prodrug which is hydrolysed *in vivo* to its active metabolite, benazeprilat. Benazeprilat is a highly potent and selective inhibitor of ACE, thus preventing the conversion of inactive angiotensin I to active angiotensin II and thereby also reducing the synthesis of aldosterone. Therefore, benazepril blocks effects mediated by angiotensin II and aldosterone, including the vasoconstriction of arteries and veins, the retention of

sodium and water by the kidney, and remodelling effects (including pathological cardiac hypertrophy and degenerative renal changes).

In dogs with congestive heart failure benazepril hydrochloride reduces the blood pressure and volume load on the heart. Benazepril increased the time to worsening of heart failure, and the time to death, improved clinical condition, reduced cough and improved exercise tolerance in dogs with symptomatic congestive heart failure caused by valvular disease or dilated cardiomyopathy.

Pimobendan, a benzimidazole-pyridazinone derivative, is a non-sympathomimetic, non-glycoside inotropic substance with potent vasodilating properties. It increases the calcium sensitivity of cardiac myofilaments and inhibits phosphodiesterase (type III). It also exhibits a vasodilatory action through the inhibition of phosphodiesterase type III activity.

## 5.2 Pharmacokinetic particulars

### Absorption

Following oral administration of pimobendan alone the absolute bioavailability of the active ingredient is 60–63%. Since this bioavailability is considerably reduced when pimobendan is administered with food or shortly thereafter, it is recommended to treat animals approximately 1 hour before feeding.

After oral administration of benazepril hydrochloride alone, the systemic bioavailability is incomplete (~13%) in dogs due to incomplete absorption (38%) and first pass metabolism. Levels of benazepril decline quickly as the drug is partially metabolised by liver enzymes to benazeprilat. There is no significant difference in the pharmacokinetics of benazeprilat when benazepril hydrochloride is administered to fed or fasted dogs. After the oral administration of FORTEKOR PLUS tablets at twice the recommended dose to dogs, peak levels of both compounds are attained rapidly ( $T_{max}$  0.5 h for benazepril hydrochloride and 0.85 h for pimobendan) with peak concentrations ( $C_{max}$ ) for benazepril hydrochloride of 35.1 ng/ml and 16.5 ng/ml for pimobendan. Peak benazeprilat levels are seen after 1.9 h with peak concentrations ( $C_{max}$ ) of 43.4 ng/ml.

### Distribution

The volume of distribution at steady state is 2.6 l/kg after intravenous administration of pimobendan alone, indicating that pimobendan is distributed readily into the tissues. The mean plasma protein binding *in vitro* is 93%.

Benazeprilat concentrations decline biphasically: the initial fast phase ( $t_{1/2} = 1.7$  h) represents elimination of the free drug, while the terminal phase ( $t_{1/2} = 19$  h) reflects the

release of benazeprilat that was bound to ACE, mainly in the tissues. Benazepril and benazeprilat are extensively bound to plasma proteins (85– 90%), and in the tissues they are found mainly in the lung, liver and kidney.

Repeated administration of benazepril hydrochloride leads to slight bioaccumulation of benazeprilat ( $R = 1.47$ ), steady state being achieved within a few days (4 days).

#### Metabolism

Pimobendan is oxidatively demethylated to its major active metabolite, O-desmethyl pimobendan. Further metabolic pathways are phase II, glucuronides and sulfates. Benazepril hydrochloride is partially metabolised by liver enzymes to the active metabolite benazeprilat.

#### Elimination

The plasma elimination half-life of pimobendan when dosed with FORTEKOR PLUS tablets is 0.5 h, consistent with the high clearance of the compound. The main active metabolite of pimobendan is eliminated with a plasma elimination half-life of 2.6 h. Pimobendan is excreted principally in the faeces and to a lesser extent in the urine.

The plasma elimination half-life of benazepril hydrochloride and benazeprilat, when dosed with FORTEKOR PLUS tablets is 0.36 h and 8.36 h, respectively. Benazeprilat is excreted via the biliary (54%) and urinary (46%) routes in dogs. The clearance of benazeprilat is not affected in dogs with impaired renal function; therefore, no adjustment of the FORTEKOR PLUS dose is required in dogs with renal insufficiency.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Artificial Special Dry Flavour  
Basic butylated methacrylate  
copolymer Copovidone  
Croscarmellose  
sodium Crospovidone  
Dibutyl sebacate  
Hypromellose  
Iron oxide brown (E172)  
Lactose monohydrate  
Magnesium stearate  
Maize starch  
Microcrystalline cellulose  
Polysorbate 80  
Povidone  
Silica, colloidal  
anhydrous Silicon  
dioxide anhydrous  
Sodium lauryl sulfate  
Starch pregelatinised  
Succinic acid  
Sucrose

## **6.2 Major incompatibilities**

Not applicable.

## **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Any remaining half tablets should be discarded after 1 day.

## **6.4 Special precautions for storage**

Store below 25 °C.

Keep the blister in the outer carton in order to protect from moisture.

Any remaining half tablet should be placed back in the opened blister and stored (for a maximum of 1 day) in the original cardboard carton out of the sight and reach of children.

## **6.5 Nature and composition of immediate packaging**

The tablets are packaged in aluminium/aluminium blisters packaged into an outer cardboard box.

### Pack sizes:

FORTEKOR PLUS 1.25 mg/2.5 mg tablets:

Cardboard box containing 30 tablets

Cardboard box containing 60 tablets

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Elanco GmbH  
Heinz-Lohmann Strasse 4  
Groden  
D-27472 Cuxhaven  
Germany

**8. MARKETING AUTHORISATION NUMBER**

Vm 52127/5014

**9. DATE OF FIRST AUTHORISATION**

08 September 2015

**10. DATE OF REVISION OF THE TEXT**

October 2023

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

Approved 13 October 2023

A handwritten signature in black ink, consisting of a stylized initial followed by the name "Hunter.".