

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Capstar 11.4, mg tablets for cats and small dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One tablet contains:

**Active substance:**  
Nitenpyram 11.4 mg

**Excipient(s):**  
For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablet

White to light yellow, round, biconvex tablets, with bevelled edges, imprinted on one side with "RB", on the other side with "CG".

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cats and dogs

#### **4.2 Indications for use, specifying the target species**

Treatment of flea infestations (*C. felis*).

#### **4.3 Contraindications**

None

#### **4.4 Special warnings for each target species**

None

#### **4.5 Special precautions for use**

(i) Special precautions for use in animals

Do not use on animals less than 4 weeks old or weighing less than 1 kg as these groups of animals were not studied.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

None.

Special precautions for the protection of the environment:

Not applicable.

#### 4.6 Adverse reactions (frequency and seriousness)

Cats and Dogs:

Very rare ( $<1$ animal / 10,000 animals treated, including isolated reports):	Excessive chewing, licking and/or grooming <sup>1</sup> , Hyperactivity, Vocalisation <sup>1</sup> Neurological signs (e.g., muscle tremor, ataxia, convulsion) <sup>1</sup> Panting <sup>1</sup> Increased scratching <sup>2</sup>
-----------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<sup>1</sup> Transient

<sup>2</sup> For the first hour following administration; presumably caused by flea response to the veterinary medicinal product.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

#### 4.7 Use during pregnancy, lactation or lay

Pregnancy:

Can be used during pregnancy.

Lactation:

Can be used during lactation.

Laboratory studies in rats and rabbits have produced no evidence of teratogenic or foetotoxic effects and the safety of the product was demonstrated in pregnant and lactating cats and dogs.

#### 4.8 Interaction with other medicinal products and other forms of interaction

None known. No adverse drug reactions were seen in clinical studies when nitenpyram was administered with other veterinary medicinal products including commonly used flea products, anthelmintics, vaccines or antibiotics.

#### **4.9 Amount(s) to be administered and administration route**

Oral Use.

The minimum effective recommended dose is 1 mg/kg, with the following recommendations:

One tablet should be given to cats and small dogs weighing 1 kg to 11.0 kg when a flea infestation is detected. The frequency of treatment depends on the degree of infestation. In the case of a severe flea infestation, it may be necessary to treat the animals every day or every other day, until the flea infestation is controlled. Treatment may be resumed if fleas reappear. No more than one treatment should be given per day.

Tablets should be given orally, with or without food. In order to improve palatability, tablets can be disguised in a small quantity of food immediately prior to administration.

The veterinary medicinal product does not have persistent activity. To prevent re-infestation, a suitable treatment to control immature stages of the flea life cycle is recommended. The veterinary surgeon should establish an appropriate treatment regime.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Nitenpyram is well-tolerated by the target species. Overdoses up to 50 mg/kg in cats and up to 70 mg/kg in dogs were asymptomatic.

Side-effects such as salivation, vomiting, soft stools, seizures, or decreased activity are observed at higher dosages and their seriousness increases as dosages increase. Symptoms disappear quickly and recovery is complete by 24 hours after overdosing because of the rapid elimination of nitenpyram. During 6 months of daily dosing in cats and dogs no clinically significant treatment-related side effects were observed.

#### **4.11 Withdrawal period(s)**

Not applicable.

### **5. PHARMACOLOGICAL PROPERTIES**

**Pharmacotherapeutic group:** Other ectoparasiticides for systemic use,

**ATC Vet Code:** QP53BX02

#### **5.1 Pharmacodynamic properties**

The active ingredient, nitenpyram belongs to the chemical class of neonicotinoids which bind and inhibit insect specific nicotinic acetylcholine receptors. Nitenpyram interferes with nerve transmission and leads to the

death of adult fleas. Nitenpyram does not inhibit acetylcholinesterase.

Effects on fleas (*Ctenocephalides felis*) may be seen as soon as 15-30 minutes after administration of the product to the host animal. This coincides with the first blood meal taken by fleas after sufficient blood levels are reached. Between 95% and 100% efficacy is observed within the first 6 hours and 100% efficacy is reached within 24 hours with no residual activity.

## **5.2 Pharmacokinetic particulars**

Nitenpyram is rapidly and to over 90% absorbed from the gastrointestinal tract of cats and dogs. Feeding does not affect absorption in dogs. Feeding slightly delays T<sub>max</sub> in cats without affecting the other pharmacokinetic properties and without affecting efficacy. The maximum blood concentration is reached after 0.5 to 2 hours in both fasted target species and the elimination half-life is about 4 hours in dogs and 8 hours in cats. More than 90% is eliminated in the urine within 1 day in dogs and 2 days in cats, mainly as the unchanged molecule.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Microcrystalline cellulose  
Maize starch  
Lactose monohydrate  
Anhydrous silica  
Magnesium stearate

### **6.2 Major Incompatibilities**

Not applicable.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

### **6.4 Special precautions for storage**

Do not store above 25°C

### **6.5 Nature and composition of immediate packaging**

Cardboard box with 1 polyamide/aluminium/PVC-aluminium blister. Each blister contains 1 tablet.

Cardboard box with 1 or 10 polyamide/aluminium/PVC-aluminium blisters. Each blister contains 6 tablets.

Not all pack sizes may be marketed.

**6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.  
Any unused product or waste material should be disposed of in accordance with national requirements.

**7. MARKETING AUTHORISATION HOLDER**

Elanco Europe Ltd  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

**8. MARKETING AUTHORISATION NUMBER**

Vm 00879/5023

**9. DATE OF FIRST AUTHORISATION**

03 August 2001

**10. DATE OF REVISION OF THE TEXT**

August 2023

**11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product not subject to prescription.

Approved 29 August 2023

A handwritten signature in black ink, appearing to read "A. Hunter". The signature is stylized and written in a cursive-like font.