

SUMMARY OF PRODUCT CHARACTERISTICS

1. Name of the Veterinary Medicinal Product:

Johnson's Cat Flea Spray, cutaneous spray, solution.

2. Qualitative and Quantitative Composition:

Pyrethrum Extract 25%	0.8% w/w
Piperonyl Butoxide	1.0% w/w

For full list of excipients, see section 6.1.

3. Pharmaceutical form:

Cutaneous spray, solution.
A clear, yellow liquid.

4. Clinical particulars:

4.1 Target species:

Cats and Dogs over 12 weeks of age.

4.2 Indications for use, specifying the target species:

Insecticidal spray to kill fleas on cats and dogs over 12 weeks of age.

4.3 Contra-Indications:

Do not use on animals under 12 weeks of age or on nursing females.

4.4 Special warnings for each target species:

CATS: Spray cats for 3-5 seconds (NO MORE) according to size, then comb coat back into place and towel lightly to remove excess Spray. Do not allow cat to groom until this has been done and coat is dry. Cats should not be allowed to lick the hair of treated dogs.

DOGS & CATS:

HAZARD WARNING: Keep treated animals away from fires and other sources of heat for at least 30 minutes following spraying and until the fur is totally dry.

Remove fleas from young animals and nursing females by careful use of a flea comb.

May cause gastric irritation and nervous signs if ingested.

4.5 Special precautions for use:

4.5 i) Special precautions for use in animals:

For external use only.

4.5 ii) Special precautions to be taken by the person administering the medicinal product to animals:

Extremely flammable. Do not spray on a naked flame or any incandescent material.

Do not smoke when using this product.

Do not breathe spray mist and use in well ventilated area.

Avoid contact with eyes. In case of accidental eye contact, rinse with clean, fresh water. If eye irritation persists, seek medical advice.

Persons who are hypersensitive (allergic) to pyrethrum extract should handle the product with care.

Treated animals should not be allowed to sleep or play with people, particularly children, until the animal is dry.

Keep away from food and drink, including animal feeding stuffs.

Wash hands and exposed skin after use.

4.6 Adverse reactions (frequency and seriousness)

None reported.

4.7 Use during pregnancy, lactation or lay:

Do not use on nursing females.

4.8 Interaction with other medicinal products and other forms of interaction:

None reported.

4.9 Amounts to be administered and administration route:

Directions for Use: Shake can before use. Spray lightly from a distance of about 12" (30cms), all over body, commencing at tail, while brushing against lie of coat to permit spray penetration. Avoid eyes and other sensitive areas. Repeat every 3-4 days if necessary until infestation is cleared. Keep animal under observation for 30 minutes after treatment, then brush and comb to remove dead fleas.

CATS: Use Spray with caution on cats over 12 weeks of age but spray only for 3-5 seconds (NO MORE) according to size. Comb coat back into place and towel lightly to remove excess Spray. DO NOT ALLOW CAT TO GROOM UNTIL THIS HAS BEEN DONE AND COAT IS DRY. Some cats are sensitive to aerosols, in which case use a cat flea powder or cat flea collar.

DOGS: This Spray may be used on dogs and puppies over 12 weeks of age, spraying small dogs for about 6 seconds, medium dogs - 9 seconds, large dogs - 14 seconds.

Fleas from pets often infest animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated with a suitable insecticide and vacuumed regularly. Wash or change pet bedding regularly.

4.10 Overdose (symptoms, emergency procedures, antidotes) if necessary:

Hypersalivation, vomiting, diarrhoea and nervous signs are reported. Tranquilisers may be used to control CNS effects.

4.11 Withdrawal periods:

Not applicable.

5. Pharmacological properties:

ATC Vet code: QP53AC51

5.1 Pharmacodynamic properties:

Pyrethrins/Piperonyl Butoxide

The plant Pyrethrum is a chrysanthemate; the powdered, dried flower heads contain active principles mainly in the form of pyrethrins. Pyrethrins have insecticidal activity producing both knock-down and mortality in a range of insects, and low mammalian toxicity. Common names: Dalmatian insect powder and Persian insect powder.

Piperonyl butoxide acts as a synergist when combined with pyrethrin and prolongs its action.

5.2 Pharmacokinetic particulars:

No other information available.

6. Pharmaceutical particulars:

6.1 List of excipients:

Kerosene Odourless
Isopropyl Alcohol
Propane/Isobutane/Butane 30 Propellant

6.2 Incompatibilities:

None known.

6.3 Shelf-life:

Shelf life of the veterinary medicinal product as packaged for sale: 30 months.

6.4 Special precautions for storage:

Extremely flammable. Pressurised container: Protect from sunlight and do not expose to temperatures exceeding 50°C. Do not pierce or burn even after use. Do not spray on a naked flame or any incandescent material. Keep away from sources of ignition – No smoking.
Do not store above 25°C.

6.5 Nature and composition of immediate packaging:

Pack Size: 150ml

Container: Unlined tinfoil aerosol can.

Closure: Aerosol valve assembly capable of delivering 0.85g of product per second.

6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products, if appropriate:

Harmful to fish and crustaceans. Do not contaminate aquaria and fish bowls with the product.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7.0 Marketing Authorisation holder:

Johnson's Veterinary Products Ltd
5 Reddicap Trading Estate
Sutton Coldfield
West Midlands B75 7DF

8. Marketing Authorisation number:

Vm 01759/5004

9. Date of first Authorisation:

8 October 2001

10. Date of revision of the text:

December 2025

Gavin Hall
Approved: 08 December 2025